

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention
of the grant of the patent:
30.08.2006 Bulletin 2006/35

(21) Application number: **01922524.2**

(22) Date of filing: **21.03.2001**

(51) Int Cl.:
C12N 15/63 ^(2006.01) **C12N 15/85** ^(2006.01)
C12N 15/12 ^(2006.01) **C12N 15/62** ^(2006.01)
C07K 14/705 ^(2006.01) **C07K 14/72** ^(2006.01)
A01H 5/00 ^(2006.01) **C12Q 1/68** ^(2006.01)
G01N 33/68 ^(2006.01)

(86) International application number:
PCT/US2001/009050

(87) International publication number:
WO 2001/070816 (27.09.2001 Gazette 2001/39)

(54) **ECDYSONE RECEPTOR-BASED INDUCIBLE GENE EXPRESSION SYSTEM**

INDUZIERTES GENEXPRESSIONSSYSTEM BASIEREND AUF ECDYSONREZEPTOREN

NOUVEAU SYSTEME D'EXPRESSION GENETIQUE INDUCIBLE FONDE SUR LE RECEPTEUR
A L'ECDYSONE

(84) Designated Contracting States:
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE TR**

(30) Priority: **22.03.2000 US 191355 P**
20.02.2001 US 269799 P

(43) Date of publication of application:
18.12.2002 Bulletin 2002/51

(73) Proprietor: **RheoGene Holdings Inc.**
Norristown, PA 19403 (US)

(72) Inventors:
• **PALLI, Subba, Reddy**
Lansdale, PA 19446 (US)
• **KAPITSKAYA, Marianna Zinovjevna**
North Wales, PA 19454 (US)
• **CRESS, Dean, Ervin**
Sounderton, PA 18964 (US)

(74) Representative: **Harding, Charles Thomas et al**
D Young & Co
120 Holborn
London EC1N 2DY (GB)

(56) References cited:
EP-A- 0 798 378 EP-A- 0 965 644
WO-A-96/27673 WO-A-99/02683
WO-A-99/10510 WO-A-99/58155

- **CHRISTOPHERSON K S ET AL: "ECDYSTEROID-DEPENDENT REGULATION OF GENES IN MAMMALIAN CELLS BY A DROSOPHILA ECDYSONE RECEPTOR AND CHIMERIC TRANSACTIVATORS" PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA, NATIONAL ACADEMY OF SCIENCE. WASHINGTON, US, vol. 89, July 1992 (1992-07), pages 6314-6318, XP002921831 ISSN: 0027-8424**
- **KAKIZAWA TOMOKO ET AL: "Ligand-dependent heterodimerization of thyroid hormone receptor and retinoid X receptor." JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 272, no. 38, 1997, pages 23799-23804, XP002200320 ISSN: 0021-9258**
- **PERERA SRINI C ET AL: "An analysis of ecdysone receptor domains required for heterodimerization with ultraspiracle." ARCHIVES OF INSECT BIOCHEMISTRY AND PHYSIOLOGY, vol. 41, no. 2, 1999, pages 61-70, XP001077866 ISSN: 0739-4462**
- **KOELLE M R ET AL: "THE DROSOPHILA ECR GENE ENCODES AN ECDYSONE RECEPTOR, A NEW MEMBER OF THE STEROID RECEPTOR SUPERFAMILY" CELL, CELL PRESS, CAMBRIDGE, MA, US, vol. 67, no. 1, 4 October 1991 (1991-10-04), pages 59-77, XP001010154 ISSN: 0092-8674**

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

- LEID M ET AL: "PURIFICATION, CLONING, AND RXR IDENTITY OF THE HELA CELL FACTOR WITH WHICH RAR OR TRHETERODIMERIZES TO BIND TARGET SEQUENCES EFFICIENTLY" CELL, CELL PRESS, CAMBRIDGE, NA, US, vol. 68, 24 January 1992 (1992-01-24), pages 377-395, XP002007203 ISSN: 0092-8674
- PERERA SRINI C ET AL: "Studies on two ecdysone receptor isoforms of the spruce budworm, *Choristoneura fumiferana*." MOLECULAR AND CELLULAR ENDOCRINOLOGY, vol. 152, no. 1-2, 25 June 1999 (1999-06-25), pages 73-84, XP002200321 ISSN: 0303-7207

Description**FIELD OF THE INVENTION**

[0001] This invention relates to the field of biotechnology or genetic engineering. Specifically, this invention relates to the field of gene expression. More specifically, this invention relates to a novel ecdysone receptor-based inducible gene expression system and methods of modulating the expression of a gene within a host cell using this inducible gene expression system.

BACKGROUND OF THE INVENTION

[0002] In the field of genetic engineering, precise control of gene expression is a valuable tool for studying, manipulating, and controlling development and other physiological processes. Gene expression is a complex biological process involving a number of specific protein-protein interactions. In order for gene expression to be triggered, such that it produces the RNA necessary as the first step in protein synthesis, a transcriptional activator must be brought into proximity of a promoter that controls gene transcription. Typically, the transcriptional activator itself is associated with a protein that has at least one DNA binding domain that binds to DNA binding sites present in the promoter regions of genes. Thus, for gene expression to occur, a protein comprising a DNA binding domain and a transactivation domain located at an appropriate distance from the DNA binding domain must be brought into the correct position in the promoter region of the gene.

[0003] The traditional transgenic approach utilizes a cell-type specific promoter to drive the expression of a designed transgene. A DNA construct containing the transgene is first incorporated into a host genome. When triggered by a transcriptional activator, expression of the transgene occurs in a given cell type.

[0004] Another means to regulate expression of foreign genes in cells is through inducible promoters. Examples of the use of such inducible promoters include the **PR1-a** promoter, prokaryotic repressor-operator systems, immunosuppressive-immunophilin systems, and higher eukaryotic transcription activation systems such as steroid hormone receptor systems and are described below.

[0005] The PR1-a promoter from tobacco is induced during the systemic acquired resistance response following pathogen attack. The use of PR1-a may be limited because it often responds to endogenous materials and external factors such as pathogens, UV-B radiation, and pollutants. Gene regulation systems based on promoters induced by heat shock, interferon and heavy metals have been described (Wurn et al., 1986, Proc. Natl. Acad. Sci. USA 83: 5414-5418; Arnheiter et al., 1990 Cell 62:51-61; Filmus et al., 1992 Nucleic Acids Research 20:27550-27560). However, these systems have limitations due to their effect on expression of non-target genes. These systems are also leaky.

[0006] Prokaryotic repressor-operator systems utilize bacterial repressor proteins and the unique operator DNA sequences to which they bind. Both the tetracycline ("Tet") and lactose ("Lac") repressor-operator systems from the bacterium *Escherichia coli* have been used in plants and animals to control gene expression. In the Tet system, tetracycline binds to the TetR repressor protein, resulting in a conformational change which releases the repressor protein from the operator which as a result allows transcription to occur. In the Lac system, a lac operon is activated in response to the presence of lactose, or synthetic analogs such as isopropyl-b-D-thiogalactoside. Unfortunately, the use of such systems is restricted by unstable chemistry of the ligands, *i.e.* tetracycline and lactose, their toxicity, their natural presence, or the relatively high levels required for induction or repression. For similar reasons, utility of such systems in animals is limited.

[0007] Immunosuppressive molecules such as FK506, rapamycin and cyclosporine A can bind to immunophilins FKBP12, cyclophilin, *etc.* Using this information, a general strategy has been devised to bring together any two proteins simply by placing FK506 on each of the two proteins or by placing FK506 on one and cyclosporine A on another one. A synthetic homodimer of FK506 (FK1012) or a compound resulted from fusion of FK506-cyclosporine (FKCsA) can then be used to induce dimerization of these molecules (Spencer et al., 1993, *Science* 262:1019-24; Belshaw et al., 1996 *Proc Natl Acad Sci U S A* 93:4604-7). Gal4 DNA binding domain fused to FKBP12 and VP16 activator domain fused to cyclophilin, and FKCsA compound were used to show heterodimerization and activation of a reporter gene under the control of a promoter containing Gal4 binding sites. Unfortunately, this system includes immunosuppressants that can have unwanted side effects and therefore, limits its use for various mammalian gene switch applications.

[0008] Higher eukaryotic transcription activation systems such as steroid hormone receptor systems have also been employed. Steroid hormone receptors are members of the nuclear receptor superfamily and are found in vertebrate and invertebrate cells. Unfortunately, use of steroidal compounds that activate the receptors for the regulation of gene expression, particularly in plants and mammals, is limited due to their involvement in many other natural biological pathways in such organisms. In order to overcome such difficulties, an alternative system has been developed using insect ecdysone receptors (EcR).

[0009] Growth, molting, and development in insects are regulated by the ecdysone steroid hormone (molting hormone)

and the juvenile hormones (Dhadialla, et al., 1998. Annu. Rev. Entomol. 43: 545-569). The molecular target for ecdysone in insects consists of at least ecdysone receptor (EcR) and ultraspiracle protein (USP). EcR is a member of the nuclear steroid receptor super family that is characterized by signature DNA and ligand binding domains, and an activation domain (Koelle et al. 1991, Cell, 67:59-77). EcR receptors are responsive to a number of steroidal compounds such as ponasterone A and muristerone A. Recently, non-steroidal compounds with ecdysteroid agonist activity have been described, including the commercially available insecticides tebufenozide and methoxyfenozide that are marketed world wide by Rohm and Haas Company (see International Patent Application No. PCT/EP96/00686 (WO 96/27673) and US Patent 5,530,028). Both analogs have exceptional safety profiles to other organisms.

[0010] International Patent Application No. PCT/US97/05330 (WO 97/38117) discloses methods for modulating the expression of an exogenous gene in which a DNA construct comprising the exogenous gene and an ecdysone response element is activated by a second DNA construct comprising an ecdysone receptor that, in the presence of a ligand therefor, and optionally in the presence of a receptor capable of acting as a silent partner, binds to the ecdysone response element to induce gene expression. The ecdysone receptor of choice was isolated from *Drosophila melanogaster*. Typically, such systems require the presence of the silent partner, preferably retinoid X receptor (RXR), in order to provide optimum activation. In mammalian cells, insect ecdysone receptor (EcR) heterodimerizes with retinoid X receptor (RXR) and regulates expression of target genes in a ligand dependent manner. International Patent Application No. PCT/US98/14215 (WO 99/02683) discloses that the ecdysone receptor isolated from the silk moth *Bombyx mori* is functional in mammalian systems without the need for an exogenous dimer partner.

[0011] U.S. Patent No. 5,880,333 discloses a *Drosophila melanogaster* EcR and ultraspiracle (USP) heterodimer system used in plants in which the transactivation domain and the DNA binding domain are positioned on two different hybrid proteins. Unfortunately, this system is not effective for inducing reporter gene expression in animal cells (for comparison, see Example 1.2, below).

[0012] In each of these cases, the transactivation domain and the DNA binding domain (either as native EcR as in international Patent Application No. PCT/US98/14215 or as modified EcR as in International Patent Application No. PCT/US97/05330) were incorporated into a single molecule and the other heterodimeric partners, either USP or RXR, were used in their full length form.

[0013] Drawbacks of the above described EcR-based gene regulation systems include a considerable background activity in the absence of ligands and that these systems are not applicable for use in both plants and animals (see U.S. Patent No. 5,880,333). For most applications that rely on modulating gene expression, these EcR-based systems are undesirable. Therefore, a need exists in the art for improved systems to precisely modulate the expression of exogenous genes in both plants and animals. Such improved systems would be useful for applications such as gene therapy, large scale production of proteins and antibodies, cell-based high throughput screening assays, functional genomics and regulation of traits in transgenic animals. Improved systems that are simple, compact, and dependent on ligands that are relatively inexpensive, readily available, and of low toxicity to the host would prove useful for regulating biological systems.

[0014] Various publications are cited herein, the disclosures of which are incorporated by reference in their entireties. However, the citation of any reference herein should not be construed as an admission that such reference is available as "Prior Art" to the instant application.

SUMMARY OF THE INVENTION

[0015] The present invention relates to a novel ecdysone receptor-based inducible gene expression system, novel receptor polynucleotides and polypeptides for use in the novel inducible gene expression system, and methods of modulating the expression of a gene within a host cell using this inducible gene expression system. In particular, Applicants' invention relates to an improved gene expression modulation system comprising a polynucleotide encoding a receptor polypeptide comprising a truncation mutation.

[0016] Specifically, the present invention relates to a gene expression modulation system comprising: a) a first gene expression cassette that is capable of being expressed in a host cell comprising a polynucleotide that encodes a first polypeptide comprising: i) a DNA-binding domain that recognizes a response element associated with a gene whose expression is to be modulated; and ii) a ligand binding domain comprising a ligand binding domain from a nuclear receptor; and b) a second gene expression cassette that is capable of being expressed in the host cell comprising a polynucleotide sequence that encodes a second polypeptide comprising: i) a transactivation domain; and ii) a ligand binding domain comprising a ligand binding domain from a nuclear receptor other than an ultraspiracle receptor; wherein the DNA binding domain and the transactivation domain are from a polypeptide other than an ecdysone receptor, a retinoid X receptor, or an ultraspiracle receptor, wherein the ligand binding domains from the first polypeptide and the second polypeptide are different and dimerize.

[0017] In a specific embodiment, the ligand binding domain of the first polypeptide comprises an ecdysone receptor (EcR) ligand binding domain

[0018] In another specific embodiment, the ligand binding domain of the second polypeptide comprises a retinoid X receptor (RXR) ligand binding domain.

[0019] In a preferred embodiment, the ligand binding domain of the first polypeptide comprises an ecdysone receptor ligand binding domain and the ligand binding domain of the second polypeptide comprises a retinoid X receptor ligand binding domain

[0020] The present invention also relates to a gene expression modulation system according to the invention further comprising c) a third gene expression cassette comprising: i) a response element to which the DNA-binding domain of the first polypeptide binds; ii) a promoter that is activated by the transactivation domain of the second polypeptide; and iii) the gene whose expression is to be modulated.

[0021] The present invention also relates to an isolated polynucleotide encoding a truncated EcR or a truncated RXR polypeptide, wherein the truncation mutation affects ligand binding activity or ligand sensitivity.

[0022] In particular, the present invention relates to an isolated polynucleotide encoding a truncated EcR or a truncated RXR polypeptide comprising a truncation mutation that reduces ligand binding activity or ligand sensitivity of said EcR or RXR polypeptide. In a specific embodiment, the present invention relates to an isolated polynucleotide encoding a truncated EcR or a truncated RXR polypeptide comprising a truncation mutation that reduces steroid binding activity or steroid sensitivity of said EcR or RXR polypeptide. In another specific embodiment, the present invention relates to an isolated polynucleotide encoding a truncated EcR or a truncated RXR polypeptide comprising a truncation mutation that reduces non-steroid binding activity or non-steroid sensitivity of said EcR or RXR polypeptide.

[0023] The present invention also relates to an isolated polynucleotide encoding a truncated EcR or a truncated RXR polypeptide comprising a truncation mutation that enhances ligand binding activity or ligand sensitivity of said EcR or RXR polypeptide. In a specific embodiment, the present invention relates to an isolated polynucleotide encoding a truncated EcR or a truncated RXR polypeptide comprising a truncation mutation that enhances steroid binding activity or steroid sensitivity of said EcR or RXR polypeptide. In another specific embodiment, the present invention relates to an isolated polynucleotide encoding a truncated EcR or a truncated RXR polypeptide comprising a truncation mutation that enhances non-steroid binding activity or non-steroid sensitivity of said EcR or RXR polypeptide.

[0024] The present invention also relates to an isolated polynucleotide encoding a truncated RXR polypeptide comprising a truncation mutation that increases ligand sensitivity of a heterodimer comprising the truncated retinoid X receptor polypeptide and a dimerization partner. In a specific embodiment, the dimerization partner is an ecdysone receptor polypeptide.

[0025] The present invention also relates to an isolated polypeptide encoded by a polynucleotide according to Applicants' invention. In particular, the present invention relates to an isolated truncated EcR or truncated RXR polypeptide comprising a truncation mutation, wherein the EcR or RXR polypeptide is encoded by a polynucleotide according to the invention.

[0026] Thus, the present invention also relates to an isolated truncated EcR or truncated RXR polypeptide comprising a truncation mutation that affects ligand binding activity or ligand sensitivity of said EcR or RXR polypeptide.

[0027] Applicants' invention also relates to methods of modulating gene expression in a host cell using a gene expression modulation system according to the invention. Specifically, Applicants' invention provides a method of modulating the expression of a gene in a host cell comprising the gene to be modulated comprising the steps of: a) introducing into the host cell a gene expression modulation system according to the invention; and b) introducing into the host cell a ligand that independently combines with the ligand binding domains of the first polypeptide and the second polypeptide of the gene expression modulation system; wherein the gene to be expressed is a component of a chimeric gene comprising: i) a response element comprising a domain to which the DNA binding domain from the first polypeptide binds; ii) a promoter that is activated by the transactivation domain of the second polypeptide; and iii) the gene whose expression is to be modulated, whereby a complex is formed comprising the ligand, the first polypeptide, and the second polypeptide, and whereby the complex modulates expression of the gene in the host cell.

[0028] Applicants' invention also provides an isolated host cell comprising an inducible gene expression system according to the invention. The present invention also relates to an isolated host cell comprising a polynucleotide or polypeptide according to the invention. Accordingly, Applicants' invention also relates to a non-human organism comprising a host cell according to the invention.

BRIEF DESCRIPTION OF THE DRAWINGS

[0029]

Figure 1: An ecdysone receptor-based gene expression system comprising a first gene expression cassette encoding a Gal4DBD-CfEcRDEF chimeric polypeptide and a second gene expression cassette encoding a VP16AD-MmRXR-DEF chimeric polypeptide; prepared as described in Example 1 (switch 1.1).

Figure 2: An ecdysone receptor-based gene expression system comprising a first gene expression cassette encoding

a Gal4DBD-CfEcRDEF chimeric polypeptide and a second gene expression cassette encoding a VP16AD-CfUSP-DEF chimeric polypeptide; prepared as described in Example 1 (switch 1.2).

Figure 3: An ecdysone receptor-based gene expression system comprising a first gene expression cassette encoding a Ga14DBD-MmRXRDEF chimeric polypeptide and a second gene expression cassette encoding a VP16AD-CfE-cRCDEF chimeric polypeptide; prepared as described in Example 1 (switch 1.3).

Figure 4: An ecdysone receptor-based gene expression system comprising a first gene expression cassette encoding a Ga14DBD-MmRXRDEF chimeric polypeptide and a second gene expression cassette encoding a VP16AD-DmE-cRCDEF chimeric polypeptide; prepared as described in Example 1 (switch 1.4).

Figure 5: An ecdysone receptor-based gene expression system comprising a first gene expression cassette encoding a Ga14DBD-CfUSPDEF chimeric polypeptide and a second gene expression cassette encoding a VP16AD-CfE-cRCDEF chimeric polypeptide; prepared as described in Example 1 (switch 1.5).

Figure 6: An ecdysone receptor-based gene expression system comprising a first gene expression cassette encoding a Ga14DBD-CfEcRDEF-VP16AD chimeric polypeptide; prepared as described in Example 1 (switch 1.6).

Figure 7: An ecdysone receptor-based gene expression system comprising a first gene expression cassette encoding a VP16AD-CfEcRCDEF chimeric polypeptide; prepared as described in Example 1 (switch 1.7).

Figure 8: An ecdysone receptor-based gene expression system comprising a first gene expression cassette encoding a VP16AD-DmEcRCDEF chimeric polypeptide and a second gene expression cassette encoding a MmRXR polypeptide; prepared as described in Example 1 (switch 1.8).

Figure 9: An ecdysone receptor-based gene expression system comprising a first gene expression cassette encoding a VP16AD-CfEcRCDEF chimeric polypeptide and a second gene expression cassette encoding a MmRXR polypeptide; prepared as described in Example 1 (switch 1.9).

Figure 10: An ecdysone receptor-based gene expression system comprising a gene expression cassette encoding a Gal4DBD-CfEcRCDEF chimeric polypeptide; prepared as described in Example 1 (switch 1.10).

Figure 11: Expression data of GAL4CfEcRA/BCDEF, GAL4CfEcRCDEF, GAL4CfEcR1/2CDEF, GAL4CfEcRDEF, GAL4CfEcREF, GAL4CfEcRDE truncation mutants transfected into NIH3T3 cells along with VP16MmRXRDE, pFRLUc and pTKRL plasmid DNAs.

Figure 12: Expression data of GAL4CfEcRA/BCDEF, GAL4CfEcRCDEF, GAL4CfEcR1/2CDEF, GAL4CfEcRDEF, GAL4CfEcREF, GAL4CfEcRDE truncation mutants transfected into 3T3 cells along with VP16MmRXRE, pFRLUc and pTKRL plasmid DNAs.

Figure 13: Expression data of VP16MmRXRA/BCDEF, VP16MmRXRCDEF, VP16MmRXRDEF, VP16MmRXREF, VP16MmRXRBam-EF, VP16MmRXRAF2del constructs transfected into NIH3T3 cells along with GALACfEcRCDEF, pFRLUc and pTKRL plasmid DNAs.

Figure 14: Expression data of VP16MmRXRA/BCDEF, VP16MmRXRCDEF, VP16MmRXRDEF, VP16MmRXREF, VP16MmRXRBam-EF, VP16MmRXRAF2del constructs transfected into NLH3T3 cells along with GALACfEcRDEF, pFRLUc and pTKRL plasmid DNAs.

Figure 15: Expression data of various truncated CfEcR and MmRXR receptor pairs transfected into NIH3T3 cells along with GAL4CfEcRDEF, pFRLUc and pTKRL plasmid DNAs.

DETAILED DESCRIPTION OF THE INVENTION

[0030] Applicants have now developed an improved ecdysone receptor-based inducible gene expression system comprising a truncation mutant of an ecdysone receptor or a retinoid X receptor (RXR) polypeptide that affects ligand binding activity or ligand sensitivity. This mutational effect may increase or reduce ligand binding activity or ligand sensitivity and may be steroid or non-steroid specific. Thus, Applicants' invention provides an improved ecdysone receptor-based inducible gene expression system useful for modulating expression of a gene of interest in a host cell. In a particularly desirable embodiment, Applicants' invention provides an inducible gene expression system that has a reduced level of background gene expression and responds to submicromolar concentrations of non-steroidal ligand. Thus, Applicants' novel inducible gene expression system and its use in methods of modulating gene expression in a host cell overcome the limitations of currently available inducible expression systems and provide the skilled artisan with an effective means to control gene expression.

[0031] The present invention provides a novel inducible gene expression system that can be used to modulate gene expression in both prokaryotic and eukaryotic host cells. Applicants' invention is useful for applications such as gene therapy, large scale production of proteins and antibodies, cell-based high throughput screening assays, functional genomics and regulation of traits in transgenic organisms.

DEFINITIONS

[0032] In this disclosure, a number of terms and abbreviations are used. The following definitions are provided and

should be helpful in understanding the scope and practice of the present invention.

[0033] In a specific embodiment, the term "about" or "approximately" means within 20%, preferably within 10%, more preferably within 5%, and even more preferably within 1% of a given value or range.

[0034] The term "substantially free" means that a composition comprising "A" (where "A" is a single protein, DNA molecule, vector, recombinant host cell, etc.) is substantially free of "B" (where "B" comprises one or more contaminating proteins, DNA molecules, vectors, etc.) when at least about 75% by weight of the proteins, DNA, vectors (depending on the category of species to which A and B belong) in the composition is "A". Preferably, "A" comprises at least about 90% by weight of the A+B species in the composition, most preferably at least about 99% by weight. It is also preferred that a composition, which is substantially free of contamination, contain only a single molecular weight species having the activity or characteristic of the species of interest.

[0035] The term "isolated" for the purposes of the present invention designates a biological material (nucleic acid or protein) that has been removed from its original environment (the environment in which it is naturally present).

[0036] For example, a polynucleotide present in the natural state in a plant or an animal is not isolated. The same polynucleotide separated from the adjacent nucleic acids in which it is naturally present. The term "purified" does not require the material to be present in a form exhibiting absolute purity, exclusive of the presence of other compounds. It is rather a relative definition.

[0037] A polynucleotide is in the "purified" state after purification of the starting material or of the natural material by at least one order of magnitude, preferably 2 or 3 and preferably 4 or 5 orders of magnitude.

[0038] A "nucleic acid" is a polymeric compound comprised of covalently linked subunits called nucleotides. Nucleic acid includes polyribonucleic acid (RNA) and polydeoxyribonucleic acid (DNA), both of which may be single-stranded or double-stranded. DNA includes but is not limited to cDNA, genomic DNA, plasmids DNA, synthetic DNA, and semi-synthetic DNA. DNA may be linear, circular, or supercoiled.

[0039] A "nucleic acid molecule" refers to the phosphate ester polymeric form of nucleosides (adenosine, guanosine, uridine or cytidine; "RNA molecules") or deoxyribonucleosides (deoxyadenosine, deoxyguanosine, deoxythymidine, or deoxycytidine; "DNA molecules"), or any phosphoester analogs thereof, such as phosphorothioates and thioesters, in either single stranded form, or a double-stranded helix. Double stranded DNA-DNA, DNA-RNA and RNA-RNA helices are possible. The term nucleic acid molecule, and in particular DNA or RNA molecule, refers only to the primary and secondary structure of the molecule, and does not limit it to any particular tertiary forms. Thus, this term includes double-stranded DNA found, *inter alia*, in linear or circular DNA molecules (*e.g.*, restriction fragments), plasmids, and chromosomes. In discussing the structure of particular double-stranded DNA molecules, sequences may be described herein according to the normal convention of giving only the sequence in the 5' to 3' direction along the non-transcribed strand of DNA (*i.e.*, the strand having a sequence homologous to the mRNA). A "recombinant DNA molecule" is a DNA molecule that has undergone a molecular biological manipulation.

[0040] The term "fragment" will be understood to mean a nucleotide sequence of reduced length relative to the reference nucleic acid and comprising, over the common portion, a nucleotide sequence identical to the reference nucleic acid. Such a nucleic acid fragment according to the invention may be, where appropriate, included in a larger polynucleotide of which it is a constituent. Such fragments comprise, or alternatively consist of, oligonucleotides ranging in length from at least 8, 10, 12, 15, 18, 20 to 25, 30, 40, 50, 70, 80, 100, 200, 500, 1000 or 1500 consecutive nucleotides of a nucleic acid according to the invention.

[0041] As used herein, an "isolated nucleic acid fragment" is a polymer of RNA or DNA that is single- or double-stranded, optionally containing synthetic, non-natural or altered nucleotide bases. An isolated nucleic acid fragment in the form of a polymer of DNA may be comprised of one or more segments of cDNA, genomic DNA or synthetic DNA.

[0042] A "gene" refers to an assembly of nucleotides that encode a polypeptide, and includes cDNA and genomic DNA nucleic acids. "Gene" also refers to a nucleic acid fragment that expresses a specific protein or polypeptide, including regulatory sequences preceding (5' non-coding sequences) and following (3' non-coding sequences) the coding sequence. "Native gene" refers to a gene as found in nature with its own regulatory sequences. "Chimeric gene" refers to any gene that is not a native gene, comprising regulatory and/or coding sequences that are not found together in nature. Accordingly, a chimeric gene may comprise regulatory sequences and coding sequences that are derived from different sources, or regulatory sequences and coding sequences derived from the same source, but arranged in a manner different than that found in nature. A chimeric gene may comprise coding sequences derived from different sources and/or regulatory sequences derived from different sources. "Endogenous gene" refers to a native gene in its natural location in the genome of an organism. A "foreign" gene or "heterologous" gene refers to a gene not normally found in the host organism, but that is introduced into the host organism by gene transfer. Foreign genes can comprise native genes inserted into a non-native organism, or chimeric genes. A "transgene" is a gene that has been introduced into the genome by a transformation procedure.

[0043] "Heterologous" DNA refers to DNA not naturally located in the cell, or in a chromosomal site of the cell. Preferably, the heterologous DNA includes a gene foreign to the cell.

[0044] The term "genome" includes chromosomal as well as mitochondrial, chloroplast and viral DNA or RNA.

[0045] A nucleic acid molecule is "hybridizable" to another nucleic acid molecule, such as a cDNA, genomic DNA, or RNA, when a single stranded form of the nucleic acid molecule can anneal to the other nucleic acid molecule under the appropriate conditions of temperature and solution ionic strength (see *Sambrook et al.*, 1989 *infra*). Hybridization and washing conditions are well known and exemplified in Sambrook, J., Fritsch, E. F. and Maniatis, T. *Molecular Cloning: A Laboratory Manual*, Second Edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor (1989), particularly Chapter 11 and Table 11.1 therein (entirely incorporated herein by reference). The conditions of temperature and ionic strength determine the "stringency" of the hybridization.

[0046] Stringency conditions can be adjusted to screen for moderately similar fragments, such as homologous sequences from distantly related organisms, to highly similar fragments, such as genes that duplicate functional enzymes from closely related organisms. For preliminary screening for homologous nucleic acids, low stringency hybridization conditions, corresponding to a T_m of 55°, can be used, e.g., 5x SSC, 0.1% SDS, 0.25% milk, and no formamide; or 30% formamide, 5x SSC, 0.5% SDS. Moderate stringency hybridization conditions correspond to a higher T_m , e.g., 40% formamide, with 5x or 6x SSC. High stringency hybridization conditions correspond to the highest T_m , e.g., 50% formamide, 5x or 6x SSC. Hybridization requires that the two nucleic acids contain complementary sequences, although depending on the stringency of the hybridization, mismatches between bases are possible.

[0047] The term "complementary" is used to describe the relationship between nucleotide bases that are capable of hybridizing to one another. For example, with respect to DNA, adenosine is complementary to thymine and cytosine is complementary to guanine. Accordingly, the instant invention also includes isolated nucleic acid fragments that are complementary to the complete sequences as disclosed or used herein as well as those substantially similar nucleic acid sequences.

[0048] In a specific embodiment, the term "standard hybridization conditions" refers to a T_m of 55°C, and utilizes conditions as set forth above. In a preferred embodiment, the T_m is 60°C; in a more preferred embodiment, the T_m is 65°C.

[0049] Post-hybridization washes also determine stringency conditions. One set of preferred conditions uses a series of washes starting with 6X SSC, 0.5% SDS at room temperature for 15 minutes (min), then repeated with 2X SSC, 0.5% SDS at 45°C for 30 minutes, and then repeated twice with 0.2X SSC, 0.5% SDS at 50°C for 30 minutes. A more preferred set of stringent conditions uses higher temperatures in which the washes are identical to those above except for the temperature of the final two 30 min washes in 0.2X SSC, 0.5% SDS was increased to 60°C. Another preferred set of highly stringent conditions uses two final washes in 0.1X SSC, 0.1% SDS at 65°C. Hybridization requires that the two nucleic acids comprise complementary sequences, although depending on the stringency of the hybridization, mismatches between bases are possible.

[0050] The appropriate stringency for hybridizing nucleic acids depends on the length of the nucleic acids and the degree of complementation, variables well known in the art. The greater the degree of similarity or homology between two nucleotide sequences, the greater the value of T_m for hybrids of nucleic acids having those sequences. The relative stability (corresponding to higher T_m) of nucleic acid hybridizations decreases in the following order: RNA:RNA, DNA:RNA, DNA:DNA. For hybrids of greater than 100 nucleotides in length, equations for calculating T_m have been derived (see *Sambrook et al.*, *supra*, 9.50-0.51). For hybridization with shorter nucleic acids, i.e., oligonucleotides, the position of mismatches becomes more important, and the length of the oligonucleotide determines its specificity (see *Sambrook et al.*, *supra*, 11.7-11.8).

[0051] In one embodiment the length for a hybridizable nucleic acid is at least about 10 nucleotides. Preferable a minimum length for a hybridizable nucleic acid is at least about 15 nucleotides; more preferably at least about 20 nucleotides; and most preferably the length is at least 30 nucleotides. Furthermore, the skilled artisan will recognize that the temperature and wash solution salt concentration may be adjusted as necessary according to factors such as length of the probe.

[0052] The term "probe" refers to a single-stranded nucleic acid molecule that can base pair with a complementary single stranded target nucleic acid to form a double-stranded molecule.

[0053] As used herein, the term "oligonucleotide" refers to a nucleic acid, generally of at least 18 nucleotides, that is hybridizable to a genomic DNA molecule, a cDNA molecule, a plasmid DNA or an mRNA molecule. Oligonucleotides can be labeled, e.g., with ³²P-nucleotides or nucleotides to which a label, such as biotin, has been covalently conjugated. A labeled oligonucleotide can be used as a probe to detect the presence of a nucleic acid. Oligonucleotides (one or both of which may be labeled) can be used as PCR primers, either for cloning full length or a fragment of a nucleic acid, or to detect the presence of a nucleic acid. An oligonucleotide can also be used to form a triple helix with a DNA molecule. Generally, oligonucleotides are prepared synthetically, preferably on a nucleic acid synthesizer. Accordingly, oligonucleotides can be prepared with non-naturally occurring phosphoester analog bonds, such as thioester bonds, etc.

[0054] A "primer" is an oligonucleotide that hybridizes to a target nucleic acid sequence to create a double stranded nucleic acid region that can serve as an initiation point for DNA synthesis under suitable conditions. Such primers may be used in a polymerase chain reaction.

[0055] "Polymerase chain reaction" is abbreviated PCR and means an *in vitro* method for enzymatically amplifying specific nucleic acid sequences. PCR involves a repetitive series of temperature cycles with each cycle comprising three

stages: denaturation of the template nucleic acid to separate the strands of the target molecule, annealing a single stranded PCR oligonucleotide primer to the template nucleic acid, and extension of the annealed primer(s) by DNA polymerase. PCR provides a means to detect the presence of the target molecule and, under quantitative or semi-

quantitative conditions, to determine the relative amount of that target molecule within the starting pool of nucleic acids.

[0056] "Reverse transcription-polymerase chain reaction" is abbreviated RT-PCR and means an *in vitro* method for enzymatically producing a target cDNA molecule or molecules from an RNA molecule or molecules, followed by enzymatic amplification of a specific nucleic acid sequence or sequences within the target cDNA molecule or molecules as described above. RT-PCR also provides a means to detect the presence of the target molecule and, under quantitative or semi-quantitative conditions, to determine the relative amount of that target molecule within the starting pool of nucleic acids.

[0057] A DNA "coding sequence" is a double-stranded DNA sequence that is transcribed and translated into a polypeptide in a cell *in vitro* or *in vivo* when placed under the control of appropriate regulatory sequences. "Suitable regulatory sequences" refer to nucleotide sequences located upstream (5' non-coding sequences), within, or downstream (3' non-coding sequences) of a coding sequence, and which influence the transcription, RNA processing or stability, or translation of the associated coding sequence. Regulatory sequences may include promoters, translation leader sequences, introns, polyadenylation recognition sequences, RNA processing site, effector binding site and stem-loop structure. The boundaries of the coding sequence are determined by a start codon at the 5' (amino) terminus and a translation stop codon at the 3' (carboxyl) terminus. A coding sequence can include, but is not limited to, prokaryotic sequences, cDNA from mRNA, genomic DNA sequences, and even synthetic DNA sequences. If the coding sequence is intended for expression in a eukaryotic cell, a polyadenylation signal and transcription termination sequence will usually be located 3' to the coding sequence.

[0058] "Open reading frame" is abbreviated ORF and means a length of nucleic acid sequence, either DNA, cDNA or RNA, that comprises a translation start signal or initiation codon, such as an ATG or AUG, and a termination codon and can be potentially translated into a polypeptide sequence.

[0059] The term "head-to-head" is used herein to describe the orientation of two polynucleotide sequences in relation to each other. Two polynucleotides are positioned in a head-to-head orientation when the 5' end of the coding strand of one polynucleotide is adjacent to the 5' end of the coding strand of the other polynucleotide, whereby the direction of transcription of each polynucleotide proceeds away from the 5' end of the other polynucleotide. The term "head-to-head" may be abbreviated (5')-to-(5') and may also be indicated by the symbols ($\leftarrow \rightarrow$) or (3'←5'→3').

[0060] The term "tail-to-tail" is used herein to describe the orientation of two polynucleotide sequences in relation to each other. Two polynucleotides are positioned in a tail-to-tail orientation when the 3' end of the coding strand of one polynucleotide is adjacent to the 3' end of the coding strand of the other polynucleotide, whereby the direction of transcription of each polynucleotide proceeds toward the other polynucleotide. The term "tail-to-tail" may be abbreviated (3')-to-(3') and may also be indicated by the symbols ($\rightarrow \leftarrow$) or (5'→3'3'←5').

[0061] The term "head-to-tail" is used herein to describe the orientation of two polynucleotide sequences in relation to each other. Two polynucleotides are positioned in a head-to-tail orientation when the 5' end of the coding strand of one polynucleotide is adjacent to the 3' end of the coding strand of the other polynucleotide, whereby the direction of transcription of each polynucleotide proceeds in the same direction as that of the other polynucleotide. The term "head-to-tail" may be abbreviated (5')-to-(3') and may also be indicated by the symbols ($\rightarrow \rightarrow$) or (5'→3'5'→3').

[0062] The term "downstream" refers to a nucleotide sequence that is located 3' to reference nucleotide sequence. In particular, downstream nucleotide sequences generally relate to sequences that follow the starting point of transcription. For example, the translation initiation codon of a gene is located downstream of the start site of transcription.

[0063] The term "upstream" refers to a nucleotide sequence that is located 5' to reference nucleotide sequence. In particular, upstream nucleotide sequences generally relate to sequences that are located on the 5' side of a coding sequence or starting point of transcription. For example, most promoters are located upstream of the start site of transcription.

[0064] The terms "restriction endonuclease" and "restriction enzyme" refer to an enzyme that binds and cuts within a specific nucleotide sequence within double stranded DNA.

[0065] "Homologous recombination" refers to the insertion of a foreign DNA sequence into another DNA molecule, e.g., insertion of a vector in a chromosome. Preferably, the vector targets a specific chromosomal site for homologous recombination. For specific homologous recombination, the vector will contain sufficiently long regions of homology to sequences of the chromosome to allow complementary binding and incorporation of the vector into the chromosome. Longer regions of homology, and greater degrees of sequence similarity, may increase the efficiency of homologous recombination.

[0066] Several methods known in the art may be used to propagate a polynucleotide according to the invention. Once a suitable host system and growth conditions are established, recombinant expression vectors can be propagated and prepared in quantity. As described herein, the expression vectors which can be used include, but are not limited to, the following vectors or their derivatives: human or animal viruses such as vaccinia virus or adenovirus; insect viruses such as baculovirus; yeast vectors; bacteriophage vectors (e.g., lambda), and plasmid and cosmid DNA vectors, to name but

a few.

[0067] A "vector" is any means for the cloning of and/or transfer of a nucleic acid into a host cell. A vector may be a replicon to which another DNA segment may be attached so as to bring about the replication of the attached segment. A "replicon" is any genetic element (e.g., plasmid, phage, cosmid, chromosome, virus) that functions as an autonomous unit of DNA replication *in vivo*, i.e., capable of replication under its own control. The term "vector" includes both viral and nonviral means for introducing the nucleic acid into a cell *in vitro*, *ex vivo* or *in vivo*. A large number of vectors known in the art may be used to manipulate nucleic acids, incorporate response elements and promoters into genes, etc. Possible vectors include, for example, plasmids or modified viruses including, for example bacteriophages such as lambda derivatives, or plasmids such as PBR322 or pUC plasmid derivatives, or the Bluescript vector. For example, the insertion of the DNA fragments corresponding to response elements and promoters into a suitable vector can be accomplished by ligating the appropriate DNA - fragments into a chosen vector that has complementary cohesive termini. Alternatively, the ends of the DNA molecules may be enzymatically modified or any site may be produced by ligating nucleotide sequences (linkers) into the DNA termini. Such vectors may be engineered to contain selectable marker genes that provide for the selection of cells that have incorporated the marker into the cellular genome. Such markers allow identification and/or selection of host cells that incorporate and express the proteins encoded by the marker.

[0068] Viral vectors, and particularly retroviral vectors, have been used in a wide variety of gene delivery applications in cells, as well as living animal subjects. Viral vectors that can be used include but are not limited to retrovirus, adeno-associated virus, pox, baculovirus, vaccinia, herpes simplex, Epstein-Barr, adenovirus, geminivirus, and caulimovirus vectors. Non-viral vectors include plasmids, liposomes, electrically charged lipids (cytofectins), DNA-protein complexes, and biopolymers. In addition to a nucleic acid, a vector may also comprise one or more regulatory regions, and/or selectable markers useful in selecting, measuring, and monitoring nucleic acid transfer results (transfer to which tissues, duration of expression, etc.).

[0069] The term "plasmid" refers to an extra chromosomal element often carrying a gene that is not part of the central metabolism of the cell, and usually in the form of circular double-stranded DNA molecules. Such elements may be autonomously replicating sequences, genome integrating sequences, phage or nucleotide sequences, linear, circular, or supercoiled, of a single- or double-stranded DNA or RNA, derived from any source, in which a number of nucleotide sequences have been joined or recombined into a unique construction which is capable of introducing a promoter fragment and DNA sequence for a selected gene product along with appropriate 3' untranslated sequence into a cell.

[0070] A "cloning vector" is a "replicon", which is a unit length of a nucleic acid, preferably DNA, that replicates sequentially and which comprises an origin of replication, such as a plasmid, phage or cosmid, to which another nucleic acid segment may be attached so as to bring about the replication of the attached segment. Cloning vectors may be capable of replication in one cell type and expression in another ("shuttle vector").

[0071] Vectors may be introduced into the desired host cells by methods known in the art, e.g., transfection, electroporation, microinjection, transduction, cell fusion, DEAE dextran, calcium phosphate precipitation, lipofection (lysosome fusion), use of a gene gun, or a DNA vector transporter (see, e.g., Wu et al., 1992, J. Biol. Chem. 267:963-967; Wu and Wu, 1988, J. Biol. Chem. 263:14621-14624; and Hartmut et al., Canadian Patent Application No. 2,012,311, filed March 15, 1990).

[0072] A polynucleotide according to the invention can also be introduced *in vivo* by lipofection. For the past decade, there has been increasing use of liposomes for encapsulation and transfection of nucleic acids *in vitro*. Synthetic cationic lipids designed to limit the difficulties and dangers encountered with liposome mediated transfection can be used to prepare liposomes for *in vivo* transfection of a gene encoding a marker (Felper et al., 1987, PNAS 84:7413; Mackey, et al., 1988, Proc. Natl. Acad. Sci. U.S.A 85:8027-8031; and Ulmer et al., 1993, Science 259:1745-1748). The use of cationic lipids may promote encapsulation of negatively charged nucleic acids, and also promote fusion with negatively charged cell membranes (Felgner and Ringold, 1989, Science 337:387-388). Particularly useful lipid compounds and compositions for transfer of nucleic acids are described in International Patent Publications WO95/18863 and WO96/17823, and in U.S. Patent No. 5,459,127. The use of lipofection to introduce exogenous genes into the specific organs *in vivo* has certain practical advantages. Molecular targeting of liposomes to specific cells represents one area of benefit. It is clear that directing transfection to particular cell types would be particularly preferred in a tissue with cellular heterogeneity, such as pancreas, liver, kidney, and the brain. Lipids may be chemically coupled to other molecules for the purpose of targeting (Mackey, et al., 1988, *supra*). Targeted peptides, e.g., hormones or neurotransmitters, and proteins such as antibodies, or non-peptide molecules could be coupled to liposomes chemically.

[0073] Other molecules are also useful for facilitating transfection of a nucleic acid *in vivo*, such as a cationic oligopeptide (e.g., WO95/21931), peptides derived from DNA binding proteins (e.g., WO96/25508), or a cationic polymer (e.g., WO95/21931).

[0074] It is also possible to introduce a vector *in vivo* as a naked DNA plasmid (see U.S. Patents 5,693,622, 5,589,466 and 5,580,859). Receptor-mediated DNA delivery approaches can also be used (Curiel et al., 1992, Hum. Gene Ther. 3:147-154; and Wu and Wu, 1987, J. Biol. Chem 262:4429-4432).

[0075] The term "transfection" means the uptake of exogenous or heterologous RNA or DNA by a cell. A cell has been

"transfected" by exogenous or heterologous RNA or DNA when such RNA or DNA has been introduced inside the cell. A cell has been "transformed" by exogenous or heterologous RNA or DNA when the transfected RNA or DNA effects a phenotypic change. The transforming RNA or DNA can be integrated (covalently linked) into chromosomal DNA making up the genome of the cell.

[0076] "Transformation" refers to the transfer of a nucleic acid fragment into the genome of a host organism, resulting in genetically stable inheritance. Host organisms containing the transformed nucleic acid fragments are referred to as "transgenic" or "recombinant" or "transformed" organisms.

[0077] The term "genetic region" will refer to a region of a nucleic acid molecule or a nucleotide sequence that comprises a gene encoding a polypeptide.

[0078] In addition, the recombinant vector comprising a polynucleotide according to the invention may include one or more origins for replication in the cellular hosts in which their amplification or their expression is sought, markers or selectable markers.

[0079] The term "selectable marker" means an identifying factor, usually an antibiotic or chemical resistance gene, that is able to be selected for based upon the marker gene's effect, i.e., resistance to an antibiotic, resistance to a herbicide, colorimetric markers, enzymes, fluorescent markers, and the like, wherein the effect is used to track the inheritance of a nucleic acid of interest and/or to identify a cell or organism that has inherited the nucleic acid of interest. Examples of selectable marker genes known and used in the art include: genes providing resistance to ampicillin, streptomycin, gentamycin, kanamycin, hygromycin, bialaphos herbicide, sulfonamide, and the like; and genes that are used as phenotypic markers, i.e., anthocyanin regulatory genes, isopentany transferase gene, and the like.

[0080] The term "reporter gene" means a nucleic acid encoding an identifying factor that is able to be identified based upon the reporter gene's effect, wherein the effect is used to track the inheritance of a nucleic acid of interest, to identify a cell or organism that has inherited the nucleic acid of interest, and/or to measure gene expression induction or transcription. Examples of reporter genes known and used in the art include: luciferase (Luc), green fluorescent protein (GFP), chloramphenicol acetyltransferase (CAT), β -galactosidase (LacZ), β -glucuronidase (Gus), and the like. Selectable marker genes may also be considered reporter genes.

[0081] "Promoter" refers to a DNA sequence capable of controlling the expression of a coding sequence or functional RNA. In general, a coding sequence is located 3' to a promoter sequence. Promoters may be derived in their entirety from a native gene, or be composed of different elements derived from different promoters found in nature, or even comprise synthetic DNA segments. It is understood by those skilled in the art that different promoters may direct the expression of a gene in different tissues or cell types, or at different stages of development, or in response to different environmental or physiological conditions. Promoters that cause a gene to be expressed in most cell types at most times are commonly referred to as "constitutive promoters". Promoters that cause a gene to be expressed in a specific cell type are commonly referred to as "cell-specific promoters" or "tissue-specific promoters". Promoters that cause a gene to be expressed at a specific stage of development or cell differentiation are commonly referred to as "developmentally-specific promoters" or "cell differentiation-specific promoters". Promoters that are induced and cause a gene to be expressed following exposure or treatment of the cell with an agent, biological molecule, chemical, ligand, light, or the like that induces the promoter are commonly referred to as "inducible promoters" or "regulatable promoters". It is further recognized that since in most cases the exact boundaries of regulatory sequences have not been completely defined, DNA fragments of different lengths may have identical promoter activity.

[0082] A "promoter sequence" is a DNA regulatory region capable of binding RNA polymerase in a cell and initiating transcription of a downstream (3' direction) coding sequence. For purposes of defining the present invention, the promoter sequence is bounded at its 3' terminus by the transcription initiation site and extends upstream (5' direction) to include the minimum number of bases or elements necessary to initiate transcription at levels detectable above background. Within the promoter sequence will be found a transcription initiation site (conveniently defined for example, by mapping with nuclease S1), as well as protein binding domains (consensus sequences) responsible for the binding of RNA polymerase.

[0083] A coding sequence is "under the control" of transcriptional and translational control sequences in a cell when RNA polymerase transcribes the coding sequence into mRNA, which is then trans-RNA spliced (if the coding sequence contains introns) and translated into the protein encoded by the coding sequence.

[0084] "Transcriptional and translational control sequences" are DNA regulatory sequences, such as promoters, enhancers, terminators, and the like, that provide for the expression of a coding sequence in a host cell. In eukaryotic cells, polyadenylation signals are control sequences.

[0085] The term "response element" means one or more cis-acting DNA elements which confer responsiveness on a promoter mediated through interaction with the DNA-binding domains of the first chimeric gene. This DNA element may be either palindromic (perfect or imperfect) in its sequence or composed of sequence motifs or half sites separated by a variable number of nucleotides. The half sites can be similar or identical and arranged as either direct or inverted repeats or as a single half site or multimers of adjacent half sites in tandem. The response element may comprise a minimal promoter isolated from different organisms depending upon the nature of the cell or organism into which the

response element will be incorporated. The DNA binding domain of the first hybrid protein binds, in the presence or absence of a ligand, to the DNA sequence of a response element to initiate or suppress transcription of downstream gene(s) under the regulation of this response element. Examples of DNA sequences for response elements of the natural ecdysone receptor include: RRGG/TTCANTGAC/ACY (see Cherbas L., et al., (1991), *Genes Dev.* 5, 120-131); AG-GTCAN_(n)AGGTCA, where N_(n) can be one or more spacer nucleotides (see D'Avino PP., et. al., (1995), *Mol. Cell. Endocrinol.* 113,1-9); and GGGTTGAATGAATTT (see Antoniewski C., et. al., (1994). *Mol. Cell Biol.* 14, 4465-4474).

[0086] The term "operably linked" refers to the association of nucleic acid sequences on a single nucleic acid fragment so that the function of one is affected by the other. For example, a promoter is operably linked with a coding sequence when it is capable of affecting the expression of that coding sequence (i.e., that the coding sequence is under the transcriptional control of the promoter). Coding sequences can be operably linked to regulatory sequences in sense or antisense orientation.

[0087] The term "expression", as used herein, refers to the transcription and stable accumulation of sense (mRNA) or antisense RNA derived from a nucleic acid or polynucleotide. Expression may also refer to translation of mRNA into a protein or polypeptide.

[0088] The terms "cassette", "expression cassette" and "gene expression cassette" refer to a segment of DNA that can be inserted into a nucleic acid or polynucleotide at specific restriction sites or by homologous recombination. The segment of DNA comprises a polynucleotide that encodes a polypeptide of interest, and the cassette and restriction sites are designed to ensure insertion of the cassette in the proper reading frame for transcription and translation. "Transformation cassette" refers to a specific vector comprising a polynucleotide that encodes a polypeptide of interest and having elements in addition to the polynucleotide that facilitate transformation of a particular host cell. Cassettes, expression cassettes, gene expression cassettes and transformation cassettes of the invention may also comprise elements that allow for enhanced expression of a polynucleotide encoding a polypeptide of interest in a host cell. These elements may include, but are not limited to: a promoter, a minimal promoter, an enhancer, a response element, a terminator sequence, a polyadenylation sequence, and the like.

[0089] For purposes of this invention, the term "gene switch" refers to the combination of a response element associated with a promoter, and an EcR based system which, in the presence of one or more ligands, modulates the expression of a gene into which the response element and promoter are incorporated.

[0090] The terms "modulate" and "modulates" mean to induce, reduce or inhibit nucleic acid or gene expression, resulting in the respective induction, reduction or inhibition of protein or polypeptide production.

[0091] The plasmids or vectors according to the invention may further comprise at least one promoter suitable for driving expression of a gene in a host cell. The term "expression vector" means a vector, plasmid or vehicle designed to enable the expression of an inserted nucleic acid sequence following transformation into the host. The cloned gene, i.e., the inserted nucleic acid sequence, is usually placed under the control of control elements such as a promoter, a minimal promoter, an enhancer, or the like. Initiation control regions or promoters, which are useful to drive expression of a nucleic acid in the desired host cell are numerous and familiar to those skilled in the art. Virtually any promoter capable of driving these genes is suitable for the present invention including but not limited to: viral promoters, plant promoters, bacterial promoters, animal promoters, mammalian promoters, synthetic promoters, constitutive promoters, tissue specific promoter, developmental specific promoters, inducible promoters, light regulated promoters; *CYC1*, *HIS3*, *GAL1*, *GAL4*, *GAL10*, *ADHI*, *PGK*, *PHO5*, *GAPDH*, *ADC1*, *TRP1*, *URA3*, *LEU2*, *ENO*, *TPI*, alkaline phosphatase promoters (useful for expression in *Saccharomyces*); *AOX1* promoter (useful for expression in *Pichia*); b-lactamase, *lac*, *ara*, *tet*, *trp*, *IP_L*, *IP_R*, *T7*, *tac*, and *trc* promoters (useful for expression in *Escherichia coli*); and light regulated-, seed specific-, pollen specific-, ovary specific-, pathogenesis or disease related-, cauliflower mosaic virus 35S, CMV 35S minimal, cassava vein mosaic virus (CsVMV), chlorophyll a/b binding protein, ribulose 1, 5-bisphosphate carboxylase, shoot-specific, root specific, chitinase, stress inducible, rice tungro bacilliform virus, plant super-promoter, potato leucine aminopeptidase, nitrate reductase, mannopine synthase, nopaline synthase, ubiquitin, zein protein, and anthocyanin promoters (useful for expression in plant cells); animal and mammalian promoters known in the art include, but are not limited to, the SV40 early (SV40e) promoter region, the promoter contained in the 3' long terminal repeat (LTR) of Rous sarcoma virus (RSV), the promoters of the E1A or major late promoter (MLP) genes of adenoviruses, the cytomegalovirus early promoter, the herpes simplex virus (HSV) thymidine kinase (TK) promoter, an elongation factor 1 alpha (EF1) promoter, a phosphoglycerate kinase (PGK) promoter, a ubiquitin (Ubc) promoter, an albumin promoter, the regulatory sequences of the mouse metallothionein-L promoter, and transcriptional control regions, the ubiquitous promoters (HPRT, vimentin, α -actin, tubulin and the like), the promoters of the intermediate filaments (desmin, neurofilaments, keratin, GFAP, and the like), the promoters of therapeutic genes (of the MDR, CFTR or factor VIII type, and the like), and promoters that exhibit tissue specificity and have been utilized in transgenic animals, such as the elastase I gene control region which is active in pancreatic acinar cells; insulin gene control region active in pancreatic beta cells, immunoglobulin gene control region active in lymphoid cells, mouse mammary tumor virus control region active in testicular, breast, lymphoid and mast cells; albumin gene, Apo AI and Apo AII control regions active in liver, alpha-fetoprotein gene control region active in liver, alpha 1-antitrypsin gene control region active in the liver, beta-globin gene

control region active in myeloid cells, myelin basic protein gene control region) active in oligodendrocyte cells in the brain, myosin light chain-2 gene control region active in skeletal muscle, and gonadotropic releasing hormone gene control region active in the hypothalamus, pyruvate kinase promoter, villin promoter, promoter of the fatty acid binding intestinal protein, promoter of the smooth muscle cell α -actin, and the like. In a preferred embodiment of the invention,

the promoter is selected from the group consisting of a cauliflower mosaic virus 35S promoter, a cassava vein mosaic virus promoter, and a cauliflower mosaic virus 35S minimal promoter, an elongation factor 1 alpha (EF1) promoter, a phosphoglycerate kinase (PGK) promoter, a ubiquitin (Ubc) promoter, and an albumin promoter. In addition, these expression sequences may be modified by addition of enhancer or regulatory sequences and the like.

[0092] Enhancers that may be used in embodiments of the invention include but are not limited to: tobacco mosaic virus enhancer, cauliflower mosaic virus 35S enhancer, tobacco etch virus enhancer, ribulose 1, 5-bisphosphate carboxylase enhancer, rice tungro bacilliform virus enhancer, and other plant and viral gene enhancers, and the like.

[0093] Termination control regions, *i.e.*, terminator or polyadenylation sequences, may also be derived from various genes native to the preferred hosts. Optionally, a termination site may be unnecessary, however, it is most preferred if included. In a preferred embodiment of the invention, the termination control region may be comprise or be derived from a synthetic sequence, synthetic polyadenylation signal, an SV40 late polyadenylation signal, an SV40 polyadenylation signal, a bovine growth hormone (BGH) polyadenylation signal, nopaline synthase (nos), cauliflower mosaic virus (CaMV), octopine synthase (ocs), Agrocateum, viral, and plant terminator sequences, or the like.


[0094] The terms "3' non-coding sequences" or "3' untranslated region (UTR)" refer to DNA sequences located downstream (3') of a coding sequence and may comprise polyadenylation [poly(A)] recognition sequences and other sequences encoding regulatory signals capable of affecting mRNA processing or gene expression. The polyadenylation signal is usually characterized by affecting the addition of polyadenylic acid tracts to the 3' end of the mRNA precursor.

[0095] "Regulatory region" means a nucleic acid sequence which regulates the expression of a second nucleic acid sequence. A regulatory region may include sequences which are naturally responsible for expressing a particular nucleic acid (a homologous region) or may include sequences of a different origin that are responsible for expressing different proteins or even synthetic proteins (a heterologous region). In particular, the sequences can be sequences of prokaryotic, eukaryotic, or viral genes or derived sequences that stimulate or repress transcription of a gene in a specific or non-specific manner and in an inducible or non-inducible manner. Regulatory regions include origins of replication, RNA splice sites, promoters, enhancers, transcriptional termination sequences, and signal sequences which direct the polypeptide into the secretory pathways of the target cell.

[0096] A regulatory region from a "heterologous source" is a regulatory region that is not naturally associated with the expressed nucleic acid. Included among the heterologous regulatory regions are regulatory regions from a different species, regulatory regions from a different gene, hybrid regulatory sequences, and regulatory sequences which do not occur in nature, but which are designed by one having ordinary skill in the art.

[0097] "RNA transcript" refers to the product resulting from RNA polymerase-catalyzed transcription of a DNA sequence. When the RNA transcript is a perfect complementary copy of the DNA sequence, it is referred to as the primary transcript or it may be a RNA sequence derived from post-transcriptional processing of the primary transcript and is referred to as the mature RNA. "Messenger RNA (mRNA)" refers to the RNA that is without introns and that can be translated into protein by the cell. "cDNA" refers to a double-stranded DNA that is complementary to and derived from mRNA. "Sense" RNA refers to RNA transcript that includes the mRNA and so can be translated into protein by the cell. "Antisense RNA" refers to a RNA transcript that is complementary to all or part of a target primary transcript or mRNA and that blocks the expression of a target gene. The complementarity of an antisense RNA may be with any part of the specific gene transcript, *i.e.*, at the 5' non-coding sequence, 3' non-coding sequence, or the coding sequence. "Functional RNA" refers to antisense RNA, ribozyme RNA, or other RNA that is not translated yet has an effect on cellular processes.

[0098] A "polypeptide" is a polymeric compound comprised of covalently linked amino acid residues. Amino acids have the following general structure:



Amino acids are classified into seven groups on the basis of the side chain R: (1) aliphatic side chains, (2) side chains containing a hydroxylic (OH) group, (3) side chains containing sulfur atoms, (4) side chains containing an acidic or amide

group, (5) side chains containing a basic group, (6) side chains containing an aromatic ring, and (7) proline, an imino acid in which the side chain is fused to the amino group. A polypeptide of the invention preferably comprises at least about 14 amino acids.

[0099] A "protein" is a polypeptide that performs a structural or functional role in a living cell.

[0100] An "isolated polypeptide" or "isolated protein" is a polypeptide or protein that is substantially free of those compounds that are normally associated therewith in its natural state (e.g., other proteins or polypeptides, nucleic acids, carbohydrates, lipids). "Isolated" is not meant to exclude artificial or synthetic mixtures with other compounds, or the presence of impurities which do not interfere with biological activity, and which may be present, for example, due to incomplete purification, addition of stabilizers, or compounding into a pharmaceutically acceptable preparation.

[0101] "Fragment" of a polypeptide according to the invention will be understood to mean a polypeptide whose amino acid sequence is shorter than that of the reference polypeptide and which comprises, over the entire portion with these reference polypeptides, an identical amino acid sequence. Such fragments may, where appropriate, be included in a larger polypeptide of which they are a part. Such fragments of a polypeptide according to the invention may have a length of 10, 15, 20, 30 to 40, 50, 100, 200 or 300 amino acids.

[0102] A "variant" of a polypeptide or protein is any analogue, fragment, derivative, or mutant which is derived from a polypeptide or protein and which retains at least one biological property of the polypeptide or protein. Different variants of the polypeptide or protein may exist in nature. These variants may be allelic variations characterized by differences in the nucleotide sequences of the structural gene coding for the protein, or may involve differential splicing or post-translational modification. The skilled artisan can produce variants having single or multiple amino acid substitutions, deletions, additions, or replacements. These variants may include, *inter alia*: (a) variants in which one or more amino acid residues are substituted with conservative or non-conservative amino acids, (b) variants in which one or more amino acids are added to the polypeptide or protein, (c) variants in which one or more of the amino acids includes a substituent group, and (d) variants in which the polypeptide or protein is fused with another polypeptide such as serum albumin. The techniques for obtaining these variants, including genetic (suppressions, deletions, mutations, etc.), chemical, and enzymatic techniques, are known to persons having ordinary skill in the art. A variant polypeptide preferably comprises at least about 14 amino acids.

[0103] A "heterologous protein" refers to a protein not naturally produced in the cell.

[0104] A "mature protein" refers to a post-translationally processed polypeptide; i.e., one from which any pre- or propeptides present in the primary translation product have been removed. "Precursor" protein refers to the primary product of translation of mRNA; i.e., with pre- and propeptides still present. Pre- and propeptides may be but are not limited to intracellular localization signals.

[0105] The term "signal peptide" refers to an amino terminal polypeptide preceding the secreted mature protein. The signal peptide is cleaved from and is therefore not present in the mature protein. Signal peptides have the function of directing and translocating secreted proteins across cell membranes. Signal peptide is also referred to as signal protein.

[0106] A "signal sequence" is included at the beginning of the coding sequence of a protein to be expressed on the surface of a cell. This sequence encodes a signal peptide, N-terminal to the mature polypeptide, that directs the host cell to translocate the polypeptide. The term "translocation signal sequence" is used herein to refer to this sort of signal sequence. Translocation signal sequences can be found associated with a variety of proteins native to eukaryotes and prokaryotes, and are often functional in both types of organisms.

[0107] The term "homology" refers to the percent of identity between two polynucleotide or two polypeptide moieties. The correspondence between the sequence from one moiety to another can be determined by techniques known to the art. For example, homology can be determined by a direct comparison of the sequence information between two polypeptide molecules by aligning the sequence information and using readily available computer programs. Alternatively, homology can be determined by hybridization of polynucleotides under conditions that form stable duplexes between homologous regions, followed by digestion with single-stranded-specific nuclease(s) and size determination of the digested fragments.

[0108] As used herein, the term "homologous" in all its grammatical forms and spelling variations refers to the relationship between proteins that possess a "common evolutionary origin," including proteins from superfamilies (e.g., the immunoglobulin superfamily) and homologous proteins from different species (e.g., myosin light chain, etc.) (Reeck et al., 1987, Cell 50:667.). Such proteins (and their encoding genes) have sequence homology, as reflected by their high degree of sequence similarity.

[0109] Accordingly, the term "sequence similarity" in all its grammatical forms refers to the degree of identity or correspondence between nucleic acid or amino acid sequences of proteins that may or may not share a common evolutionary origin (see Reeck et al., 1987, Cell 50:667). As used herein, the term "homologous" in all its grammatical forms and spelling variations refers to the relationship between proteins that possess a "common evolutionary origin," including proteins from superfamilies and homologous proteins from different species (Reeck et al., *supra*). Such proteins (and their encoding genes) have sequence homology, as reflected by their high degree of sequence similarity. However, in common usage and in the instant application, the term "homologous," when modified with an adverb such as "highly,"

may refer to sequence similarity and not a common evolutionary origin.

[0110] In a specific embodiment, two DNA sequences are "substantially homologous" or "substantially similar" when at least about 50% (preferably at least about 75%, and most preferably at least about 90 or 95%) of the nucleotides match over the defined length of the DNA sequences. Sequences that are substantially homologous can be identified by comparing the sequences using standard software available in sequence data banks, or in a Southern hybridization experiment under, for example, stringent conditions as defined for that particular system. Defining appropriate hybridization conditions is within the skill of the art. See, e.g., Sambrook *et al.*, 1989, *supra*.

[0111] As used herein, "substantially similar" refers to nucleic acid fragments wherein changes in one or more nucleotide bases results in substitution of one or more amino acids, but do not affect the functional properties of the protein encoded by the DNA sequence. "Substantially similar" also refers to nucleic acid fragments wherein changes in one or more nucleotide bases does not affect the ability of the nucleic acid fragment to mediate alteration of gene expression by antisense or co-suppression technology. "Substantially similar" also refers to modifications of the nucleic acid fragments of the instant invention such as deletion or insertion of one or more nucleotide bases that do not substantially affect the functional properties of the resulting transcript. It is therefore understood that the invention encompasses more than the specific exemplary sequences. Each of the proposed modifications is well within the routine skill in the art, as is determination of retention of biological activity of the encoded products.

[0112] Moreover, the skilled artisan recognizes that substantially similar sequences encompassed by this invention are also defined by their ability to hybridize, under stringent conditions (0.1X SSC, 0.1% SDS, 65°C and washed with 2X SSC, 0.1% SDS followed by 0.1X SSC, 0.1% SDS), with the sequences exemplified herein. Substantially similar nucleic acid fragments of the instant invention are those nucleic acid fragments whose DNA sequences are at least 70% identical to the DNA sequence of the nucleic acid fragments reported herein. Preferred substantially nucleic acid fragments of the instant invention are those nucleic acid fragments whose DNA sequences are at least 80% identical to the DNA sequence of the nucleic acid fragments reported herein. More preferred nucleic acid fragments are at least 90% identical to the DNA sequence of the nucleic acid fragments reported herein. Even more preferred are nucleic acid fragments that are at least 95% identical to the DNA sequence of the nucleic acid fragments reported herein.

[0113] Two amino acid sequences are "substantially homologous" or "substantially similar" when greater than about 40% of the amino acids are identical, or greater than 60% are similar (functionally identical). Preferably, the similar or homologous sequences are identified by alignment using, for example, the GCG (Genetics Computer Group, Program Manual for the GCG Package, Version 7, Madison, Wisconsin) pileup program

[0114] The term "corresponding to" is used herein to refer to similar or homologous sequences, whether the exact position is identical or different from the molecule to which the similarity or homology is measured. A nucleic acid or amino acid sequence alignment may include spaces. Thus, the term "corresponding to" refers to the sequence similarity, and not the numbering of the amino acid residues or nucleotide bases.

[0115] A "substantial portion" of an amino acid or nucleotide sequence comprises enough of the amino acid sequence of a polypeptide or the nucleotide sequence of a gene to putatively identify that polypeptide or gene, either by manual evaluation of the sequence by one skilled in the art, or by computer-automated sequence comparison and identification using algorithms such as BLAST (Basic Local Alignment Search Tool; Altschul, S. F., et al., (1993) *J. Mol. Biol.* 215: 403-410; see also www.ncbi.nlm.nih.gov/BLAST/). In general, a sequence of ten or more contiguous amino acids or thirty or more nucleotides is necessary in order to putatively identify a polypeptide or nucleic acid sequence as homologous to a known protein or gene. Moreover, with respect to nucleotide sequences, gene specific oligonucleotide probes comprising 20-30 contiguous nucleotides may be used in sequence-dependent methods of gene identification (e.g., Southern hybridization) and isolation (e.g., *in situ* hybridization of bacterial colonies or bacteriophage plaques). In addition, short oligonucleotides of 12-15 bases may be used as amplification primers in PCR in order to obtain a particular nucleic acid fragment comprising the primers. Accordingly, a "substantial portion" of a nucleotide sequence comprises enough of the sequence to specifically identify and/or isolate a nucleic acid fragment comprising the sequence.

[0116] The term "percent identity", as known in the art, is a relationship between two or more polypeptide sequences or two or more polynucleotide sequences, as determined by comparing the sequences. In the art, "identity" also means the degree of sequence relatedness between polypeptide or polynucleotide sequences, as the case may be, as determined by the match between strings of such sequences. "Identity" and "similarity" can be readily calculated by known methods, including but not limited to those described in: *Computational Molecular Biology* (Lesk, A. M., ed.) Oxford University Press, New York (1988); *Biocomputing: Informatics and Genome Projects* (Smith, D. W., ed.) Academic Press, New York (1993); *Computer Analysis of Sequence Data, Part I* (Griffin, A. M., and Griffin, H. G., eds.) Humana Press, New Jersey (1994); *Sequence Analysis in Molecular Biology* (von Heinje, G., ed.) Academic Press (1987); and *Sequence Analysis Primer* (Gribskov, M. and Devereux, J., eds.) Stockton Press, New York (1991). Preferred methods to determine identity are designed to give the best match between the sequences tested. Methods to determine identity and similarity are codified in publicly available computer programs. Sequence alignments and percent identity calculations may be performed using the Megalign program of the LASERGENE bioinformatics computing suite (DNASTAR Inc., Madison, WI). Multiple alignment of the sequences may be performed using the Clustal method of alignment (Higgins

and Sharp (1989) *CABIOS*, 5:151-153) with the default parameters (GAP PENALTY=10, GAP LENGTH PENALTY=10). Default parameters for pairwise alignments using the Clustal method may be selected: KTUPLE 1, GAP PENALTY=3, WINDOW=5 and DIAGONALS SAVED=5.

[0117] The term "sequence analysis software" refers to any computer algorithm or software program that is useful for the analysis of nucleotide or amino acid sequences. "Sequence analysis software" may be commercially available or independently developed. Typical sequence analysis software will include but is not limited to the GCG suite of programs (Wisconsin Package Version 9.0, Genetics Computer Group (GCG), Madison, WI), BLASTP, BLASTN, BLASTX (Altschul et al., *J. Mol. Biol.* 215:403-410 (1990), and DNASTAR (DNASTAR, Inc. 1228 S. Park St. Madison, WI 53715 USA). Within the context of this application it will be understood that where sequence analysis software is used for analysis, that the results of the analysis will be based on the "default values" of the program referenced, unless otherwise specified. As used herein "default values" will mean any set of values or parameters which originally load with the software when first initialized.

[0118] "Synthetic genes" can be assembled from oligonucleotide building blocks that are chemically synthesized using procedures known to those skilled in the art. These building blocks are ligated and annealed to form gene segments that are then enzymatically assembled to construct the entire gene. "Chemically synthesized", as related to a sequence of DNA, means that the component nucleotides were assembled *in vitro*. Manual chemical synthesis of DNA may be accomplished using well established procedures, or automated chemical synthesis can be performed using one of a number of commercially available machines. Accordingly, the genes can be tailored for optimal gene expression based on optimization of nucleotide sequence to reflect the codon bias of the host cell. The skilled artisan appreciates the likelihood of successful gene expression if codon usage is biased towards those codons favored by the host. Determination of preferred codons can be based on a survey of genes derived from the host cell where sequence information is available.

GENE EXPRESSION MODULATION SYSTEM OF THE INVENTION

[0119] Applicants have now shown that separating the transactivation and DNA binding domains by placing them on two different proteins results in greatly reduced background activity in the absence of a ligand and significantly increased activity over background in the presence of a ligand. Applicants' improved gene expression system comprises two chimeric gene expression; the first encoding a DNA binding domain fused to a nuclear receptor polypeptide and the second encoding a transactivation domain fused to a nuclear receptor polypeptide. The interaction of the first protein with the second protein effectively tethers the DNA binding domain to the transactivation domain. Since the DNA binding and transactivation domains reside on two different molecules, the background activity in the absence of ligand is greatly reduced.

[0120] In general, the inducible gene expression modulation system of the invention comprises a) a first chimeric gene that is capable of being expressed in a host cell comprising a polynucleotide sequence that encodes a first hybrid polypeptide comprising i) a DNA binding domain that recognizes a response element associated with a gene whose expression is to be modulated; and ii) a ligand binding domain comprising the ligand binding domain from a nuclear receptor, and b) a second chimeric gene that is capable of being expressed in the host cell comprising a polynucleotide sequence that encodes a second hybrid polypeptide comprising: i) a transactivation domain; and ii) a ligand binding domain comprising the ligand binding domain from a nuclear receptor other than ultraspiracle (USP); wherein the transactivation domain are from other than EcR, RXR, or USP; and wherein the ligand binding domains from the first hybrid polypeptide and the second hybrid polypeptide are different and dimerize.

[0121] This two-hybrid system exploits the ability of a pair of interacting proteins to bring the transcription activation domain into a more favorable position relative to the DNA binding domain such that when the DNA binding domain binds to the DNA binding site on the gene, the transactivation domain more effectively activates the promoter (see, for example, U.S. Patent No. 5,283,173). This two-hybrid system is a significantly improved inducible gene expression modulation system compared to the two systems disclosed in International Patent Applications PCT/US97/05330 and PCT/US98/14215.

[0122] The ecdysone receptor-based gene expression modulation system of the invention may be either heterodimeric and homodimeric. A functional EcR complex generally refers to a heterodimeric protein complex consisting of two members of the steroid receptor family, an ecdysone receptor protein obtained from various insects, and an ultraspiracle (USP) protein or the vertebrate homolog of USP, retinoid X receptor protein (see Yao, et al. (1993) *Nature* 366, 476-479; Yao, et al., (1992) *Cell* 71, 63-72). However, the complex may also be a homodimer as detailed below. The functional ecdysteroid receptor complex may also include additional protein(s) such as immunophilins. Additional members of the steroid receptor family of proteins, known as transcriptional factors (such as DHR38 or *betaFTZ-1*), may also be ligand dependent or independent partners for EcR, USP, and/or RXR. Additionally, other cofactors may be required such as proteins generally known as coactivators (also termed adapters or mediators). These proteins do not bind sequence-specifically to DNA and are not involved in basal transcription. They may exert their effect on transcription activation

through various mechanisms, including stimulation of DNA-binding of activators, by affecting chromatin structure, or by mediating activator-initiation complex interactions. Examples of such coactivators include RIP140, TIF1, RAP46Bag-1, ARA70, SRC-1/NCoA-1, TIF2/GRIP/NCoA-2, ACTR/AIB 1/RAC3/pCIP as well as the promiscuous co activator C response element B binding protein, CBP/p300 (for review see Glass et al, Curr. Opin. Cell Biol. 9:222-232, 1997). Also, protein cofactors generally known as corepressors (also known as repressors, silencers, or silencing mediators) may be required to effectively inhibit transcriptional activation in the absence of ligand. These corepressors may interact with the unliganded ecdysone receptor to silence the activity at the response element. Current evidence suggests that binding of ligand changes the conformation of the receptor, which results in release of the corepressor and recruitment of the above described coactivators, thereby, abolishing their silencing activity. Examples of corepressors include N-CoR and SMRT (for review, see Horwitz et al. Mol Endocrinol. 10: 1167-1177, 1996). These cofactors may either be endogenous within the cell or organism, or may be added exogenously as transgenes to be expressed in either a regulated or unregulated fashion. Homodimer complexes of the ecdysone receptor protein, USP, or RXR may also be functional under some circumstances.

[0123] The ecdysone receptor complex typically includes proteins which are members of the nuclear receptor superfamily wherein all members are characterized by the presence of an amino-terminal transactivation domain, a DNA binding domain ("DBD"), and a ligand binding domain ("LBD") separated from the DBD by a hinge region. As used herein, the term "DNA binding domain" comprises a minimal peptide sequence of a DNA binding protein, up to the entire length of a DNA binding protein, so long as the DNA binding domain functions to associate with a particular response element. Members of the nuclear receptor superfamily are also characterized by the presence of four or five domains: A/B, C, D, E, and in some members F (see Evans, *Science* 240:889-895 (1988)). The "AB" domain corresponds to the transactivation domain, "C" corresponds to the DNA binding domain, "D" corresponds to the hinge region, and "E" corresponds to the ligand binding domain. Some members of the family may also have another transactivation domain on the carboxy-terminal side of the LBD corresponding to "F".

[0124] The DBD is characterized by the presence of two cysteine zinc fingers between which are two amino acid motifs, the P-box and the D-box, which confer specificity for ecdysone response elements. These domains may be either native, modified, or chimeras of different domains of heterologous receptor proteins. This EcR receptor, like a subset of the steroid receptor family, also possesses less well defined regions responsible for heterodimerization properties. Because the domains of EcR, USP, and RXR are modular in nature, the LBD, DBD, and transactivation domains may be interchanged.

[0125] Gene switch systems are known that incorporate components from the ecdysone receptor complex. However, in these known systems, whenever EcR is used it is associated with native or modified DNA binding domains and transactivation domains on the same molecule. USP or RXR are typically used as silent partners. We have now shown that when DNA binding domains and transactivation domains are on the same molecule the background activity in the absence of ligand is high and that such activity is dramatically reduced when DNA binding domains and transactivation domains are on different molecules, that is, on each of two partners of a heterodimeric or homodimeric complex. This two-hybrid system also provides improved sensitivity to non-steroidal ligands for example, diacylhydrazines, when compared to steroidal ligands for example, ponasterone A ("PonA") or muristerone A ("MurA"). That is, when compared to steroids, the non-steroidal ligands provide higher activity at a lower concentration. In addition, since transactivation based on EcR gene switches is often cell-line dependent, it is easier to tailor switching system to obtain maximum transactivation capability for each application. Furthermore, this two-hybrid system avoids some side effects due to overexpression of RXR that often occur when unmodified RXR is used as a switching partner. In this two-hybrid system, native DNA binding and transactivation domains of EcR or RXR are eliminated. As a result, these chimeric molecules have less chance of interacting with other steroid hormone receptors present in the cell resulting in reduced side effects.

[0126] Specifically, Applicants' invention relates to a gene expression modulation system comprising: a) a first gene expression cassette that is capable of being expressed in a host cell, wherein the first gene expression cassette comprises a polynucleotide that encodes a first polypeptide comprising i) a DNA-binding domain that recognizes a response element associated with a gene whose expression is to be modulated; and ii) a ligand binding domain comprising a ligand binding domain from a nuclear receptor; and b) a second gene expression cassette that is capable of being expressed in the host cell, wherein the second gene expression cassette comprises a polynucleotide sequence that encodes a second polypeptide comprising i) a transactivation domain; and ii) a ligand binding domain comprising a ligand binding domain from a nuclear receptor other than ultraspiracle (USP); wherein the DNA binding domain and the transactivation domain are from other than EcR, RXR, or USP; wherein the ligand binding domains from the first polypeptide and the second polypeptide are different and dimerize.

[0127] The present invention also relates to a gene expression modulation system according to the present invention further comprising c) a third gene expression cassette comprising: i) the response element to which the DNA-binding domain of the first polypeptide binds; ii) a promoter that is activated by the transactivation domain of the second polypeptide; and iii) the gene whose expression is to be modulated.

[0128] In a specific embodiment, the gene whose expression is to be modulated is a homologous gene with respect

to the host cell. In another specific embodiment, the gene whose expression is to be modulated is a heterologous gene with respect to the host cell.

[0129] In a specific embodiment, the ligand binding domain of the first polypeptide comprises an ecdysone receptor ligand binding domain.

[0130] In another specific embodiment, the ligand binding domain of the first polypeptide comprises a retinoid X receptor ligand binding domain.

[0131] In a specific embodiment, the ligand binding domain of the second polypeptide comprises an ecdysone receptor ligand binding domain.

[0132] In another specific embodiment, the ligand binding domain of the second polypeptide comprises a retinoid X receptor ligand binding domain.

[0133] In a preferred embodiment, the ligand binding domain of the first polypeptide comprises an ecdysone receptor ligand binding domain, and the ligand binding domain of the second polypeptide comprises a retinoid X receptor ligand binding domain.

[0134] In another preferred embodiment, the ligand binding domain of the first polypeptide is from a retinoid X receptor polypeptide, and the ligand binding domain of the second polypeptide is from an ecdysone receptor polypeptide.

[0135] Preferably, the ligand binding domain is an EcR or RXR related steroid/thyroid hormone nuclear receptor family member ligand binding domain, or analogs, combinations, or modifications thereof. More preferably, the LBD is from EcR or RXR. Even more preferably, the LBD is from a truncated EcR or RXR. A truncation mutation may be made by any method used in the art, including but not limited to restriction endonuclease digestion/deletion, PCR-mediated/oligonucleotide-directed deletion, chemical mutagenesis, UV strand breakage, and the like.

[0136] Preferably, the EcR is an insect EcR selected from the group consisting of a Lepidopteran EcR, a Dipteran EcR, an Arthropod EcR, a Homopteran EcR and a Hemipteran EcR. More preferably, the EcR for use is a spruce budworm *Choristoneura fumiferana* EcR ("CfEcR"), a *Tenebrio molitor* EcR ("TmEcR"), a *Manduca sexta* EcR ("MsEcR"), a *Heliothis virescens* EcR ("HvEcR"), a silk moth *Bombyx mori* EcR ("BmEcR"), a fruit fly *Drosophila melanogaster* EcR ("DmEcR"), a mosquito *Aedes aegypti* EcR ("AaEcR"), a blowfly *Lucilia capitata* EcR ("LcEcR"), a Mediterranean fruit fly *Ceratitis capitata* EcR ("CcEcR"), a locust *Locusta migratoria* EcR ("LmEcR"), an aphid *Myzus persicae* EcR ("MpEcR"), a fiddler crab *Uca pugilator* EcR ("UpEcR"), or an ixodid tick *Amblyomma americanum* EcR ("AmaEcR"). Even more preferably, the LBD is from spruce budworm (*Choristoneura fumiferana*) EcR ("CfEcR") or fruit fly *Drosophila melanogaster* EcR ("DmEcR").

[0137] Preferably, the LBD is from a truncated insect EcR. The insect EcR polypeptide truncation comprises a deletion of at least 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 105, 110, 115, 120, 125, 130, 135, 140, 145, 150, 155, 160, 165, 170, 175, 180, 185, 190, 195, 200, 205, 210, 215, 220, 225, 230, 235, 240, 245, 250, 255, 260, or 265 amino acids. More preferably, the insect EcR polypeptide truncation comprises a deletion of at least a partial polypeptide domain. Even more preferably, the insect EcR polypeptide truncation comprises a deletion of at least an entire polypeptide domain. In a specific embodiment, the insect EcR polypeptide truncation comprises a deletion of at least an A/B-domain deletion, a C-domain deletion, a D-domain deletion, an E-domain deletion, an F-domain deletion, an A/B/C-domains deletion, an AB/1/2-C-domains deletion, an A/B/C/D-domains deletion, an AB/C/D/F-domains deletion, an A/B/F-domains, and an A/B/C/F-domains deletion. A combination of several complete and/or partial domain deletions may also be performed.

[0138] In a preferred embodiment, the ecdysone receptor ligand binding domain is encoded by a polynucleotide comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, and SEQ ID NO: 10.

[0139] In another preferred embodiment, the ecdysone receptor ligand binding domain comprises a polypeptide sequence selected from the group consisting of SEQ ID NO: 11, SEQ ID NO: 12, SEQ ID NO: 13, SEQ ID NO: 14, SEQ ID NO: 15, SEQ ID NO: 16, SEQ ID NO: 17, SEQ ID NO: 18, SEQ ID NO: 19, and SEQ ID NO: 20.

[0140] Preferably, the RXR polypeptide is a mouse *Mus musculus* RXR ("MmRXR") or a human *Homo sapiens* RXR ("HsRXR"). The RXR polypeptide may be an RXR_α, RXR_β, or RXR_γ isoform.

[0141] Preferably, the LBD is from a truncated RXR. The RXR polypeptide truncation comprises a deletion of at least 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 105, 110, 115, 120, 125, 130, 135, 140, 145, 150, 155, 160, 165, 170, 175, 180, 185, 190, 195, 200, 205, 210, 215, 220, 225, 230, 235, 240, 245, 250, 255, 260, or 265 amino acids. More preferably, the RXR polypeptide truncation comprises a deletion of at least a partial polypeptide domain. Even more preferably, the RXR polypeptide truncation comprises a deletion of at least an entire polypeptide domain. In a specific embodiment, the RXR polypeptide truncation comprises a deletion of at least an A/B-domain deletion, a C-domain deletion, a D-domain deletion, an E-domain deletion, an F-domain - deletion, an A/B/C-domains deletion, an AB/1/2-C-domains deletion, an A/B/C/D-domains deletion, an A/B/C/D/F-domains deletion, an A/B/F-domains, and an A/B/C/F-domains deletion. A combination of several complete and/or partial domain deletions may also be performed.

[0142] In a preferred embodiment, the retinoid X receptor ligand binding domain is encoded by a polynucleotide

comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO: 21, SEQ ID NO: 22, SEQ ID NO: 23, SEQ ID NO: 24, SEQ ID NO: 25, SEQ ID NO: 26, SEQ ID NO: 27, SEQ ID NO: 28, SEQ ID NO: 29, and SEQ ID NO: 30.

[0143] In another preferred embodiment, the retinoid X receptor ligand binding domain comprises a polypeptide sequence selected from the group consisting of SEQ ID NO: 31, SEQ ID NO: 32, SEQ ID NO: 33, SEQ ID NO: 34, SEQ ID NO: 35, SEQ ID NO: 36, SEQ ID NO: 37, SEQ ID NO: 38, SEQ ID NO: 39, and SEQ ID NO: 40.

[0144] For purposes of this invention EcR and RXR also include synthetic and chimeric EcR and RXR and their homologs.

[0145] The DNA binding domain can be any DNA binding domain with a known response element, including synthetic and chimeric DNA binding domains, or analogs, combinations, or modifications thereof. Preferably, the DBD is a GAL4 DBD, a LexA DBD, a transcription factor DBD, a steroid/thyroid hormone nuclear receptor superfamily member DBD, a bacterial LacZ DBD, or a yeast put DBD. More preferably, the DBD is a GAL4 DBD [SEQ ID NO: 41 (polynucleotide) or SEQ ID NO: 42 (polypeptide)] or a LexA DBD [(SEQ ID NO: 43 (polynucleotide) or SEQ ID NO: 44 (polypeptide)].

[0146] The transactivation domain (abbreviated "AD" or "TA") may be any steroid/thyroid hormone nuclear receptor AD, synthetic or chimeric AD, polyglutamine AD, basic or acidic amino acid AD, a VP16 AD, a GAL4 AD, an NF- κ B AD, a BP64 AD, or an analog, combination, or modification thereof. Preferably, the AD is a synthetic or chimeric AD, or is obtained from a VP16, GAL4, or NF- κ B. Most preferably, the AD is a VP16 AD [SEQ ID NO: 45 (polynucleotide) or SEQ ID NO: 46 (polypeptide)].

[0147] The response element ("RE") may be any response element with a known DNA binding domain, or an analog, combination, or modification thereof. Preferably, the RE is an RE from GAL4 ("GAL4RE"), LexA, a steroid/thyroid hormone nuclear receptor RE, or a synthetic RE that recognizes a synthetic DNA binding domain. More preferably, the RE is a GAL4RE comprising a polynucleotide sequence of SEQ ID NO: 47 or a LexA 8X operon comprising a polynucleotide sequence of SEQ ID NO: 48. Preferably, the first hybrid protein is substantially free of a transactivation domain and the second hybrid protein is substantially free of a DNA binding domain. For purposes of this invention, "substantially free" means that the protein in question does not contain a sufficient sequence of the domain in question to provide activation or binding activity.

[0148] The ligands for use in the present invention as described below, when combined with the ligand binding domain of an EcR, USP, RXR, or another polypeptide which in turn are bound to the response element linked to a gene, provide the means for external temporal regulation of expression of the gene. The binding mechanism or the order in which the various components of this invention bind to each other, that is, ligand to receptor, first polypeptide to response element, second polypeptide to promoter, etc., is not critical. Binding of the ligand to the ligand binding domains of an EcR, USP, RXR, or another protein, enables expression or suppression of the gene. This mechanism does not exclude the potential for ligand binding to EcR, USP, or RXR, and the resulting formation of active homodimer complexes (e.g. EcR+EcR or USP+USP). Preferably, one or more of the receptor domains can be varied producing a chimeric gene switch. Typically, one or more of the three domains, DBD, LBD, and transactivation domain, may be chosen from a source different than the source of the other domains so that the chimeric genes and the resulting hybrid proteins are optimized in the chosen host cell or organism for transactivating activity, complementary binding of the ligand, and recognition of a specific response element. In addition, the response element itself can be modified or substituted with response elements for other DNA binding protein domains such as the GAL-4 protein from yeast (see Sadowski, et al. (1988) Nature, 335: 563-564) or LexA protein from *E. coli* (see Brent and Ptashne (1985), Cell, 43:729-736), or synthetic response elements specific for targeted interactions with proteins designed, modified, and selected for such specific interactions (see, for example, Kim, et al. (1997), *Proc. Natl. Acad. Sci., USA*, 94:3616-3620) to accommodate chimeric receptors. Another advantage of chimeric systems is that they allow choice of a promoter used to drive the gene expression according to a desired end result. Such double control can be particularly important in areas of gene therapy, especially when cytotoxic proteins are produced, because both the timing of expression as well as the cells wherein expression occurs can be controlled. When genes, operatively linked to a suitable promoter, are introduced into the cells of the subject, expression of the exogenous genes is controlled by the presence of the system of this invention. Promoters may be constitutively or inducibly regulated or may be tissue-specific (that is, expressed only in a particular type of cells) or specific to certain developmental stages of the organism.

GENE EXPRESSION CASSETTES OF THE INVENTION

[0149] The novel ecdysone receptor-based inducible gene expression system of the invention comprises a novel gene expression cassette that is capable of being expressed in a host cell, wherein the gene expression cassette comprises a polynucleotide encoding a hybrid polypeptide. Thus, Applicants' invention also provides novel gene expression cassettes for use in the gene expression system of the invention.

[0150] Specifically, the present invention provides a gene expression cassette comprising a polynucleotide encoding a hybrid polypeptide. The hybrid polypeptide comprises either 1) a DNA-binding domain that recognizes a response element and a ligand binding domain of a nuclear receptor or 2) a transactivation domain and a ligand binding domain

of a nuclear receptor, wherein the transactivation domain is from a nuclear receptor other than an EcR, an RXR, or a USP.

[0151] In a specific embodiment, the gene expression cassette encodes a hybrid polypeptide comprising a DNA-binding domain that recognizes a response element and an ecdysone receptor ligand binding domain, wherein the DNA binding domain is from a nuclear receptor other than an ecdysone receptor.

[0152] In another specific embodiment, the gene expression cassette encodes a hybrid polypeptide comprising a DNA binding domain that recognizes a response element and a retinoid X receptor ligand binding domain, wherein the DNA binding domain is from a nuclear receptor other than a retinoid X receptor.

[0153] The DNA binding domain can be any DNA binding domain with a known response element, including synthetic and chimeric DNA binding domains, or analogs, combinations, or modifications thereof. Preferably, the DBD is a GAL4 DBD, a LexA DBD, a transcription factor DBD, a steroid/thyroid hormone nuclear receptor superfamily member DBD, a bacterial LacZ DBD, or a yeast put DBD. More preferably, the DBD is a GAL4 DBD [SEQ ID NO: 41 (polynucleotide) or SEQ ID NO: 42 (polypeptide)] or a LexA DBD [(SEQ ID NO: 43 (polynucleotide) or SEQ ID NO: 44 (polypeptide)].

[0154] In another specific embodiment, the gene expression cassette encodes a hybrid polypeptide comprising a transactivation domain and an ecdysone receptor ligand binding domain, wherein the transactivation domain is from a nuclear receptor other than an ecdysone receptor.

[0155] In another specific embodiment, the gene expression cassette encodes a hybrid polypeptide comprising a transactivation domain and a retinoid X receptor ligand binding domain, wherein the transactivation domain is from a nuclear receptor other than a retinoid X receptor.

[0156] The transactivation domain (abbreviated "AD" or "TA") may be any steroid/thyroid hormone nuclear receptor AD, synthetic or chimeric AD, polyglutamine AD, basic or acidic amino acid AD, a VP16 AD, a GAL4 AD, an NF- κ B AD, a BP64 AD, or an analog, combination, or modification thereof. Preferably, the AD is a synthetic or chimeric AD, or is obtained from a VP16, GAL4, or NF- κ B. Most preferably, the AD is a VP16 AD [SEQ ID NO: 45 (polynucleotide) or SEQ ID NO: 46 (polypeptide)].

[0157] Preferably, the ligand binding domain is an EcR or RXR related steroid/thyroid hormone nuclear receptor family member ligand binding domain, or analogs, combinations, or modifications thereof. More preferably, the LBD is from EcR or RXR. Even more preferably, the LBD is from a truncated EcR or RXR.

[0158] Preferably, the EcR is an insect EcR selected from the group consisting of a Lepidopteran EcR, a Dipteran EcR, an Arthropod EcR, a Homopteran EcR and a Hemipteran EcR. More preferably, the EcR for use is a spruce budworm *Choristoneura fumiferana* EcR ("CfEcR"), a *Tenebrio molitor* EcR ("TmEcR"), a *Manduca sexta* EcR ("MsEcR"), a *Heliothis virescens* EcR ("HvEcR"), a silk moth *Bombyx mori* EcR ("BmEcR"), a fruit fly *Drosophila melanogaster* EcR ("DmEcR"), a mosquito *Aedes aegypti* EcR ("AaEcR"), a blowfly *Lucilia capitata* EcR ("LcEcR"), a Mediterranean fruit fly *Ceratitis capitata* EcR ("CcEcR"), a locust *Locusta migratoria* EcR ("LmEcR"), an aphid *Myzus persicae* EcR ("MpEcR"), a fiddler crab *Uca pugilator* EcR ("UpEcR"), or an ixodid tick *Amblyomma americanum* EcR ("AmaEcR"). Even more preferably, the LBD is from spruce budworm (*Choristoneura fumiferana*) EcR ("CfEcR") or fruit fly *Drosophila melanogaster* EcR ("DmEcR").

[0159] Preferably, the LBD is from a truncated insect EcR. The insect EcR polypeptide truncation comprises a deletion of at least 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 105, 110, 115, 120, 125, 130, 135, 140, 145, 150, 155, 160, 165, 170, 175, 180, 195, 190, 195, 200, 205, 210, 215, 220, 225, 230, 235, 240, 245, 250, 255, 260, or 265 amino acids. More preferably, the insect EcR polypeptide truncation comprises a deletion of at least a partial polypeptide domain. Even more preferably, the insect EcR polypeptide truncation comprises a deletion of at least an entire polypeptide domain. In a specific embodiment, the insect EcR polypeptide truncation comprises a deletion of at least an A/B-domain deletion, a C-domain deletion, a D-domain deletion, an E-domain deletion, an F-domain deletion, an A/B/C-domains deletion, an AB/1/2-C-domains deletion, an A/B/C/D-domains deletion, an A/B/C/D/F-domains deletion, an A/B/F-domains, and an A/B/C/F-domains deletion. A combination of several complete and/or partial domain deletions may also be performed.

[0160] In a preferred embodiment, the ecdysone receptor ligand binding domain is encoded by a polynucleotide comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, and SEQ ID NO: 10.

[0161] In another preferred embodiment, the ecdysone receptor ligand binding domain comprises an amino acid sequence selected from the group consisting of SEQ ID NO: 11, SEQ ID NO: 12, SEQ ID NO: 13, SEQ ID NO: 14, SEQ ID NO: 15, SEQ ID NO: 16, SEQ ID NO: 17, SEQ ID NO: 18, SEQ ID NO: 19, and SEQ ID NO: 20.

[0162] Preferably, the RXR polypeptide is a mouse *Mus musculus* RXR ("MmRXR") or a human *Homo sapiens* RXR ("HsRXR"). The RXR polypeptide may be an RXR $_{\alpha}$, RXR $_{\beta}$, or RXR $_{\gamma}$ isoform.

[0163] Preferably, the LBD is from a truncated RXR. The RXR polypeptide truncation comprises a deletion of at least 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 105, 110, 115, 120, 125, 130, 135, 140, 145, 150, 155, 160, 165, 170, 175, 180, 185, 190, 195, 200, 205, 210, 215, 220, 225, 230, 235, 240, 245, 250, 255, 260, or 265 amino acids. More preferably, the RXR polypeptide truncation comprises a deletion of at least a partial polypeptide domain. Even more preferably, the RXR polypeptide truncation comprises a deletion of at least an entire

polypeptide domain. In a specific embodiment, the RXR polypeptide truncation comprises a deletion of at least an A/B-domain deletion, a C-domain deletion, a D-domain deletion, an E-domain deletion, an F-domain deletion, an A/B/C-domains deletion, an AB/1/2-C-domains deletion, an A/B/C/D-domains deletion, an A/B/C/D/P-domains deletion, an A/B/F-domains, and an A/B/C/F-domains deletion. A combination of several complete and/or partial domain deletions may also be performed.

[0164] In a preferred embodiment, the retinoid X receptor ligand binding domain is encoded by a polynucleotide comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO: 21, SEQ ID NO: 22, SEQ ID NO: 23, SEQ ID NO: 24, SEQ ID NO: 25, SEQ ID NO: 26, SEQ ID NO: 27, SEQ ID NO: 28, SEQ ID NO: 29, and SEQ ID NO: 30.

[0165] In another preferred embodiment, the retinoid X receptor ligand binding domain comprises an amino acid sequence selected from the group consisting of SEQ ID NO: 31, SEQ ID NO: 32, SEQ ID NO: 33, SEQ ID NO: 34, SEQ ID NO: 35, SEQ ID NO: 36, SEQ ID NO: 37, SEQ ID NO: 38, SEQ ID NO: 39, and SEQ ID NO: 40.

[0166] In a preferred embodiment, the gene expression cassette encodes a hybrid polypeptide comprising a DNA-binding domain encoded by a polynucleotide comprising a nucleic acid sequence selected from the group consisting of a GAL4 DBD (SEQ ID NO: 41) or a LexA DBD (SEQ ID NO: 43) and an ecdysone receptor ligand binding domain encoded by a polynucleotide comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, and SEQ ID NO: 10.

[0167] In another preferred embodiment, the gene expression cassette encodes a hybrid polypeptide comprising a DNA-binding domain comprising a polypeptide sequence selected from the group consisting of a GAL4 DBD (SEQ ID NO: 42) or a LexA DBD (SEQ ID NO: 44) and an ecdysone receptor ligand binding domain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 11, SEQ ID NO: 12, SEQ ID NO: 13, SEQ ID NO: 14, SEQ ID NO: 15, SEQ ID NO: 16, SEQ ID NO: 17, SEQ ID NO: 18, SEQ ID NO: 19, and SEQ ID NO: 20.

[0168] In another preferred embodiment, the gene expression cassette encodes a hybrid polypeptide comprising a DNA-binding domain encoded by a polynucleotide comprising a nucleic acid sequence selected from the group consisting of a GAL4 DBD (SEQ ID NO: 41) or a LexA DBD (SEQ ID NO: 43) and a retinoid X receptor ligand binding domain encoded by a polynucleotide comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO: 21, SEQ ID NO: 22, SEQ ID NO: 23, SEQ ID NO: 24, SEQ ID NO: 25, SEQ ID NO: 26, SEQ ID NO: 27, SEQ ID NO: 28, SEQ ID NO: 29, and SEQ ID NO: 30.

[0169] In another preferred embodiment, the gene expression cassette encodes a hybrid polypeptide comprising a DNA-binding domain comprising a polypeptide sequence selected from the group consisting of a GAL4 DBD (SEQ ID NO: 42) or a LexA DBD (SEQ ID NO: 44) and a retinoid X receptor ligand binding domain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 31, SEQ ID NO: 32, SEQ ID NO: 33, SEQ ID NO: 34, SEQ ID NO: 35, SEQ ID NO: 36, SEQ ID NO: 37, SEQ ID NO: 38, SEQ ID NO: 39, and SEQ ID NO: 40.

[0170] In another preferred embodiment, the gene expression cassette encodes a hybrid polypeptide comprising a transactivation domain encoded by a polynucleotide comprising a nucleic acid sequence of SEQ ID NO: 45 and an ecdysone receptor ligand binding domain encoded by a polynucleotide comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, and SEQ ID NO: 10.

[0171] In another preferred embodiment, the gene expression cassette encodes a hybrid polypeptide comprising a transactivation domain comprising a polypeptide sequence of SEQ ID NO: 46 and an ecdysone receptor ligand binding domain comprising a polypeptide sequence selected from the group consisting of SEQ ID NO: 11, SEQ ID NO: 12, SEQ ID NO: 13, SEQ ID NO: 14, SEQ ID NO: 15, SEQ ID NO: 16, SEQ ID NO: 17, SEQ ID NO: 18, SEQ ID NO: 19, and SEQ ID NO: 20.

[0172] In another preferred embodiment, the gene expression cassette encodes a hybrid polypeptide comprising a transactivation domain encoded by a polynucleotide comprising a nucleic acid sequence of SEQ ID NO: 45 and a retinoid X receptor ligand binding domain encoded by a polynucleotide comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO: 21, SEQ ID NO: 22, SEQ ID NO: 23, SEQ ID NO: 24, SEQ ID NO: 25, SEQ ID NO: 26, SEQ ID NO: 27, SEQ ID NO: 28, SEQ ID NO: 29, and SEQ ID NO: 30.

[0173] In another preferred embodiment, the gene expression cassette encodes a hybrid polypeptide comprising a transactivation domain comprising a polypeptide sequence of SEQ ID NO: 46 and a retinoid X receptor ligand binding domain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 31, SEQ ID NO: 32, SEQ ID NO: 33, SEQ ID NO: 34, SEQ ID NO: 35, SEQ ID NO: 36, SEQ ID NO: 37, SEQ ID NO: 38, SEQ ID NO: 39, and SEQ ID NO: 40.

[0174] For purposes of this invention EcR and RXR also include synthetic and chimeric EcR and RXR and their homologs.

POLYNUCLEOTIDES OF THE INVENTION

[0175] The novel ecdysone receptor-based inducible gene expression system of the invention comprises a gene expression cassette comprising a polynucleotide that encodes a truncated EcR or RXR polypeptide comprising a truncation mutation and is useful in methods of modulating the expression of a gene within a host cell.

[0176] Thus, the present invention also relates to a polynucleotide that encodes an EcR or RXR polypeptide comprising a truncation mutation. Specifically, the present invention relates to an isolated polynucleotide encoding an EcR or RXR polypeptide comprising a truncation mutation that affects ligand binding activity or ligand sensitivity.

[0177] Preferably, the truncation mutation results in a polynucleotide that encodes a truncated EcR polypeptide or a truncated RXR polypeptide comprising a deletion of at least 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 105, 110, 115, 120, 125, 130, 135, 140, 145, 150, 155, 160, 165, 170, 175, 180, 185, 190, 195, 200, 205, 210, 215, 220, 225, 230, 235, 240, 245, 250, 255, 260, or 265 amino acids. More preferably, the EcR or RXR polypeptide truncation comprises a deletion of at least a partial polypeptide domain. Even more preferably, the EcR or RXR polypeptide truncation comprises a deletion of at least an entire polypeptide domain. In a specific embodiment, the EcR or RXR polypeptide truncation comprises a deletion of at least an A/B-domain deletion, a C-domain deletion, a D-domain deletion, an E-domain deletion, an F-domain deletion, an A/B/C-domains deletion, an A/B/1/2-C-domains deletion, an A/B/C/D-domains deletion, an A/B/C/D/F-domains deletion, an A/B/F-domains, and an A/B/C/F-domains deletion. A combination of several complete and/or partial domain deletions may also be performed.

[0178] In a specific embodiment, the EcR polynucleotide according to the invention comprises a polynucleotide sequence selected from the group consisting of SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, and SEQ ID NO: 10. In a specific embodiment, the polynucleotide according to the invention encodes a ecdysone receptor polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 11 (CfEcR-CDEF), SEQ ID NO: 12 (CfEcR-1/2CDEF, which comprises the last 33 carboxy-terminal amino acids of C domain), SEQ ID NO: 13 (CfEcR-DEF), SEQ ID NO: 14 (CfEcR-EF), SEQ ID NO: 15 (CfEcR-DE), SEQ ID NO: 16 (DmEcR-CDEF), SEQ ID NO: 17 (DmEcR-1/2CDEF), SEQ ID NO: 18 (DmEcR-DEF), SEQ ID NO: 19 (DmEcR-EF), and SEQ ID NO: 20 (DmEcR-DE).

[0179] In another specific embodiment, the RXR polynucleotide according to the invention comprises a polynucleotide sequence selected from the group consisting of SEQ ID NO: 21, SEQ ID NO: 22, SEQ ID NO: 23, SEQ ID NO: 24, SEQ ID NO: 25, SEQ ID NO: 26, SEQ ID NO: 27, SEQ ID NO: 28, SEQ ID NO: 29, and SEQ ID NO: 30. In another specific embodiment, the polynucleotide according to the invention encodes a truncated RXR polypeptide comprising an amino acid sequence consisting of SEQ ID NO: 31 (MmRXR-CDEF), SEQ ID NO: 32 (MmRXR-DEF), SEQ ID NO: 33 (MmRXR-EF), SEQ ID NO: 34 (MmRXR-truncatedEF), SEQ ID NO: 35 (MmRXR-E), SEQ ID NO: 36 (HsRXR-CDEF), SEQ ID NO: 37 (HsRXR-DEF), SEQ ID NO: 38 (HsRXR-EF), SEQ ID NO: 39 (HsRXR-truncated EF), and SEQ ID NO: 40 (HsRXR-E).

[0180] In particular, the present invention relates to an isolated polynucleotide encoding an EcR or RXR polypeptide comprising a truncation mutation, wherein the mutation reduces ligand binding activity or ligand sensitivity of the EcR or RXR polypeptide. In a specific embodiment, the present invention relates to an isolated polynucleotide encoding an EcR or RXR polypeptide comprising a truncation mutation that reduces steroid binding activity or steroid sensitivity of the EcR or RXR polypeptide. In a preferred embodiment, the present invention relates to an isolated polynucleotide encoding an EcR polypeptide comprising a truncation mutation that reduces steroid binding activity or steroid sensitivity of the EcR polypeptide, wherein the polynucleotide comprises a nucleic acid sequence of SEQ ID NO: 3 (CfEcR-DEF), SEQ ID NO: 4 (CfEcR-EF), SEQ ID NO: 8 (DmEcR-DEF), or SEQ ID NO: 9 (DmEcR-EF). In another specific embodiment, the present invention relates to an isolated polynucleotide encoding an EcR or RXR polypeptide comprising a truncation mutation that reduces non-steroid binding activity or non-steroid sensitivity of the EcR or RXR polypeptide. In a preferred embodiment, the present invention relates to an isolated polynucleotide encoding an EcR polypeptide comprising a truncation mutation that reduces non-steroid binding activity or non-steroid sensitivity of the EcR polypeptide, wherein the polynucleotide comprises a nucleic acid sequence of SEQ ID NO: 4 (CfEcR-EF) or SEQ ID NO: 9 (DmEcR-EF).

[0181] The present invention also relates to an isolated polynucleotide encoding an EcR or RXR polypeptide comprising a truncation mutation, wherein the mutation enhances ligand binding activity or ligand sensitivity of the EcR or RXR polypeptide. In a specific embodiment, the present invention relates to an isolated polynucleotide encoding an EcR or RXR polypeptide comprising a truncation mutation that enhances steroid binding activity or steroid sensitivity of the EcR or RXR polypeptide. In another specific embodiment, the present invention relates to an isolated polynucleotide encoding an EcR or RXR polypeptide comprising a truncation mutation that enhances non-steroid binding activity or non-steroid sensitivity of the EcR or RXR polypeptide. In a preferred embodiment, the present invention relates to an isolated polynucleotide encoding an EcR polypeptide comprising a truncation mutation that enhances non-steroid binding activity or non-steroid sensitivity of the EcR polypeptide, wherein the polynucleotide comprises a nucleic acid sequence of SEQ ID NO: 3 (CfEcR-DEF) or SEQ ID NO: 8 (DmEcR-DEF).

[0182] The present invention also relates to an isolated polynucleotide encoding a retinoid X receptor polypeptide

comprising a truncation mutation that increases ligand sensitivity of a heterodimer comprising the mutated retinoid X receptor polypeptide and a dimerization partner. Preferably, the isolated polynucleotide encoding a retinoid X receptor polypeptide comprising a truncation mutation that increases ligand sensitivity of a heterodimer comprises a polynucleotide sequence selected from the group consisting of SEQ ID NO: 23 (MmRXR-EF), SEQ ID NO: 24 (114mRXR-truncatedEF),
 5 SEQ ID NO: 28 (HsRXR-EF), or SEQ ID NO: 29 (HsRXR-truncated EF). In a specific embodiment, the dimerization partner is an ecdysone receptor polypeptide. Preferably, the dimerization partner is a truncated EcR polypeptide. More preferably, the dimerization partner is an EcR polypeptide in which domains A/B/C have been deleted. Even more preferably, the dimerization partner is an EcR polypeptide comprising an amino acid sequence of SEQ ID NO: 13 (CfEcR-DEF) or SEQ ID NO: 18 (DmEcR-DEF).

POLYPEPTIDES OF THE INVENTION

[0183] The novel ecdysone receptor-based inducible gene expression system of the invention comprises a polynucleotide that encodes a truncated EcR or RXR polypeptide and is useful in methods of modulating the expression of a
 15 gene within a host cell. Thus, the present invention also relates to an isolated truncated EcR or RXR polypeptide encoded by a polynucleotide or a gene expression cassette according to the invention. Specifically, the present invention relates to an isolated truncated EcR or RXR polypeptide comprising a truncation mutation that affects ligand binding activity or ligand sensitivity encoded by a polynucleotide according to the invention.

[0184] The present invention also relates to an isolated truncated EcR or RXR polypeptide comprising a truncation
 20 mutation. Specifically, the present invention relates to an isolated EcR or RXR polypeptide comprising a truncation mutation that affects ligand binding activity or ligand sensitivity.

[0185] Preferably, the truncation mutation results in a truncated EcR polypeptide or a truncated RXR polypeptide comprising a deletion of at least 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 105, 110, 115, 120, 125, 130, 135, 140, 145, 150, 155, 160, 165, 170, 175, 180, 185, 190, 195, 200, 205, 210, 215, 220, 225,
 25 230, 235, 240, 245, 250, 255, 260, or 265 amino acids. More preferably, the EcR or RXR polypeptide truncation comprises a deletion of at least a partial polypeptide domain. Even more preferably, the EcR or RXR polypeptide truncation comprises a deletion of at least an entire polypeptide domain. In a specific embodiment, the EcR or RXR polypeptide truncation comprises a deletion of at least an A/B-domain deletion, a C-domain deletion, a D-domain deletion, an E-domain deletion, an F-domain deletion, an AB/C-domains deletion, an A/B/1/2-C-domains deletion, an AB/C/D-domains deletion, an
 30 AB/C/D/F-domains deletion, an A/B/F-domains, and an A/B/C/F-domains deletion. A combination of several complete and/or partial domain deletions may also be performed.

[0186] In a preferred embodiment, the isolated truncated ecdysone receptor polypeptide is encoded by a polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO: 1 (CfEcR-CDEF), SEQ ID NO: 2 (CfEcR-1/2CDEF), SEQ ID NO: 3 (CfEcR-DEF), SEQ ID NO: 4 (CfEcR-EF), SEQ ID NO: 5 (CfEcR-DE), SEQ ID NO: 6 (DmEcR-CDEF), SEQ ID NO: 7 (DmEcR-1/2CDEF), SEQ ID NO: 8 (DmEcR-DEF), SEQ ID NO: 9 (DmEcR-EF), and
 35 SEQ ID NO: 10 (DmEcR-DE). In another preferred embodiment, the isolated truncated ecdysone receptor polypeptide comprises an amino acid sequence selected from the group consisting of SEQ ID NO: 11 (CfEcR-CDEF), SEQ ID NO: 12 (CfEcR-1/2CDEF), SEQ ID NO: 13 (CfEcR-DEF), SEQ ID NO: 14 (CfEcR-EP), SEQ ID NO: 15 (CfEcR-DE), SEQ ID NO: 16 (DmEcR-CDEF), SEQ ID NO: 17 (DmEcR-1/2CDEF), SEQ ID NO: 18 (DmEcR-DEF), SEQ ID NO: 19 (DmEcR-EF), and SEQ ID NO: 20 (DmEcR-DE).

[0187] In a preferred embodiment, the isolated truncated RXR polypeptide is encoded by a polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO: 21 (MmRXR-CDEF), SEQ ID NO: 22 (MmRXR-DEF), SEQ ID NO: 23 (MmRXR-EF), SEQ ID NO: 24 (MmRXR-truncatedEF), SEQ ID NO: 25 (MmRXR-E),
 45 SEQ ID NO: 26 (HsRXR-CDEF), SEQ ID NO: 27 (HsRXR-DEF), SEQ ID NO: 28 (HsRXR-EF), SEQ ID NO: 29 (HsRXR-truncatedEF) and SEQ ID NO: 30 (HsRXR-E). In another preferred embodiment, the isolated truncated RXR polypeptide comprises an amino acid sequence selected from the group consisting of SEQ ID NO: 31 (MmRXR-CDEF), SEQ ID NO: 32 (MmRXR-DEF), SEQ ID NO: 33 (MmRXR-EF), SEQ ID NO: 34 (MmRXR-truncatedEF), SEQ ID NO: 35 (MmRXR-E), SEQ ID NO: 36 (HsRXR-CDEF), SEQ ID NO: 37 (HsRXR-DEF), SEQ ID NO: 38 (HsRXR-EF), SEQ ID NO: 39 (HsRXR-truncatedEF), and SEQ ID NO: 40 (HsRXR-E).

[0188] The present invention relates to an isolated EcR or RXR polypeptide comprising a truncation mutation that reduces ligand binding activity or ligand sensitivity of the EcR or RXR polypeptide, wherein the polypeptide is encoded by a polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO: 1 (CfEcR-CDEF), SEQ ID NO: 2 (CfEcR-1/2CDEF), SEQ ID NO: 3 (CfEcR-DEF), SEQ ID NO: 4 (CfEcR-EF), SEQ ID NO: 5 (CfEcR-DE),
 55 SEQ ID NO: 6 (DmEcR-CDEF), SEQ ID NO: 7 (DmEcR-1/2CDEF), SEQ ID NO: 8 (DmEcR-DEF), SEQ ID NO: 9 (DmEcR-EF), SEQ ID NO: 10 (DmEcR-DE), SEQ ID NO: 21 (MmRXR-CDEF), SEQ ID NO: 22 (MmRXR-DEF), SEQ ID NO: 23 (MmRXR-EF), SEQ ID NO: 24 (MmRXR-truncatedEF), SEQ ID NO: 25 (MmRXR-E), SEQ ID NO: 26 (HsRXR-CDEF), SEQ ID NO: 27 (HsRXR-DEF), SEQ ID NO: 28 (HsRXR-EF), SEQ ID NO: 29 (HsRXR-truncatedEF), and SEQ ID NO: 30 (HsRXR-E).

[0189] Thus, the present invention relates to an isolated truncated EcR or RXR polypeptide comprising a truncation mutation that reduces ligand binding activity or ligand sensitivity of the EcR or RXR polypeptide, wherein the polypeptide comprises an amino acid sequence selected from the group consisting of SEQ ID NO: 11 (CfEcR-CDEF), SEQ ID NO: 12 (CfEcR-1/2CDEF), SEQ ID NO: 13 (CfEcR-DEF), SEQ ID NO: 14 (CfEcR-EF), SEQ ID NO: 15 (CfEcR-DE), SEQ ID NO: 16 (DmEcR-CDEF), SEQ ID NO: 17 (DmEcR-1/2CDEF), SEQ ID NO: 18 (DmEcR-DEF), SEQ ID NO: 19 (DmEcR-EF), SEQ ID NO: 20 (DmEcR-DE), SEQ ID NO: 31 (MmRXR-CDEF), SEQ ID NO: 32 (MmRXR-DEF), SEQ ID NO: 33 (MmRXR-EF), SEQ ID NO: 34 (MmRXR-truncatedEF), SEQ ID NO: 35 (MmRXR-E), SEQ ID NO: 36 (HsRXR-CDEF), SEQ ID NO: 37 (HsRXR-DEF), SEQ ID NO: 38 (HsRXR-EF), SEQ ID NO: 39 (HsRXR-truncatedEF), and SEQ ID NO: 40 (HsRXR-E).

[0190] In a specific embodiment, the present invention relates to an isolated EcR or RXR polypeptide comprising a truncation mutation that reduces steroid binding activity or steroid sensitivity of the EcR or RXR polypeptide. In a preferred embodiment, the present invention relates to an isolated EcR polypeptide comprising a truncation mutation that reduces steroid binding activity or steroid sensitivity of the EcR polypeptide, wherein the EcR polypeptide is encoded by a polynucleotide comprising a nucleic acid sequence of SEQ ID NO: 3 (CfEcR-DEF), SEQ ID NO: 4 (CfEcR-EF), SEQ ID NO: 8 (DmEcR-DEF), or SEQ ID NO: 9 (DmEcR-EF). Accordingly, the present invention also relates to an isolated truncated EcR or RXR polypeptide comprising a truncation mutation that reduces steroid binding activity or steroid sensitivity of the EcR or RXR polypeptide. In a preferred embodiment, the present invention relates to an isolated EcR polypeptide comprising a truncation mutation that reduces steroid binding activity or steroid sensitivity of the EcR polypeptide, wherein the EcR polypeptide comprises an amino acid sequence of SEQ ID NO: 13 (CfEcR-DEF), SEQ ID NO: 14 (CfEcR-EF), SEQ ID NO: 18 (DmEcR-DEF), or SEQ ID NO: 19 (DmEcR-EF).

[0191] In another specific embodiment, the present invention relates to an isolated EcR or RXR polypeptide comprising a truncation mutation that reduces non-steroid binding activity or non-steroid sensitivity of the EcR or RXR polypeptide. In a preferred embodiment, the present invention relates to an isolated EcR polypeptide comprising a truncation mutation that reduces non-steroid binding activity or non-steroid sensitivity of the EcR polypeptide, wherein the EcR polypeptide is encoded by a polynucleotide comprising a nucleic acid sequence of SEQ ID NO: 4 (CfEcR-EF) or SEQ ID NO: 9 (DmEcR-EF). Accordingly, the present invention also relates to an isolated truncated EcR or RXR polypeptide comprising a truncation mutation that reduces non-steroid binding activity or steroid sensitivity of the EcR or RXR polypeptide. In a preferred embodiment, the present invention relates to an isolated EcR polypeptide comprising a truncation mutation that reduces non-steroid binding activity or non-steroid sensitivity of the EcR polypeptide, wherein the EcR polypeptide comprises an amino acid sequence of SEQ ID NO: 14 (CfEcR-EF) or SEQ ID NO: 19 (DmEcR-EF).

[0192] In particular, the present invention relates to an isolated EcR or RXR polypeptide comprising a truncation mutation that enhances ligand binding activity or ligand sensitivity of the EcR or RXR polypeptide, wherein the polypeptide is encoded by a polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO: 1 (CfEcR-CDEF), SEQ ID NO: 2 (CfEcR-1/2CDEF), SEQ ID NO: 3 (CfEcR-DEF), SEQ ID NO: 4 (CfEcR-EF), SEQ ID NO: 5 (CfEcR-DE), SEQ ID NO: 6 (DmEcR-CDEF), SEQ ID NO: 7 (DmEcR-1/2CDEF), SEQ ID NO: 8 (DmEcR-DEF), SEQ ID NO: 9 (DmEcR-EF), SEQ ID NO: 10 (DmEcR-DE), SEQ ID NO: 21 (MmRXR-CDEF), SEQ ID NO: 22 (MmRXR-DEF), SEQ ID NO: 23 (MmRXR-EF), SEQ ID NO: 24 (MmRXR-truncatedEF), SEQ ID NO: 25 (MmRXR-E), SEQ ID NO: 26 (HsRXR-CDEF), SEQ ID NO: 27 (HsRXR-DEF), SEQ ID NO: 28 (HsRXR-EF), SEQ ID NO: 29 (HsRXR-truncatedEF), and SEQ ID NO: 30 (HsRXR-E).

[0193] The present invention relates to an isolated EcR or RXR polypeptide comprising a truncation mutation that enhances ligand binding activity or ligand sensitivity of the EcR or RXR polypeptide, wherein the polypeptide comprises an amino acid sequence selected from the group consisting of SEQ ID NO: 11 (CfEcR-CDEF), SEQ ID NO: 12 (CfEcR-1/2CDEF), SEQ ID NO: 13 (CfEcR-DEF), SEQ ID NO: 14 (CfEcR-EF), SEQ ID NO: 15 (CfEcR-DE), SEQ ID NO: 16 (DmEcR-CDEF), SEQ ID NO: 17 (DmEcR-1/2CDEF), SEQ ID NO: 18 (DmEcR-DEF), SEQ ID NO: 19 (DmEcR-EF), SEQ ID NO: 20 (DmEcR-DE), SEQ ID NO: 31 (MmRXR-CDEF), SEQ ID NO: 32 (MmRXR-DEF), SEQ ID NO: 33 (MmRXR-EF), SEQ ID NO: 34 (MmRXR-truncatedEF), SEQ ID NO: 35 (MmRXR-E), SEQ ID NO: 36 (HsRXR-CDEF), SEQ ID NO: 37 (HsRXR-DEF), SEQ ID NO: 39 (HsRXR-EF), SEQ ID NO: 39 (HsRXR-truncatedEF), and SEQ ID NO: 40 (HsRXR-E).

[0194] The present invention relates to an isolated EcR or RXR polypeptide comprising a truncation mutation that enhances ligand binding activity or ligand sensitivity of the EcR or RXR polypeptide. In a specific embodiment, the present invention relates to an isolated EcR or RXR polypeptide comprising a truncation mutation that enhances steroid binding activity or steroid sensitivity of the EcR or RXR polypeptide. Accordingly, the present invention also relates to an isolated EcR or RXR polypeptide comprising a truncation mutation that enhances steroid binding activity or steroid sensitivity of the EcR or RXR polypeptide.

[0195] In another specific embodiment, the present invention relates to an isolated EcR or RXR polypeptide comprising a truncation mutation that enhances non-steroid binding activity or non-steroid sensitivity of the EcR or RXR polypeptide. In a preferred embodiment, the present invention relates to an isolated EcR polypeptide comprising a truncation mutation that enhances non-steroid binding activity or non-steroid sensitivity of the EcR polypeptide, wherein the EcR polypeptide

is encoded by a polynucleotide comprising a nucleic acid sequence of SEQ ID NO: 3 (CfEcR-DEF) or SEQ ID NO: 8 (DmEcR-DEF). Accordingly, the present invention also relates to an isolated EcR or RXR polypeptide comprising a truncation mutation that enhances non-steroid binding activity or steroid sensitivity of the EcR or RXR polypeptide. In a preferred embodiment, the present invention relates to an isolated EcR polypeptide comprising a truncation mutation that enhances non-steroid binding activity or non-steroid sensitivity of the EcR polypeptide, wherein the EcR polynucleotide comprises an amino acid sequence of SEQ ID NO: 13 (CfEcR-DEF) or SEQ ID NO: 18 (DmEcR-DEF).

[0196] The present invention also relates to an isolated retinoid X receptor polypeptide comprising a truncation mutation that increases ligand sensitivity of a heterodimer comprising the mutated retinoid X receptor polypeptide and a dimerization partner. Preferably, the isolated retinoid X receptor polypeptide comprising a truncation mutation that increases ligand sensitivity of a heterodimer is encoded by a polynucleotide comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO: 23 (MmRXR-EF), SEQ ID NO: 24 (MmRXR-truncatedEF), SEQ ID NO: 28 (HsRXR-EF), or SEQ ID NO: 29 (HsRXR-truncatedEF). More preferably, the isolated polynucleotide encoding a retinoid X receptor polypeptide comprising a truncation mutation that increases ligand sensitivity of a heterodimer comprises an amino acid sequence selected from the group consisting of SEQ ID NO: 33 (MmRXR-EF), SEQ ID NO: 34 (MmRXR-truncatedEF), SEQ ID NO: 38 (HsRXR-EF), or SEQ ID NO: 39 (HsRXR-truncatedEF).

[0197] In a specific embodiment, the dimerization partner is an ecdysone receptor polypeptide. Preferably, the dimerization partner is a truncated EcR polypeptide. More preferably, the dimerization partner is an EcR polypeptide in which domains A/B/C have been deleted. Even more preferably, the dimerization partner is an EcR polypeptide comprising an amino acid sequence of SEQ ID NO: 13 (CfEcR-DEF) or SEQ ID NO: 18 (DmEcR-DEF).

METHOD OF MODULATING GENE EXPRESSION OF THE INVENTION


[0198] Applicants' invention also relates to methods of modulating gene expression in a host cell using a gene expression modulation system according to the invention. Specifically, Applicants' invention provides a method of modulating the expression of a gene in a host cell comprising the steps of: a) introducing into the host cell a gene expression modulation system according to the invention; and b) introducing into the host cell a ligand that independently combines with the ligand binding domains of the first polypeptide and the second polypeptide of the gene expression modulation system; wherein the gene to be expressed is a component of a gene expression cassette comprising: i) a response element comprising a domain to which the DNA binding domain of the first polypeptide binds; ii) a promoter that is activated by the transactivation domain of the second polypeptide; and iii) a gene whose expression is to be modulated, whereby a complex is formed comprising the ligand, the first polypeptide of the gene expression modulation system and the second polypeptide of the gene expression modulation system, and whereby the complex modulates expression of the gene in the host cell.

[0199] Genes of interest for expression in a host cell using Applicants' methods may be endogenous genes or heterologous genes. Nucleic acid or amino acid sequence information for a desired gene or protein can be located in one of many public access databases, for example, GENBANK, EMBL, Swiss-Prot, and PIR, or in many biology related journal publications. Thus, those skilled in the art have access to nucleic acid sequence information for virtually all known genes. Such information can then be used to construct the desired constructs for the insertion of the gene of interest within the gene expression cassettes used in Applicants' methods described herein.

[0200] Examples of genes of interest for expression in a host cell using Applicants' methods include, but are not limited to: antigens produced in plants as vaccines, enzymes like alpha-amylase, phytase, glucanase, and xylanase, genes for resistance against insects, nematodes, fungi, bacteria, viruses, and abiotic stresses, nutraceuticals, pharmaceuticals, vitamins, genes for modifying amino acid content, herbicide resistance, cold, drought, and heat tolerance, industrial products, oils, protein, carbohydrates, antioxidants, male sterile plants, flowers, fuels, other output traits, genes encoding therapeutically desirable polypeptides or products, such as genes that can provide, modulate, alleviate, correct and/or restore polypeptides important in treating a condition, a disease, a disorder, a dysfunction, a genetic defect, and the like.

[0201] Acceptable ligands are any that modulate expression of the gene when binding of the DNA binding domain of the two hybrid system to the response element in the presence of the ligand results in activation or suppression of expression of the genes. Preferred ligands include ponasterone, muristerone A, N,N'-diacylhydrazines such as those disclosed in U. S. Patents No. 6,013,836; 5,117,057; 5,530,028; and 5,378,726; dibenzoylalkyl cyanohydrazines such as those disclosed in European Application No. 461,809; N-alkyl-N,N'-diarylhydrazines such as those disclosed in U. S. Patent No. 5,225,443; N-acyl-N-alkylcarbonylhydrazines such as those disclosed in European Application No. 234,994; N-aryl-N-alkyl-N'-arylhdydrazines such as those described in U. S. Patent No. 4,985,461; each of which is incorporated herein by reference and other similar materials including 3,5-di-tert-butyl-4-hydroxy-N-isobutyl-benzamide, 8-O-acetyl-harpagide, and the like.

[0202] Preferably, the ligand for use in Applicants' method of modulating expression of gene is a compound of the formula:



wherein:

- E is a (C₄-C₆)alkyl containing a tertiary carbon or a cyano(C₃-C₅)alkyl containing a tertiary carbon;
- R¹ is H, Me, Et, i-Pr, F, formyl, CF₃, CUF₂, CHCl₂, CH₂F, CH₂Cl, CH₂OH, CH₂OMe, CH₂CN, CN, C°CH, 1-propynyl, 2-propynyl, vinyl, OH, OMe, OEt, cyclopropyl, CF₂CF₃, CH=CHCN, allyl, azido, SCN, or SCHF₂;
- R² is H, Me, Et, n-Pr, i-Pr, formyl, CF₃, CHF₂, CHCl₂, CH₂F, CH₂Cl, CH₂OH, CH₂OMe, CH₂CN, CN, C°CH, 1-propynyl, 2-propynyl, vinyl, Ac, F, Cl, OH, OMe, OEt, O-n-Pr, OAc, NMe₂, NEt₂, SMe, SEt, SOCF₃, OCF₂CF₂H, COEt, cyclopropyl, CF₂CF₃, CH=CHCN, allyl, azido, OCF₃, OCHF₂, O-i-Pr, SCN, SCHF₂, SMe, NH-CN, or joined with R³ and the phenyl carbons to which R² and R³ are attached to form an ethylenedioxy, a dihydrofuryl ring with the oxygen adjacent to a phenyl carbon, or a dihydropyryl ring with the oxygen adjacent to a phenyl carbon;
- R³ is H, Et, or joined with R² and the phenyl carbons to which R² and R³ are attached to form an ethylenedioxy, a dihydrofuryl ring with the oxygen adjacent to a phenyl carbon, or a dihydropyryl ring with the oxygen adjacent to a phenyl carbon;
- R⁴, R⁵, and R⁶ are independently H, Me, Et, F, Cl, Br, formyl, CF₃, CHF₂, CHCl₂, CH₂F, CH₂Cl, CH₂OH, CN, C°CH, 1-propynyl, 2-propynyl, vinyl, OMe, OEt, SMe, or SEt.

[0203] Applicants' invention provides for modulation of gene expression in prokaryotic and eukaryotic host cells. Thus, the present invention also relates to a method for modulating gene expression in a host cell selected from the group consisting of a bacterial cell, a fungal cell, a yeast cell, a plant cell, an animal cell, and a mammalian cell. Preferably, the host cell is a yeast cell, a plant cell, a murine cell, or a human cell.

[0204] Expression in transgenic host cells may be useful for the expression of various polypeptides of interest including but not limited to therapeutic polypeptides, pathway intermediates; for the modulation of pathways already existing in the host for the synthesis of new products heretofore not possible using the host; cell based assays; and the like. Additionally the gene products may be useful for conferring higher growth yields of the host or for enabling alternative growth mode to be utilized.

HOST CELLS AND NON-HUMAN ORGANISMS OF THE INVENTION

[0205] As described above, the gene expression modulation system of the present invention may be used to modulate gene expression in a host cell. Expression in transgenic host cells may be useful for the expression of various genes of interest. Thus, Applicants' invention also provides an isolated host cell comprising a gene expression system according to the invention. The present invention also provides an isolated host cell comprising a gene expression cassette according to the invention. Applicants' invention also provides an isolated host cell comprising a polynucleotide or polypeptide according to the invention. The isolated host cell may be either a prokaryotic or a eukaryotic host cell.

[0206] Preferably, the host cell is selected from the group consisting of a bacterial cell, a fungal cell, a yeast cell, a plant cell, an animal cell, and a mammalian cell. Examples of preferred host cells include, but are not limited to, fungal or yeast species such as *Aspergillus*, *Trichoderma*, *Saccharomyces*, *Pichia*, *Candida*, *Hansenula*, or bacterial species such as those in the genera *Synechocystis*, *Synechococcus*, *Salmonella*, *Bacillus*, *Acinetobacter*, *Rhodococcus*, *Streptomyces*, *Escherichia*, *Pseudomonas*, *Methylobacter*, *Alcaligenes*, *Synechocystis*, *Anabaena*, *Thiobacillus*, *Methanobacterium* and *Klebsiella*, plant, animal, and mammalian host cells. More preferably, the host cell is a yeast cell, a plant cell, a mucine cell, or a human cell.

[0207] In a specific embodiment, the host cell is a yeast cell selected from the group consisting of a *Saccharomyces*, a *Pichia*, and a *Candida* host cell.

[0208] In another specific embodiment, the host cell is a plant cell selected from the group consisting of an apple, *Arabidopsis*, bajra, banana, barley, bean, beet, blackgram, chickpea, chili, cucumber, eggplant, favabean, maize, melon,

millet, mungbean, oat, okra, *Panicum*, papaya, peanut, pea, pepper, pigeonpea, pineapple, *Phaseolus*, potato, pumpkin, rice, sorghum, soybean, squash, sugarcane, sugarbeet, sunflower, sweet potato, tea, tomato, tobacco, watermelon, and wheat host cell.

[0209] In another specific embodiment, the host cell is a murine cell.

[0210] In another specific embodiment, the host cell is a human cell.

[0211] Host cell transformation is well known in the art and may be achieved by a variety of methods including but not limited to electroporation, viral infection, plasmid/vector transfection, non-viral vector mediated transfection, *Agrobacterium-mediated* transformation, particle bombardment, and the like. Expression of desired gene products involves culturing the transformed host cells under suitable conditions and inducing expression of the transformed gene. Culture conditions and gene expression protocols in prokaryotic and eukaryotic cells are well known in the art (see General Methods section of Examples). Cells may be harvested and the gene products isolated according to protocols specific for the gene product.

[0212] In addition, a host cell may be chosen which modulates the expression of the inserted polynucleotide, or modifies and processes the polypeptide product in the specific fashion desired. Different host cells have characteristic and specific mechanisms for the translational and post-translational processing and modification (e.g., glycosylation, cleavage [e.g., of signal sequence]) of proteins. Appropriate cell lines or host systems can be chosen to ensure the desired modification and processing of the foreign protein expressed. For example, expression in a bacterial system can be used to produce a non-glycosylated core protein product. However, a polypeptide expressed in bacteria may not be properly folded. Expression in yeast can produce a glycosylated product. Expression in eukaryotic cells can increase the likelihood of "native" glycosylation and folding of a heterologous protein. Moreover, expression in mammalian cells can provide a tool for reconstituting, or constituting, the polypeptide's activity. Furthermore, different vector/host expression systems may affect processing reactions, such as proteolytic cleavages, to a different extent.

[0213] Applicants' invention also relates to a non-human organism comprising an isolated host cell according to the invention. Preferably, the non-human organism is selected from the group consisting of a bacterium, a fungus, a yeast, a plant, an animal, and a mammal. More preferably, the non-human organism is a yeast, a plant, a mouse, a rat, a rabbit, a cat, a dog, a bovine, a goat, a pig, a horse, a sheep, a monkey, or a chimpanzee.

[0214] In a specific embodiment, the non-human organism is a yeast selected from the group consisting of *Saccharomyces*, *Pichia*, and *Candida*.

[0215] In another specific embodiment, the non-human organism is a plant selected from the group consisting of an apple, *Arabidopsis*, bajra, banana, barley, beans, beet, blackgram, chickpea, chili, cucumber, eggplant, favabean, maize, melon, millet, mungbean, oat, okra, *Panicum*, papaya, peanut, pea, pepper, pigeonpea, pineapple, *Phaseolus*, potato, pumpkin, rice, sorghum, soybean, squash, sugarcane, sugarbeet, sunflower, sweet potato, tea, tomato, tobacco, watermelon, and wheat.

[0216] In another specific embodiment, the non-human organism is a *Mus musculus* mouse.

MEASURING GENE EXPRESSION/TRANSCRIPTION

[0217] One useful measurement of Applicants' methods of the invention is that of the transcriptional state of the cell including the identities and abundances of RNA, preferably mRNA species. Such measurements are conveniently conducted by measuring cDNA abundances by any of several existing gene expression technologies.

[0218] Nucleic acid array technology is a useful technique for determining differential mRNA expression. Such technology includes, for example, oligonucleotide chips and DNA microarrays. These techniques rely on DNA fragments or oligonucleotides which correspond to different genes or cDNAs which are immobilized on a solid support and hybridized to probes prepared from total mRNA pools extracted from cells, tissues, or whole organisms and converted to cDNA. Oligonucleotide chips are arrays of oligonucleotides synthesized on a substrate using photolithographic techniques. Chips have been produced which can analyze for up to 1700 genes. DNA microarrays are arrays of DNA samples, typically PCR products, that are robotically printed onto a microscope slide. Each gene is analyzed by a full or partial-length target DNA sequence. Microarrays with up to 10,000 genes are now routinely prepared commercially. The primary difference between these two techniques is that oligonucleotide chips typically utilize 25-mer oligonucleotides which allow fractionation of short DNA molecules whereas the larger DNA targets of microarrays, approximately 1000 base pairs, may provide more sensitivity in fractionating complex DNA mixtures.

[0219] Another useful measurement of Applicants' methods of the invention is that of determining the translation state of the cell by measuring the abundances of the constituent protein species present in the cell using processes well known in the art.

[0220] Where identification of genes associated with various physiological functions is desired, an assay may be employed in which changes in such functions as cell growth, apoptosis, senescence, differentiation, adhesion, binding to a specific molecules, binding to another cell, cellular organization, organogenesis, intracellular transport, transport facilitation, energy conversion, metabolism, myogenesis, neurogenesis, and/or hematopoiesis is measured.

[0221] In addition, selectable marker or reporter gene expression may be used to measure gene expression modulation using Applicants' invention,

[0222] Other methods to detect the products of gene expression are well known in the art and include Southern blots (DNA detection), dot or slot blots (DNA, RNA), Northern blots (RNA), and RT-PCR (RNA) analyses. Although less preferred, labeled proteins can be used to detect a particular nucleic acid sequence to which it hybridizes.

[0223] In some cases it is necessary to amplify the amount of a nucleic acid sequence. This may be carried out using one or more of a number of suitable methods including, for example, polymerase chain reaction ("PCR"), ligase chain reaction ("LCR"), strand displacement amplification ("SDA"), transcription-based amplification, and the like. PCR is carried out in accordance with known techniques in which, for example, a nucleic acid sample is treated in the presence of a heat stable DNA polymerase, under hybridizing conditions, with one oligonucleotide primer for each strand of the specific sequence to be detected. An extension product of each primer that is synthesized is complementary to each of the two nucleic acid strands, with the primers sufficiently complementary to each strand of the specific sequence to hybridize therewith. The extension product synthesized from each primer can also serve as a template for further synthesis of extension products using the same primers. Following a sufficient number of rounds of synthesis of extension products, the sample may be analyzed as described above to assess whether the sequence or sequences to be detected are present.

[0224] The present invention may be better understood by reference to the following nonlimiting Examples, which are provided as exemplary of the invention.

EXAMPLES

GENERAL METHODS

[0225] Standard recombinant DNA and molecular cloning techniques used herein are well known in the art and are described by Sambrook, J., Fritsch, E. F. and Maniatis, T. *Molecular Cloning: A Laboratory Manual*; Cold Spring Harbor Laboratory Press: Cold Spring Harbor, (1989) (Maniatis) and by T. J. Silhavy, M. L. Bannan, and L. W. Enquist, *Experiments with Gene Fusions*, Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y. (1984) and by Ausubel, F. M. et al., *Current Protocols in Molecular Biology*, Greene Publishing Assoc. and Wiley-Interscience (1987).

[0226] Methods for plant tissue culture, transformation, plant molecular biology, and plant, general molecular biology may be found in *Plant Tissue Culture Concepts and Laboratory Exercises* edited by RN Trigiano and DJ Gray, 2nd edition, 2000, CRC press, New York; *Agrobacterium Protocols* edited by KMA Gartland and MR Davey, 1995, Humana Press, Totowa, New Jersey; *Methods in Plant Molecular Biology*, P. Maliga et al., 1995, Cold Spring Harbor Lab Press, New York; and *Molecular Cloning*, J. Sambrook et al., 1989, Cold Spring Harbor Lab Press, New York.

[0227] Materials and methods suitable for the maintenance and growth of bacterial cultures are well known in the art. Techniques suitable for use in the following examples may be found as set out in *Manual of Methods for General Bacteriology* (Phillipp Gerhardt, R. G. E. Murray, Ralph N. Costilow, Eugene W. Nester, Willis A. Wood, Noel R. Krieg and G. Briggs Phillips, eds), American Society for Microbiology, Washington, DC. (1994) or by Thomas D. Brock in *Biotechnology: A Textbook of Industrial Microbiology*, Second Edition, Sinauer Associates, Inc., Sunderland, MA (1989). All reagents, restriction enzymes and materials used for the growth and maintenance of host cells were obtained from Aldrich Chemicals (Milwaukee, WI), DIFCO Laboratories (Detroit, MI), GIBCO/BRL (Gaithersburg, MD), or Sigma Chemical Company (St. Louis, MO) unless otherwise specified.

[0228] Manipulations of genetic sequences may be accomplished using the suite of programs available from the Genetics Computer Group Inc. (Wisconsin Package Version 9.0, Genetics Computer Group (GCG), Madison, WI). Where the GCG program "Pileup" is used the gap creation default value of 12, and the gap extension default value of 4 may be used. Where the GCG "Gap" or "Bestfit" programs is used the default gap creation penalty of 50 and the default gap extension penalty of 3 may be used. In any case where GCG program parameters are not prompted for, in these or any other GCG program, default values may be used.

[0229] The meaning of abbreviations is as follows: "h" means hour(s), "min" means minute(s), "sec" means second(s), "d" means day(s), "μl" means microliter(s), "ml" means milliliter(s), "L" means liter(s), "μM" means micromolar, "mM" means millimolar, "μg" means microgram(s), "mg" means milligram(s), "A" means adenine or adenosine, "T" means thymine or thymidine, "G" means guanine or guanosine, "C" means cytidine or cytosine, "x g" means times gravity, "nt" means nucleotide(s), "aa" means amino acid(s), "bp" means base pair(s), "kb" means kilobase(s), "k" means kilo, "μ" means micro, and "°C" means degrees Celsius.

EXAMPLE 1

[0230] Applicants' improved EcR-based inducible gene modulation system was developed for use in various applications including gene therapy, expression of proteins of interest in host cells, production of transgenic organisms, and

cell-based assays. This Example describes the construction and evaluation of several gene expression cassettes for use in the EcR-based inducible gene expression system of the invention.

[0231] In various cellular backgrounds, including mammalian cells, insect ecdysone receptor (EcR) heterodimerizes with retinoid X receptor (RXR) and, upon binding of ligand, transactivates genes under the control of ecdysone response elements. Applicants constructed several EcR-based gene expression cassettes based on the spruce budworm *Choristoneura fumiferana* EcR ("CfEcR"; full length polynucleotide and amino acid sequences are set forth in SEQ ID NO: 49 and SEQ ID NO: 50, respectively), *C. fumiferana* ultraspiracle ("CfUSP"; full length polynucleotide and amino acid sequences are set forth in SEQ ID NO: 51 and SEQ ID NO: 52, respectively), and mouse *Mus musculus* RXR α (MmRXR α ; full length polynucleotide and amino acid sequences are set forth in SEQ ID NO: 53 and SEQ ID NO: 54, respectively). The prepared receptor constructs comprise a ligand binding domain of EcR and of RXR or of USP; a DNA binding domain of GAL4 or of EcR; and an activation domain of VP16. The reporter constructs include a reporter gene, luciferase or LacZ, operably linked to a synthetic promoter construct that comprises either GAL4 or EcR/USP binding sites (response elements). Various combinations of these receptor and reporter constructs were cotransfected into CHO, NIH3T3, CV1 and 293 cells. Gene induction potential (magnitude of induction) and ligand specificity and sensitivity were examined using four different ligands: two steroidal ligands (ponasterone A and muristerone A) and two non-steroidal ligands (N-(2-ethyl-3-methoxybenzoyl)-N'-(3,5-dimethylbenzoyl)-N'-tert-butylhydrazine and N-(3,4-(1,2-ethylenedioxy)-2-methylbenzoyl)-N-(3,5-dimethylbenzoyl)-N'-tert-butylhydrazine) in a dose-dependent induction of reporter gene expression in the transfected cells. Reporter gene expression activities were assayed at 24hr or 48hr after ligand addition.

Gene Expression Cassettes: Ecdysone receptor-based, chemically inducible gene expression cassettes (switches) were constructed as followed, using standard cloning methods available in the art. The following is brief description of preparation and composition of each switch.

1.1 - GAL4EcR/VP16RXR: The D, E, and F domains from spruce budworm *Choristoneura fumiferana* EcR ("CfEcRDEF"; SEQ ID NO: 3) were fused to GAL4 DNA binding domain ("DNABD"; SEQ ID NO: 41) and placed under the control of an SV40e promoter (SEQ ID NO: 55). The DEF domains from mouse (*Mus musculus*) RXR ("MmRXRDEF"; SEQ ID NO: 22) were fused to the activation domain from VP16 ("VP16AD"; SEQ ID NO: 45) and placed under the control of an SV40e promoter (SEQ ID NO: 55). Five consensus GAL4 binding sites ("5XGAL4RE"; comprising 5, GAL4RE comprising SEQ ID NO: 47) were fused to a synthetic E1b minimal promoter (SEQ ID NO: 56) and placed upstream of the luciferase gene (SEQ ID NO: 57).

1.2 - GAIAEcR/VP16USP: This construct was prepared in the same way as in switch 1.1 above except MmRXRDEF was replaced with the D, E and F domains from spruce budworm USP ("CfUSPDEF"; SEQ ID NO: 58). The constructs used in this example are similar to those disclosed in U. S. Patent No. 5,880,333 except that *Choristoneura fumiferana* USP rather than *Drosophila melanogaster* USP was utilized.

1.3 - GAL4RXR/VP16CfEcR: MmRXRDEF (SEQ ID NO: 22) was fused to a GAL4DNABD (SEQ ID NO: 41) and CfEcRCDEF (SEQ ID NO: 1) was fused to a VP16AD (SEQ ID NO: 45).

1.4 - GAL4RXR/VP16DmEcR: This construct was prepared in the same way as switch 1.3 except CfEcRCDEF was replaced with DmEcRCDEF (SEQ ID NO: 6).

1.5-GAL4USP/VP16CfEcR: This construct was prepared in the same way as switch 1.3 except MmRXRDEF was replaced with CfUSPDEF (SEQ ID NO: 58).

1.6 - GAIRXRctEcRVPI6: This construct was prepared so that both the GAL4 DNABD and the VP16AD were placed on the same molecule. GAL4DNABD (SEQ ID NO: 41) and VP16AD (SEQ ID NO: 45) were fused to CfEcRDEF (SEQ ID NO: 3) at N-and C-termini respectively. The fusion was placed under the control of an SV40e promoter (SEQ ID NO: 55).

1.7 - VP16CfEcR: This construct was prepared such that CfEcRCDEF (SEQ ID NO: 1) was fused to VP16AD (SEQ ID NO: 45) and placed under the control of an SV40e promoter (SEQ ID NO: 55). Six ecdysone response elements ("EcRE"; SEQ ID NO: 59) from the hsp27 gene were placed upstream of the promoter and a luciferase gene (SEQ ID NO: 57). This switch most probably uses endogenous RXR.

1.8 - DmVgRXR: This system was purchased from Invitrogen Corp., Carlsbad, California. It comprises a *Drosophila melanogaster* EcR ("DmEcR") with a modified DNABD fused to VP16AD and placed under the control of a CMV promoter (SEQ ID NO: 60). Full length MmRXR (SEQ ID NO: 53) was placed under the control of the RSV promoter (SEQ ID NO: 61). The reporter, pIND(SP1)LacZ, contains five copies of a modified ecdysone response element ("EcRE", E/GRE), three copies of an SP1 enhancer, and a minimal heat shock promoter, all of which were placed upstream to the LacZ reporter gene.

1.9 - CfVgRXR: This example was prepared in the same way as switch 1.8 except DmEcR was replaced with a truncated CfEcR comprising a partial A/B domain and the complete CDEF domains [SEQ ID NO: 62 (polynucleotide) and SEQ ID NO: 63 (polypeptide)].

1.10 - CtVgRXRdel: This example was prepared in the same way as switch 1.9 except MmRXR (SEQ ID NO:

53) was deleted.

Cell lines: Four cell lines: CHO, Chinese hamster *Cricetulus griseus* ovarian cell line; NIH3T3 (3T3) mouse *Mus musculus* cell line; 293 human *Homo sapiens* kidney cell line, and CV1 African green monkey kidney cell line were used in these experiments. Cells were maintained in their respective media and were subcultured when they reached 60% confluency. Standard methods for culture and maintenance of the cells were followed.

Transfections: Several commercially available lipofactors as well as electroporation methods were evaluated and the best conditions for transfection of each cell line were developed. CHO, NIH3T3, 293 and CV1 cells were grown to 60% confluency. DNAs corresponding to the various switch constructs outlined in Examples 1.1 through 1.10 were transfected into CHO cells, NIH3T3 cells, 293 cells, or CV1 cells as follows.

CHO cells: Cells were harvested when they reach 60-80% confluency and plated in 6- or 12- or 24- well plates at 250,000, 100,000, or 50,000 cells in 2.5, 1.0, or 0.5 ml of growth medium containing 10% Fetal bovine serum respectively. The next day, the cells were rinsed with growth medium and transfected for four hours. LipofectAMINE™ 2000 (Life Technologies Inc.) was found to be the best transfection reagent for these cells. For 12- well plates, 4 µl of LipofectAMINE™ 2000 was mixed with 100 µl of growth medium 1.0 µg of reporter construct and 0.25 µg of receptor construct(s) were added to the transfection mix. A second reporter construct was added [pTKRL (Promega), 0.1 µg/transfection mix] and comprised a *Renilla* luciferase gene (SEQ ID NO: 64) operably linked and placed under the control of a thymidine kinase (TK) constitutive promoter and was used for normalization. The contents of the transfection mix were mixed in a vortex mixer and let stand at room temperature for 30 min. At the end of incubation, the transfection mix was added to the cells maintained in 400 µl growth medium. The cells were maintained at 37 °C and 5% CO₂ for four hours. At the end of incubation, 500 µl of growth medium containing 20% FBS and either DMSO (control) or a DMSO solution of appropriate ligands were added and the cells were maintained at 37 °C and 5% CO₂ for 24-48 hr. The cells were harvested and reporter activity was assayed. The same procedure was followed for 6 and 24 well plates as well except all the reagents were doubled for 6 well plates and reduced to half for 24-well plates. _

NIH3T3 Cells: Superfect™ (Qiagen Inc.) was found to be the best transfection reagent for 3T3 cells. The same procedures described for CHO cells were followed for 3T3 cells as well with two modifications. The cells were plated when they reached 50% confluency. 125,000 or 50,000 or 25,000 cells were plated per well of 6- or 12- or 24-well plates respectively. The GA14EcR/VP16RXR and reporter vector DNAs were transfected into NIH3T3 cells, the transfected cells were grown in medium containing PonA, MurA. N-(2-ethyl-3-methoxybenzoyl)-N'-(3,5-dimethylbenzoyl)-N'-t-butylhydrazine, or N-(3,4-(1,2-ethylenedioxy)-2-methylbenzoyl)-N-(3,5-dimethylbenzoyl)-N-tert-butylhydrazine for 48 hr. The ligand treatments were performed as described in the CHO cell section above.

293 Cells: LipofectAMINE™ 2000 (Life Technologies) was found to be the best lipofactor for 293 cells. The same procedures described for CHO were followed for 293 cells except that the cells were plated in biocoated plates to avoid clumping. The ligand treatments were performed as described in the CHO cell section above.

CV1 Cells: LipofectAMINE™ plus (Life Technologies) was found to be the best lipofactor for CV1 cells. The same procedures described for NIH3T3 cells were followed for CV1 cells

Ligands: Ponasterone A and Muristerone A were purchased from Sigma Chemical Company. The two non-steroids N-(2-ethyl-3-methoxybenzoyl)-N'-(3,5-dimethylbenzoyl)-N'-t-butylhydrazine, or N-(3,4-(1,2-ethylenedioxy)-2-methylbenzoyl)-N'-(3,5-dimethylbenzoyl)-N'-tert-butylhydrazine are synthetic stable ecdysteroids synthesized at Rohm and Haas Company. All ligands were dissolved in DMSO and the final concentration of DMSO was maintained at 0.1 % in both controls and treatments.

Reporter Assays: Cells were harvested 24-48 hr after adding ligands. 125, 250, or 500 µl of passive lysis buffer (part of Dual-luciferase™ reporter assay system from Promega Corporation) were added to each well of 24- or 12- or 24-well plates respectively. The plates were placed on a rotary shaker for 15 min. Twenty µl of lysate was assayed. Luciferase activity was measured using Dual-luciferase™ reporter assay system from Promega Corporation following the manufacturer's instructions. β-Galactosidase was measured using Galacto-Star™ assay kit from TROPIX following the manufacturer's instructions. All luciferase and β-galactosidase activities were normalized using *Renilla* luciferase as a standard. Fold activities were calculated by dividing normalized relative light units ("RLU") in ligand treated cells with normalized RLU in DMSO treated cells (untreated control).

[0232] The results of these experiments are provided in the following tables.

Table 1

Transactivation of reporter genes through various switches in CHO cells		
Composition of Switch	Mean Fold Activation with 50 μ M N-(2-ethyl-3-methoxybenzoyl)-N'-(3,5-dimethylbenzoyl)-N'-t-butylhydrazine	
1.1 GAL4EcR + VP16RXR pGAL4RELuc	267	
1.2 GAL4EcR + VP16USP pGAL4RELuc	2	
1.3 GAL4RXR + VP16CtEcR pGAL4RELuc	85	
1.4 GAL4RXR+VP16DmEcR pGAL4RELuc	312	
1.5 GAL4USP + VP16CfEcR pGAL4RELuc	2	
1.6 GAL4CfEcRVP16 pGAL4RELuc	9	
1.7 VP16CfEcR pEcRELuc	36	
1.8 DmVgRXR+MmRXR pIND(SP1)LacZ	14	
1.9 CfVgRXR + MmRXR pIND(SP1)LacZ	27	
1.10 CfVgRXR pIND(SP1)LacZ	29	

Table 2

Transactivation of reporter genes through various switches in 3T3 cells		
Composition of Switch	Mean Fold Activation Through N-(2-ethyl-3-methoxybenzoyl)-N'-(3,5-dimethylbenzoyl)-N'-t-butylhydrazine	
1.1 GAL4EcR + VP16RXR pGAL4RELuc	1118	
1.2 GAL4EcR + VP16USP pGAL4RELuc	2	
1.3 GAL4RXR + VP16CfEcR pGAL4RELuc	47	
1.4 GAL4RXR + VP16DmEcR pGAL4RELuc	269	
1.5 GAL4USP + VP16CfEcR pGAL4RELuc	3	
1.6 GAL4CfEcRVP16 pGAL4RELuc	7	
1.7 VP16CfEcR pEcRELuc	1	
1.8 DmVgRXR+MmRXR pIND(SP1)LacZ	21	
1.9 CfVgRXR + MmR7Gt pIND(SP1)LacZ	19	
1.10 CfVgRXR	2	

(continued)

Transactivation of reporter genes through various switches in 3T3 cells	
Composition of Switch	Mean Fold Activation Through N-(2-ethyl-3-methoxybenzoyl)-N'-(3,5-dimethylbenzoyl)-N'-t-butylhydrazine
pIND(SP1)LacZ	

Table 3

Transactivation of reporter genes through various switches in 293 cells	
Composition of Switch	Mean Fold Activation Through N-(2-ethyl-3-methoxybenzoyl)-N'-(3,5-dimethylbenzoyl)-N'-t-butylhydrazine
1.1 GAL4EcR+VP16RXR pGAL4RELuc	125
1.2 GAL4EcR + VP16USP pGAL4RELuc	2
1.3 GAL4RXR+VP16CfEcR pGAL4RELuc	17
1.4 GAL4RXR + VP16DmEcR pGAL4RELuc	3
1.5 GAL4USP + VP16CfEcR pGAL4RELuc	2
1.6 GAL4CfEcRVP16 pGAL4RELuc	3
1.7 VP16CfEcR pEcRELuc	2
1.8 DmVgRXR+ MmRXR pIND(SP1)LacZ	21
1.9 CfVgRXR + MuaRXR pIND(SP1)LacZ	12
1.10 CfVgRXR pIND(SP1)LacZ	3

Table 4

Transactivation of reporter genes through various switches in CV1 cells	
Composition of Switch	Mean Fold Activation Through N-(2-ethyl-3-methoxybenzoyl)-N'-(3,5-dimethylbenzoyl)-N'-t-butylhydrazine
1.1 GAL4EcR + VP16RXR pGAL4RELuc	279
1.2 GAL4EcR+VP16USP pGAL4RELuc	2
1.3 GAL4RXR + VP16CfEcR pGAL4RELuc	25
1.4 GAL4RXR + VP16DmEcR pGAL4RELuc	80
1.5 GAL4USP + VP16CfEcR pGAL4RELuc	3
1.6 GAL4CfEcRVP16 pGAL4RELuc	6

(continued)

Transactivation of reporter genes through various switches in CV1 cells	
Composition of Switch	Mean Fold Activation Through N-(2-ethyl-3-methoxybenzoyl)-N'-(3,5-dimethylbenzoyl)-N'-t-butylhydrazine
1.7 VP16CfEcR pEcRELuc	1
1.8 DmVgRXR+MmRXR pIND(SP1)LacZ	12
1.9 CfVgRXR + MmRXR pIND(SP1)LacZ	7
1.10 Cf VgRXR pIND(SP1)LacZ	1

Table 5

Transactivation of reporter gene GAL4CfEcRDEF/VP16MmRXRDEF (switch 1.1) through steroids and non-steroids in 3T3 cells.	
Ligand	Mean Fold Induction at 1.0 μ M Concentration
1. Ponasterone A	1.0
2. Muristerone A	1.0
3. N-(2-ethyl-3-methoxybenzoyl)-N'-(3,5-dimethylbenzoyl)-N'-tert-butylhydrazine	116
4. N'-(3,4-(1,2-ethylenedioxy)-2-methylbenzoyl)-N'-(3,5-dimethylbenzoyl)-N'-tert-butylhydrazine	601

Table 6

Transactivation of reporter gene GAL4MniRXRDEF/VP16CfEcRDEF (switch 1.3) through steroids and non-steroids in 3T3 cells.	
Ligand	Mean Fold Induction at 1.0 μ M Concentration
1. Ponasterone A	1.0
2. Muristerone A	1.0
3. N-(2-ethyl-3-methoxybenzoyl)-N'-(3,5-dimethylbenzoyl)-N'-tert-butylhydrazine	71
4. N'-(3,4-(1,2-ethylenedioxy)-2-methylbenzoyl)-N'-(3,5-dimethylbenzoyl)-N'-tert-butylhydrazine	54

[0233] Applicants' results demonstrate that the non-steroidal ecdysone agonists, N-(2-ethyl-3-methoxybenzoyl)-N'-(3,5-dimethylbenzoyl)-N'-tert-butylhydrazine and N'-(3,4-(1,2-ethylenedioxy)-2-methylbenzoyl)-N'-(3,5-dimethylbenzoyl)-N'-tert-butylhydrazine, were more potent activators of CfEcR as compared to *Drosophila melanogaster* EcR (DmEcR). (see Tables 1-4). Also, in the mammalian cell lines tested, MmRXR performed better than CfUSP as a heterodimeric partner for CfEcR. (see Tables 1-4). Additionally, Applicants' inducible gene expression modulation system performed better when exogenous MmRXR was used than when the system relied only on endogenous RXR levels (see Tables 1-4).

[0234] Applicants' results also show that in a CfEcR-based inducible gene expression system, the non-steroidal ecdysone agonists induced reporter gene expression at a lower concentration (i.e., increased ligand sensitivity) as compared to the steroid ligands, ponasterone A and muristerone A (see Tables 5 and 6).

[0235] Out of 10 EcR based gene switches tested, the GAL4EcR/VP16RXR switch (Switch 1.1) performed better than any other switch in all four cell lines examined and was more sensitive to non-steroids than steroids. The results also demonstrate that placing the activation domain (AD) and DNA binding domain (DNABD) on each of the two partners reduced background when compared to placing both AD and DNABD together on one of the two partners. Therefore,

a switch format where the AD and DNABD are separated between two partners, works well for EcR-based gene switch applications.

[0236] In addition, the MmRXR/EcR-based switches performed better than CfUSP/EcR-based switches, which have a higher background activity than the MmRXR/EcR switches in the absence of ligand.

[0237] Finally, the GAL4EcR/VP16RXR switch (Switch 1.1) was more sensitive to non-steroid ligands than to the steroid ligands (see Tables 5 and 6). In particular, steroid ligands initiated transactivation at concentrations of 50 μ M, whereas the non-steroid ligands initiated transactivation at less than 1 μ M (submicromolar) concentration.

EXAMPLE 2

[0238] This Example describes Applicants' further analysis of truncated EcR and RXR polypeptides in the improved EcR-based inducible gene expression system of the invention. To identify the best combination and length of two receptors that give a switch with a) maximum induction in the presence of ligand; b) minimum background in the absence of ligand; c) highly sensitive to ligand concentration; and d) minimum cross-talk among ligands and receptors, Applicants made and analyzed several truncation mutations of the CfEcR and MmRXR receptor polypeptides in NIH3T3 cells.

[0239] Briefly, polynucleotides encoding EcR or RXR receptors were truncated at the junctions of A/B, C, D, E and F domains and fused to either a GAL4 DNA binding domain encoding polynucleotide (SEQ ID NO: 41) for CfEcR, or a VP16 activation domain encoding polynucleotide (SEQ ID NO: 45) for MmRXR as described in Example 1. The resulting receptor truncation/fusion polypeptides were assayed in NIH3T3 cells. Plasmid pFRLUC (Stratagene) encoding a luciferase polypeptide was used as a reporter gene construct and pTKRL (Promega) encoding a *Renilla* luciferase polypeptide under the control of the constitutive TK promoter was used to normalize the transfections as described above. The analysis was performed in triplicates and mean luciferase counts were determined as described above.

Gene Expression Cassettes Encoding Truncated Ecdysone Receptor Polypeptides

[0240] Gene expression cassettes comprising polynucleotides encoding either full length or truncated CfEcR polypeptides fused to a GAL4 DNA binding domain (SEQ ID NO: 41): GAL4CfEcRABCDEF (full length CfEcR/ABCDEF; SEQ ID NO: 49), GAL4CfEcRCDEF (CfEcRCDEF; SEQ ID NO: 1), GAL4CfEcR1/2CDEF (CfEcR1/2CDEF; SEQ ID NO: 2), GAL4CfEcRDEF (CfEcRDEF; SEQ ID NO: 3), GAL4CfEcREF (CfEcREF; SEQ ID NO: 4), and GAL4CfEcRDE (CfEcRDE; SEQ ID NO: 5) were transfected into NIH3T3 cells along with VP16MmRXRDEF (constructed as in Example 1.1; Figure 11) or VP16MmRXREF [constructed as in Example 1.1 except that MmRXRDEF was replaced with MmRXREF (SEQ ID NO: 23); Figure 12], and pFRLUC and pTKRL plasmid DNAs. The transfected cells were grown in the presence 0, 1, 5 or 25 μ M of N-(2-ethyl-3-methoxybenzoyl)-N'-(3,5-dimethylbenzoyl)-N'-tert-butylhydrazine or PonA for 48 hr. The cells were harvested, lysed and luciferase reporter activity was measured in the cell lysates. Total fly luciferase relative light units are presented. The number on the top of each bar is the maximum fold induction for that treatment.

[0241] Applicants' results show that the EF domain of MmRXR is sufficient and performs better than DEF domains of this receptor (see Figures 11 and 12). Applicants have also shown that, in general, EcR/RXR receptor combinations are insensitive to PonA (see Figures 11 and 12). As shown in the Figures 11 and 12, the GAL4CfEcRCDEF hybrid polypeptide (SEQ ID NO: 7) performed better than any other CfEcR hybrid polypeptide.

Gene Expression Cassettes Encoding Truncated Retinoid X Receptor Polypeptides

[0242] Gene expression cassettes comprising polynucleotides encoding either full length or truncated MmRXR polypeptides fused to a VP16 transactivation domain (SEQ ID NO: 45): VP16MmRXRA/BCDEF (full length MmRXRA/BCDEF; SEQ ID NO: 53), VP16MmRXRCDEF (MmRXRCDEF; SEQ ID NO: 21), VP16MmRXRDEF (MmRXR3CRDEF; SEQ ID NO: 22), VP16MmRXREF (MmRXREF; SEQ ID NO: 23), VP16MmRXRBam-EF ("MmRXRBam-EF" or "MmRXR-truncatedEF"; SEQ ID NO: 24), and VP16MmRXRAF2de1 ("MmRXRAF2de1" or "MmRXR-E"; SEQ ID NO: 25) constructs were transfected into NIH3T3 cells along with GAL4CfEcRCDEF (constructed as in Example 1.1; Figure 13) or GAL4CfEcRDEF [constructed as in Example 1.1 except CfEcRCDEF was replaced with CfEcRDEF (SEQ ID NO: 3); Figure 14], pFRLUC and pTKRL plasmid DNAs as described above. The transfected cells were grown in the presence 0, 1, 5 and 25 μ M of N-(2-ethyl-3-methoxybenzoyl)-N'-(3,5-dimethylbenzoyl)-N'-tert-butylhydrazine or PonA for 48 hr. The cells were harvested and lysed and reporter activity was measured in the cell lysate. Total fly luciferase relative light units are presented. The number on top of each bar is the maximum fold induction in that treatment.

[0243] Of all the truncations of MmRXR tested, Applicants' results show that the MmRXREF receptor was the best partner for CfEcR (Figures 13 and 14). CfEcRCDEF showed better induction than CfEcRDEF using MmRXREF. Deleting AF2 (abbreviated "EF-AF2del") or helices 1-3 of the E domain (abbreviated "EF-Bamdel") resulted in an RXR receptor that reduced gene induction and ligand sensitivity when partnered with either CfEcRCDEF (Figure 13) or CfEcRDEF (Figure 14) in NIH3T3 cells. In general, the CfEcR/RXR-based switch was much more sensitive to the non-steroid N-(2-

ethyl-3-methoxybenzoyl)-N'-(3,5-dimethylbenzoyl)-N'-tert-butylhydrazine than to the steroid PonA.

EXAMPLE 3

[0244] This Example describes Applicants' further analysis of gene expression cassettes encoding truncated EcR or RXR receptor polypeptides that affect either ligand binding activity or ligand sensitivity, or both. Briefly, six different combinations of chimeric receptor pairs, constructed as described in Examples 1 and 2, were further analyzed in a single experiment in NIH3T3 cells. These six receptor pair combinations and their corresponding sample numbers are depicted in Table 7.

Table 7

CfEcR + MmRXR Truncation Receptor Combinations in NIH3T3 Cells		
Figure 15 X-Axis Sample No.	EcR Polypeptide Construct	RXR Polypeptide Construct
Samples 1 and 2	GAL4CfEcRCDEF	VP16RXRABCDEF (Full length)
Samples 3 and 4	GAL4CfEcRCDEF	VP16RXRDEF
Samples 5 and 6	GAL4CfEcRCDEP	VP16RXREF
Samples 7 and 8	GAL4CfEcRDEF	VP16RXRA/BCDEF (Full length)
Samples 9 and 10	GAL4CfEcRDEF	VP16RXRDEF
Samples 11 and 12	GAL4CfEcRDEF	VP16RXREF

[0245] The above receptor construct pairs, along with the reporter plasmid pFRLuc were transfected into NIH3T3 cells as described above. The six CfEcR truncation receptor combinations were duplicated into two groups and treated with either steroid (odd numbers on x-axis of Figure 15) or non-steroid (even numbers on x-axis of Figure 15). In particular, the cells were grown in media containing 0, 1, 5 or 25 uM PonA (steroid) or N-(2-ethyl-3-methoxybenzoyl)-N'-(3,5-dimethylbenzoyl)-N'-tert-butylhydrazine (non-steroid) ligand. The reporter gene activity was measured and total RLU are shown. The number on top of each bar is the maximum fold induction for that treatment and is the mean of three replicates.

[0246] As shown in Figure 15, the CfEcRCDEF/MmRXREF receptor combinations were the best switch pairs both in terms of total RLU and fold induction (compare columns 1-6 to columns 7-12). This confirms Applicants' earlier findings as described in Example 2 (Figures 11-14). The same gene expression cassettes encoding the truncated EcR and RXR polypeptides were also assayed in a human lung carcinoma cell line A549 (ATCC) and similar results were observed (data not shown).

SEQUENCE LISTING

[0247]

<110> Rohm and Haas Company
Palli, Subba Reddy
Kapitskaya, Marianna Zinovjevna
Cress, Dean Ervin
<120> Novel Ecdysone Receptor-Based Inducible Gene Expression System
<130> RH0020
<150> 60/191,355
<151> 2000-03-22
<150> 60/269,799
<151> 2001-02-20
<160> 64
<170> PatentIn version 3.0 ,
<210> 1
<211> 1288
<212> DNA
<213> Artificial Sequence
<220>
<221> misc feature

EP 1 266 015 B1

<223> Novel Sequence

<400> 1

5	aagggccctg cgccccgtca gcaagaggaa ctgtgtctgg tatgcgggga cagagcctcc	60
	ggataccact acaatgcgct cacgtgtgaa ggggtgtaaag gggtcttcag acggagtgtt	120
	accaaaaaatg cggttttatat ttgtaaattc ggtcacgctt gcgaaatgga catgtacatg	180
10	cgacggaaat gccaggagtg ccgcctgaag aagtgccttag ctgtaggcat gaggcctgag	240
	tgcgtagtac ccgagactca gtgcgccatg aagcggaaaag agaagaaagc acagaaggag	300
	aaggacaaac tgcctgtcag cagcagcagc gtggacgacc acatgccgcc cattatgcag	360
	tgtgaacctc cacctcctga agcagcaagg attcacgaag tggccccaaag gtttctctcc	420
15	gacaagctgt tggagacaaa ccggcagaaa aacatcccc agttgacagc caaccagcag	480
	ttccttatcg ccaggctcat ctggtaccag gacgggtacg agcagccttc tgatgaagat	540
	ttgaagagga ttacgcagac gtggcagcaa gcggacgatg aaaacgaaga gtctgacact	600
20	cccttcgcc agatcacaga gatgactatc ctcacggctc aacttatcgt ggagttcgcg	660
	aagggattgc cagggttcgc caagatctcg cagcctgatc aaattacgct gcttaaggct	720
	tgctcaagtg aggtaatgat gctccgagtc gcgcgacgat acgatgcggc ctcagacagt	780
25	gttctgttcg cgaacaacca agcgtacact cgcgacaact accgcaaggc tggcatggcc	840
	tacgtcatcg aggatctact gcacttctgc cgggtgatgt actctatggc gttggacaac	900
	atccattacg cgctgtctac ggctgtcgtc atcttttctg accggccagg gttggagcag	960
30	ccgcaactgg tggaagaaat ccagcggtag tacctgaata cgctccgcat ctatatcctg	1020
	aaccagctga gcgggtcggc gcgttcgtcc gtcatatacg gcaagatcct ctcaatcctc	1080
35	tctgagctac gcacgctcgg catgcaaaac tccaacatgt gcattctcct caagctcaag	1140
	aacagaaaagc tgccgccttt cctcgaggag atctgggatg tggcggacat gtcgcacacc	1200
	caaccgccgc ctatcctcga gtccccacg aatctctagc ccctgcgcgc acgcatcgcc	1260
40	gatgccgcgt ccggccgcgc tgctctga	1288

<210> 2

<211> 1110

45 <212> DNA

<213> Artificial Sequence

<220>

<221> misc_feature

<223> Novel Sequence

50 <400> 2

55

EP 1 266 015 B1

	gcggtttata tttgtaaatt cggtcacgct tgcgaaatgg acatgtacat gcgacggaaa	60
	tgccaggagt gccgcctgaa gaagtgctta gctgtaggca tgaggcctga gtgcgtagta	120
5	cccgagactc agtgcgccat gaagcggaaa gagaagaaaag cacagaagga gaaggacaaa	180
	ctgcctgtca gcacgacgac ggtggacgac cacatgccgc ccattatgca gtgtgaacct	240
	ccacctcctg aagcagcaag gattcacgaa gtggtcccaa ggtttctctc cgacaagctg	300
10	ttggagacaa accggcagaa aaacatcccc cagttgacag ccaaccagca gttccttatt	360
	gccaggctca tctggtacca ggacgggtac gagcagcctt ctgatgaaga tttgaagagg	420
	attacgcaga cgtggcagca agcggacgat gaaaacgaag agtctgacac tcccttcgcg	480
15	cagatcacag agatgactat cctcacggtc caacttatcg tggagttcgc gaagggattg	540
	ccagggttcg ccaagatctc gcagcctgat caaattacgc tgcttaaggc ttgctcaagt	600
	gaggtaatga tgctccgagt cgcgcgacga tacgatgcgg cctcagacag tgttctgttc	660
20	gcgaacaacc aagcgtacac tcgcgacaac taccgcaagg ctggcatggc ctacgtcatc	720
	gaggatctac tgcacttctg ccggtgcatg tactctatgg cgttggaaca catccattac	780
	gcgctgctca cggctgtcgt catcttttct gaccggccag ggttggaaga gccgcaactg	840
25	gtggaagaaa tccagcggta ctacctgaat acgctccgca tctatatcct gaaccagctg	900
	agcgggtcgg cgcgttcgtc cgtcatatac ggcaagatcc tctcaatcct ctctgagcta	960
	cgcacgctcg gcatgcaaaa ctccaacatg tgcattctcc tcaagctcaa gaacagaaaag	1020
30	ctgccgcctt tcctcgagga gatctgggat gtggcggaca tgtcgcacac ccaaccgccg	1080
	cctatcctcg agtccccac gaatctctag	1110

<210> 3

<211> 1054

<212> DNA

<213> Artificial Sequence

<220>

<221> misc_feature

<223> Novel Sequence

<400> 3

EP 1 266 015 B1

cctgagtgcg tagtaccgga gactcagtgc gccatgaagc ggaaagagaa gaaagcacag 60
 aaggagaagg acaaactgcc tgtcagcacg acgacgggtgg acgaccacat gccgccatt 120
 5 atgcagtgtg aacctccacc tcctgaagca gcaaggattc acgaagtggc cccaaggttt 180
 ctctccgaca agctgttgga gacaaaccgg cagaaaaaca tccccagtt gacagccaac 240
 cagcagttcc ttatcgccag gctcatctgg taccaggacg ggtacgagca gccttctgat 300
 10 gaagatttga agaggattac gcagacgtgg cagcaagcgg acgatgaaaa cgaagagtct 360
 gacactccct tccgccagat cacagagatg actatcctca cgggccaaact tatcgtggag 420
 ttcggaagg gattgccagg gtctgccaag atctcgagc ctgatcaaata tacgtgctt 480
 15 aaggcttgc caagtgaggt aatgatgctc cgagtcgagc gacgatacga tgcggcctca 540
 gacagtgttc tgttcgcgaa caaccaagcg tacactcgcg acaactaccg caaggctggc 600
 atggcctacg tcatcgagga tctactgcac ttctgcccgt gcatgtactc tatggcgttg 660
 20 gacaacatcc attacgcgt gctcacggct gtcgtcatct tttctgaccg gccagggttg 720
 gagcagccgc aactgggtga agaaatccag cgggtactacc tgaatacgtc ccgcatctat 780
 atcctgaacc agctgagcgg gtcggcgagt tcgtccgtca tatacggcaa gatcctctca 840
 25 atcctctctg agctacgcac gtcggcatg caaaactcca acatgtgcat ctccctcaag 900
 ctcaagaaca gaaagctgcc gcctttcctc gaggagatct gggatgtggc ggacatgtcg 960
 cacaccaaac cgccgcctat cctcgagtcc cccacgaatc tctagcccct gcgcgcacgc 1020
 atcgccgatg ccgcgtccgg ccgcgtgct ctga 1054

30

<210> 4

<211> 735

<212> DNA

35

<213> Artificial Sequence

<220>

<221> misc feature

<223> Novel Sequence

<400> 4

40

taccaggacg ggtacgagca gccttctgat gaagatttga agaggattac gcagacgtgg 60
 cagcaagcgg acgatgaaaa cgaagagtct gacactccct tccgccagat cacagagatg 120
 45 actatcctca cgggccaaact tatcgtggag ttcggaagg gattgccagg gtctgccaag 180
 atctcgagc ctgatcaaata tacgtgctt aaggcttgc caagtgaggt aatgatgctc 240
 cgagtcgagc gacgatacga tgcggcctca gacagtgttc tgttcgcgaa caaccaagcg 300

50

55

EP 1 266 015 B1

5 tacactcgcg acaactaccg caaggctggc atggcctacg tcatcgagga tctactgcac 360
 ttctgccggt gcatgtactc tatggcggtg gacaacatcc attacgcgct gctcacggct 420
 gtcgtcatct tttctgaccg gccagggttg gagcagccgc aactggtgga agaaatccag 480
 cgggtactacc tgaatacgct ccgcatttat atcctgaacc agctgagcgg gtcggcgcggt 540
 tcgtccgtca tatacggcaa gatcctctca atcctctctg agctacgcac gctcggcatg 600
 10 caaaactcca acatgtgcat ctccctcaag ctcaagaaca gaaagctgcc gcctttcctc 660
 gaggagatct gggatgtggc ggacatgtcg cacacccaac cgcgcctat cctcgagtcc 720
 cccacgaatc tctag 735

15 <210> 5
 <211> 960
 <212> DNA
 <213> Artificial Sequence
 20 <220>
 <221> misc_feature
 <223> Novel Sequence
 <400> 5

25 cctgagtgcg tagtaccgga gactcagtgc gccatgaagc ggaaagagaa gaaagcacag 60
 aaggagaagg acaaactgcc tgtcagcacg acgacgggtg acgaccacat gccgccatt 120
 atgcagtgtg aacctccacc tcctgaagca gcaaggattc acgaagtggg cccaaggttt 180
 30 ctctccgaca agctgttgga gaaaaccgg cagaaaaaca tccccagtt gacagccaac 240
 cagcagttcc ttatcgccag gctcatctgg taccaggacg ggtacgagca gccttctgat 300
 gaagatttga agaggattac gcagacgtgg cagcaagcgg acgatgaaaa cgaagagtct 360
 35 gacactccct tccgccagat cacagagatg actatcctca cgggtccaact tatcgtggag 420
 ttcgcgaagg gattgccagg gttcgccaag atctcgagc ctgatcaaat tacgctgctt 480
 aaggcttgct caagttaggt aatgatgctc cgagtcgcgc gacgatacga tgcggcctca 540
 40 gacagtgttc tggtcgcgaa caaccaagcg tacactcgcg acaactaccg caaggctggc 600
 atggcctacg tcatcgagga tctactgcac ttctgccggt gcatgtactc tatggcggtg 660
 gacaacatcc attacgcgct gctcacggct gtcgtcatct tttctgaccg gccagggttg 720
 45 gagcagccgc aactggtgga agaaatccag cgggtactacc tgaatacgct ccgcatttat 780
 atcctgaacc agctgagcgg gtcggcgcggt tcgtccgtca tatacggcaa gatcctctca 840
 atcctctctg agctacgcac gctcggcatg caaaactcca acatgtgcat ctccctcaag 900
 50 ctcaagaaca gaaagctgcc gcctttcctc gaggagatct gggatgtggc ggacatgtcg 960

55 <210> 6
 <211> 1878
 <212> DNA
 <213> Artificial Sequence
 <220>
 <221> misc feature
 <223> Novel Sequence

EP 1 266 015 B1

<400> 6

	ggacctgctgc caccgggtgca agaggagctg tgcctgggttt gcggcgacag ggcctccggc	60
5	taccactaca acgccctcac ctgtgagggc tgcaaggggt tctttcgacg cagcgttacg	120
	aagagcgccg tctactgctg caagttcggg cgcgcctgcg aaatggacat gtacatgagg	180
	cgaaagtgtc aggagtgcg cctgaaaaag tgcctggccg tgggtatgcg gccggaatgc	240
10	gtcgtcccgg agaaccaatg tgcgatgaag cggcgcgaaa agaaggcca gaaggagaag	300
	gacaaaatga ccacttcgcc gagctctcag catggcggca atggcagctt ggcctctggt	360
	ggcggccaag actttgttaa gaaggagatt cttgacctta tgacatgcga gccgccccag	420
15	catgccacta ttccgctact acctgatgaa atattggcca agtgtcaagc gcgcaatata	480
	ccttccttaa cgtacaatca gttggccgtt atatacaagt taatttggtta ccaggatggc	540
	tatgagcagc catctgaaga ggatctcagg cgtataatga gtcaaccgca tgagaacgag	600
20	agccaaacgg acgtcagctt tcggcatata accgagataa ccatactcac ggtccagttg	660
	attgttgagt ttgctaaagg tctaccagcg ttacaaaaga taccacagga ggaccagatc	720
	acgttactaa aggcctgctc gtcggaggtg atgatgctgc gtatggcacg acgtatgac	780
25	cacagctcgg actcaatatt cttcggaat aatagatcat atacgcggga ttcttacaaa	840
	atggccggaa tggttgataa cattgaagac ctgctgcatt tctgccgcca aatgttctcg	900
	atgaagtggtg acaacgtcga atacgcgctt ctactgccca ttgtgatctt ctcgaccgg	960
30	ccgggcctgg agaaggcca actagtcgaa gcatccaga gtaactacat cgacacgcta	1020
	cgcatttata tactcaaccg ccactgcggc gactcaatga gcctcgtctt ctacgcaaag	1080
	ctgctctcga tcctcaccga gctgcgtacg ctgggcaacc agaacgccga gatgtgtttc	1140
	tcactaaagc tcaaaaaccg caaactgccc aagttcctcg aggagatctg ggacgttcat	1200
35	gccatcccgc catcggtcca gtcgcacctt cagattacce aggaggagaa cgagcgtctc	1260
	gagcgggctg agcgtatgcg ggcatcggtt gggggcgcca ttaccgccgg cattgattgc	1320
	gactctgcct ccacttcggc ggcggcagcc gcggcccage atcagcctca gcctcagccc	1380
40	cagccccaac cctctccct gaccagaac gattcccagc accagacaca gccgcagcta	1440
	caacctcagc taccacctca gctgcaaggt caactgcaac ccagctcca accacagctt	1500
	cagacgcaac tccagccaca gattcaacca cagccacagc tccttccgt ctccgtccc	1560
45	gtgccgcct ccgtaaccgc acctggttcc ttgtccgcgg tcagtaacgag cagcgaatac	1620
	atgggcggaa gtgcggccat aggaccatc acgccggcaa ccaccagcag tatcacggct	1680
	gccgttaccg ctagctccac cacatcagcg gtaccgatgg gcaacggagt tggagtcggt	1740
50		
	gttggggtgg gcggcaacgt cagcatgtat gcgaacgcc agacggcgat ggccttgatg	1800
	ggtgtagccc tgcattcgca ccaagagcag cttatcgggg gaggggcgg taagtcggag	1860
55	cactcgacga ctgcatag	1878

<210> 7

EP 1 266 015 B1

<211> 1752
 <212> DNA
 <213> Artificial Sequence
 <220>
 <221> misc feature
 <223> Novel Sequence
 <400> 7

```

10      gccgtctact gctgcaagtt cgggcgcgcc tgcgaaatgg acatgtacat gaggcgaaag      60
      tgtcaggagt gccgcctgaa aaagtgcctg gccgtgggta tgcggccgga atgcgtcgtc      120
      ccggagaacc aatgtgcgat gaagcggcgc gaaaagaagg ccagaagga gaaggacaaa      180
15      atgaccactt cgccgagctc tcagcatggc ggcaatggca gcttggcctc tggtgccggc      240
      caagactttg ttaagaagga gattcttgac cttatgacat gcgagccgcc ccagcatgcc      300
      actattccgc tactacctga tgaaatattg gccaaagtgc aagcgcgcaa tataccttcc      360
      ttaacgtaca atcagttggc cgttatatac aagttaattt ggtaccagga tggctatgag      420
20      cagccatctg aagaggatct caggcgtata atgagtcaac ccgatgagaa cgagagccaa      480
      acggacgtca gctttcggca tataaccgag ataaccatac tcacggcca gttgattggt      540
      gagtttgcta aaggtctacc agcgtttaca aagatacccc aggaggacca gatcacgtta      600
25      ctaaaggcct gctcgtcgga ggtgatgatg ctgcgtatgg cagcagccta tgaccacagc      660
      tcggactcaa tattcttcgc gaataataga tcatatacgc gggattctta caaaatggcc      720
      ggaatggctg ataacattga agacctgctg catttctgcc gccaaatgtt ctcgatgaag      780
30      gtggacaacg tcgaatacgc gcttctcact gccattgtga tcttctcgga cgggcggggc      840
      ctggagaagg cccaactagt cgaagcgatc cagagctact acatcgacac gctacgcatt      900
      tatatactca accgccactg cggcgactca atgagcctcg tcttctacgc aaagctgctc      960
35      tcgatcctca ccgagctgcg tacgctgggc aaccagaacg ccgagatgtg tttctcacta      1020
      aagctcaaaa accgcaaact gcccaagttc ctcgaggaga tctgggacgt tcatgccatc      1080
      ccgccatcgg tccagtcgca cttcagatt acccaggagg agaacgagcg tctcgagcgg      1140
40      gctgagcgta tgcgggcatc ggttgggggc gccattaccg ccggcattga ttgcgactct      1200
      gcctccactt cggcggcggc agccgcggcc cagcatcagc ctcagcctca gccccagccc      1260
      caaccctcct ccctgaccca gaacgattcc cagcaccaga cacagccgca gctacaacct      1320
45      cagctaccac ctcagctgca aggtcaactg caaccccagc tccaaccaca gtttcagacg      1380

```

EP 1 266 015 B1

	caactccagc cacagattca accacagcca cagctccttc ccgtctccgc tcccgtgccc	1440
	gcctccgtaa ccgcacctgg ttcccttgcc gcggtcagta cgagcagcga atacatgggc	1500
5	ggaagtgcgg ccataggacc catcacgccg gcaaccacca gcagtatcac ggctgccgtt	1560
	accgctagct ccaccacatc agcggtagcg atgggcaacg gagttggagt cgggtgttggg	1620
	gtggggcgga acgtcagcat gtatgcgaac gccacagacg cgatggcctt gatgggtgta	1680
10	gccctgcatt cgcaccaaga gcagcttata gggggagtg cggttaagtc ggagcactcg	1740
	acgactgcat ag	1752

	<210> 8
15	<211> 1650
	<212> DNA
	<213> Artificial Sequence
	<220>
	<221> misc feature
20	<223> Novel Sequence
	<400> 8

	cggccggaat gcgtcgtccc ggagaaccaa tgtgcgatga agcggcgcca aaagaaggcc	60
25	cagaaggaga aggacaaaat gaccacttcg ccgagctctc agcatggcgg caatggcagc	120
	ttggcctctg gtggcgcca agactttgtt aagaaggaga ttcttgacct tatgacatgc	180
	gagccgcccc agcatgccac tattccgcta ctacctgatg aaatattggc caagtgtcaa	240
30	gcgcgcaata taccttccct aacgtacaat cagttggcgg ttatatacaa gtttaatttg	300
	taccaggatg gctatgagca gccatctgaa gaggatctca ggcgtataat gagtcaaccc	360
	gatgagaacg agagccaaac ggacgtcagc tttcggcata taaccgagat aaccatactc	420
35	acgggtccagt tgattgttga gtttgctaaa ggtctaccag cgtttacaaa gataccccag	480
	gaggaccaga tcacgttact aaaggcctgc tcgtcggagg tgatgatgct gcgtatggca	540
	cgacgctatg accacagctc ggactcaata ttcttcgcca ataatagatc atatacgcg	600
40	gattcttaca aaatggccgg aatggctgat aacattgaag acctgctgca tttctgccgc	660
	caaattgtct cgatgaaggc ggacaacgtc gaatacgcgc ttctcactgc cattgtgatc	720
	ttctcggacc ggccgggcct ggagaaggcc caactagtcg aagcgatcca gagctactac	780
45	atcgacacgc tacgcattta tatactcaac cgccactgcg gcgactcaat gagcctcgtc	840
	ttctacgcaa agctgctctc gatcctcacc gagctgcgta cgctgggcaa ccagaacgcc	900
	gagatgtgtt tctcactaaa gctcaaaaac cgcaaaactgc ccaagttcct cgaggagatc	960
50	tgggacgttc atgcatccc gccatcggtc cagtcgcacc ttcagattac ccaggaggag	1020
	aacgagcgtc tcgagcgggc tgagcgtatg cgggcacgcg ttgggggcgc cattaccgcc	1080
	ggcattgatt gcgactctgc ctccacttcg gcggcggcag ccgcggccca gcacagcct	1140
55	cagcctcagc ccagcccca accctcctcc ctgaccaga acgattccca gcaccagaca	1200

EP 1 266 015 B1

5 cagccgcagc tacaacctca gctaccacct cagctgcaag gtcaactgca accccagctc 1260
 caaccacagc ttcagacgca actccagcca cagattcaac cacagccaca gtccttccc 1320
 gtctecgctc ccgtgccgc ctccgtaacc gcacctggtt ccttgccgc ggtcagtagc 1380
 agcagcgaat acatgggcgg aagtgcggcc ataggacca tcacgccgcg aaccaccagc 1440
 agtatcacgg ctgccgttac cgctagctcc accacatcag cggtagcgat gggcaacgga 1500
 10 gttggagtcg gtgttgggtt gggcggaac gtcagcatgt atgcgaacgc ccagacggcg 1560
 atggccttga tgggtgtagc cctgcattcg caccaagagc agcttatcgg gggagtggcg 1620
 gttaagtcgg agcactcgac gactgcatag 1650

15
 <210> 9
 <211> 1338
 <212> DNA
 <213> Artificial Sequence
 20 <220>
 <221> misc feature
 <223> Novel Sequence
 <400> 9

25 tatgagcagc catctgaaga ggatctcagg cgtataatga gtcaaccgga tgagaacgag 60
 agccaaacgg acgtcagctt tcggcatata accgagataa ccatactcac ggtccagttg 120
 attgttgagt ttgctaaagg tctaccagcg tttaaaaga taccacagga ggaccagatc 180
 30 acgttactaa aggcctgctc gtcggaggtg atgatgctgc gtatggcacg acgctatgac 240
 cacagctcgg actcaatatt cttcgcgaat aatagatcat atacgcggga ttcttacaaa 300
 atggccggaa tggctgataa cattgaagac ctgctgcatt tctgccgcca aatgttctcg 360
 35 atgaagggtg acaacgtcga atacgcgctt ctcaactgcca ttgtgatctt ctccgaccgg 420
 ccgggcctgg agaaggccca actagtcgaa gcgatccaga gctactacat cgacacgcta 480
 cgcatttata tactcaaccg ccactgcggc gactcaatga gcctcgtctt ctacgcaaag 540
 40 ctgctctcga tcctcaccga gctgcgtacg ctgggcaacc agaacgccga gatgtgtttc 600
 tcactaaagc tcaaaaaccg caaactgcc aagttcctcg aggagatctg ggacgttcat 660
 gccatcccg cactcggtcca gtcgcacctt cagattaccc aggaggagaa cgagcgtctc 720
 45 gagcgggctg agcgtatgcg ggcacgggtt gggggcgcca ttaccgccgg cattgattgc 780
 gactctgcct ccacttcggc ggcggcagcc ggggccagc atcagcctca gcctcagccc 840
 cagccccaac cctcctcctt gaccagaac gattccagc accagacaca gccgcagcta 900
 50 caacctcagc taccacctca gctgcaagg caactgcaac ccagctcca accacagctt 960
 cagacgcaac tccagccaca gattcaacca cagccacagc tccttcccgt ctccgctccc 1020
 gtgcccgcct ccgtaaccgc acctgggttc ttgtccgcgg tcagtacgag cagcgaatac 1080

55

EP 1 266 015 B1

atgggcggaa gtgcggccat aggacccatc acgccggcaa ccaccagcag tatcacggct 1140
 gccgttaccg ctagctccac cacatcagcg gtaccgatgg gcaacggagt tggagtcggt 1200
 5 gttggggtgg gcggcaacgt cagcatgtat gcgaacgccc agacggcgat ggccttgatg 1260
 ggtgtagccc tgcattcgca ccaagagcag cttatcgggg gagtggcggt taagtcggag 1320
 cactcgacga ctgcatag 1338

10 <210> 10
 <211> 969
 <212> DNA
 <213> Artificial Sequence
 15 <220>
 <221> misc feature
 <223> Novel Sequence
 <400> 10

20 cggccggaat gcgtcgtccc ggagaaccaa tgtgcgatga agcggcgcga aaagaaggcc 60
 cagaaggaga aggacaaaat gaccacttcg ccgagctctc agcatggcgg caatggcagc 120
 ttggcctctg gtggcggcca agactttgtt aagaaggaga ttcttgacct tatgacatgc 180
 25 gagccgcccc agcatgccac tattccgcta ctacctgatg aaatattggc caagtgtcaa 240
 gcgcgcaata taccttcctt aacgtacaat cagttggccg ttatatacaa gttaatttgg 300
 taccaggatg gctatgagca gccatctgaa gaggatctca ggcgtataat gagtcaaccc 360
 30 gatgagaacg agagccaaac ggacgtcagc ttctggcata taaccgagat aaccatactc 420
 acggtccagt tgattgttga gtttgctaaa ggtctaccag cgtttacaaa gataccccag 480
 gaggaccaga tcacgttact aaaggcctgc tcgtcggagg tgatgatgct gcgtatggca 540
 35 cgacgctatg accacagctc ggactcaata ttcttcgcga ataatagatc atatacgcg 600
 gattcttaca aaatggccgg aatggctgat aacattgaag acctgctgca tttctgccgc 660
 caaatgttct cgatgaagggt ggacaacgtc gaatacgcgc ttctcactgc cattgtgatc 720
 40 ttctcggacc ggccgggccc ggagaaggcc caactagtcg aagcgatcca gagctactac 780
 atcgacacgc tacgcattta tatactcaac cgccactgcg gcgactcaat gagcctcgtc 840
 ttctacgcaa agctgctctc gatcctcacc gagctgcgta cgctgggcaa ccagaacgcc 900
 gagatgtgtt tctcactaaa gctcaaaaac cgcaaactgc ccaagttcct cgaggagatc 960
 45 tgggacggtt 969

<210> 11
 <211> 412
 50 <212> PRT
 <213> Artificial Sequence
 <220>
 <221> misc feature
 <223> Novel Sequence
 55 <400> 11

EP 1 266 015 B1

	Lys	Gly	Pro	Ala	Pro	Arg	Gln	Gln	Glu	Glu	Leu	Cys	Leu	Val	Cys	Gly	
	1				5				10					15			
5	Asp	Arg	Ala	Ser	Gly	Tyr	His	Tyr	Asn	Ala	Leu	Thr	Cys	Glu	Gly	Cys	
				20					25					30			
	Lys	Gly	Phe	Phe	Arg	Arg	Ser	Val	Thr	Lys	Asn	Ala	Val	Tyr	Ile	Cys	
			35					40					45				
10	Lys	Phe	Gly	His	Ala	Cys	Glu	Met	Asp	Met	Tyr	Met	Arg	Arg	Lys	Cys	
		50					55					60					
	Gln	Glu	Cys	Arg	Leu	Lys	Lys	Cys	Leu	Ala	Val	Gly	Met	Arg	Pro	Glu	
		65				70					75					80	
15	Cys	Val	Val	Pro	Glu	Thr	Gln	Cys	Ala	Met	Lys	Arg	Lys	Glu	Lys	Lys	
					85					90					95		
	Ala	Gln	Lys	Glu	Lys	Asp	Lys	Leu	Pro	Val	Ser	Thr	Thr	Thr	Val	Asp	
				100					105					110			
20	Asp	His	Met	Pro	Pro	Ile	Met	Gln	Cys	Glu	Pro	Pro	Pro	Pro	Glu	Ala	
			115					120					125				
	Ala	Arg	Ile	His	Glu	Val	Val	Pro	Arg	Phe	Leu	Ser	Asp	Lys	Leu	Leu	
		130					135					140					
25	Glu	Thr	Asn	Arg	Gln	Lys	Asn	Ile	Pro	Gln	Leu	Thr	Ala	Asn	Gln	Gln	
		145				150					155					160	
	Phe	Leu	Ile	Ala	Arg	Leu	Ile	Trp	Tyr	Gln	Asp	Gly	Tyr	Glu	Gln	Pro	
					165				170						175		
30	Ser	Asp	Glu	Asp	Leu	Lys	Arg	Ile	Thr	Gln	Thr	Trp	Gln	Gln	Ala	Asp	
				180					185					190			
	Asp	Glu	Asn	Glu	Glu	Ser	Asp	Thr	Pro	Phe	Arg	Gln	Ile	Thr	Glu	Met	
			195					200					205				
35	Thr	Ile	Leu	Thr	Val	Gln	Leu	Ile	Val	Glu	Phe	Ala	Lys	Gly	Leu	Pro	
		210					215					220					
	Gly	Phe	Ala	Lys	Ile	Ser	Gln	Pro	Asp	Gln	Ile	Thr	Leu	Leu	Lys	Ala	
		225				230					235					240	
40	Cys	Ser	Ser	Glu	Val	Met	Met	Leu	Arg	Val	Ala	Arg	Arg	Tyr	Asp	Ala	
					245					250					255		
	Ala	Ser	Asp	Ser	Val	Leu	Phe	Ala	Asn	Asn	Gln	Ala	Tyr	Thr	Arg	Asp	
				260					265					270			
45	Asn	Tyr	Arg	Lys	Ala	Gly	Met	Ala	Tyr	Val	Ile	Glu	Asp	Leu	Leu	His	
				275				280					285				
	Phe	Cys	Arg	Cys	Met	Tyr	Ser	Met	Ala	Leu	Asp	Asn	Ile	His	Tyr	Ala	
		290					295					300					
50	Leu	Leu	Thr	Ala	Val	Val	Ile	Phe	Ser	Asp	Arg	Pro	Gly	Leu	Glu	Gln	
		305				310					315					320	
	Pro	Gln	Leu	Val	Glu	Glu	Ile	Gln	Arg	Tyr	Tyr	Leu	Asn	Thr	Leu	Arg	

EP 1 266 015 B1

				325					330					335		
	Ile	Tyr	Ile	Leu	Asn	Gln	Leu	Ser	Gly	Ser	Ala	Arg	Ser	Ser	Val	Ile
				340					345					350		
5	Tyr	Gly	Lys	Ile	Leu	Ser	Ile	Leu	Ser	Glu	Leu	Arg	Thr	Leu	Gly	Met
			355					360					365			
	Gln	Asn	Ser	Asn	Met	Cys	Ile	Ser	Leu	Lys	Leu	Lys	Asn	Arg	Lys	Leu
		370					375					380				
10	Pro	Pro	Phe	Leu	Glu	Glu	Ile	Trp	Asp	Val	Ala	Asp	Met	Ser	His	Thr
	385					390					395					400
	Gln	Pro	Pro	Pro	Ile	Leu	Glu	Ser	Pro	Thr	Asn	Leu				
					405					410						
15																
	<210> 12															
	<211> 412															
	<212> PRT															
	<213> Artificial Sequence															
20	<220>															
	<221> misc_feature															
	<223> Novel Sequence															
	<400> 12															
25																
	Lys	Gly	Pro	Ala	Pro	Arg	Gln	Gln	Glu	Glu	Leu	Cys	Leu	Val	Cys	Gly
	1				5					10					15	
	Asp	Arg	Ala	Ser	Gly	Tyr	His	Tyr	Asn	Ala	Leu	Thr	Cys	Glu	Gly	Cys
				20					25					30		
30	Lys	Gly	Phe	Phe	Arg	Arg	Ser	Val	Thr	Lys	Asn	Ala	Val	Tyr	Ile	Cys
			35					40					45			
	Lys	Phe	Gly	His	Ala	Cys	Glu	Met	Asp	Met	Tyr	Met	Arg	Arg	Lys	Cys
		50					55					60				
35	Gln	Glu	Cys	Arg	Leu	Lys	Lys	Cys	Leu	Ala	Val	Gly	Met	Arg	Pro	Glu
	65					70					75					80
	Cys	Val	Val	Pro	Glu	Thr	Gln	Cys	Ala	Met	Lys	Arg	Lys	Glu	Lys	Lys
					85					90					95	
40	Ala	Gln	Lys	Glu	Lys	Asp	Lys	Leu	Pro	Val	Ser	Thr	Thr	Thr	Val	Asp
				100					105						110	
	Asp	His	Met	Pro	Pro	Ile	Met	Gln	Cys	Glu	Pro	Pro	Pro	Pro	Glu	Ala
			115					120					125			
45	Ala	Arg	Ile	His	Glu	Val	Val	Pro	Arg	Phe	Leu	Ser	Asp	Lys	Leu	Leu
		130					135					140				
	Glu	Thr	Asn	Arg	Gln	Lys	Asn	Ile	Pro	Gln	Leu	Thr	Ala	Asn	Gln	Gln
	145					150					155					160
50	Phe	Leu	Ile	Ala	Arg	Leu	Ile	Trp	Tyr	Gln	Asp	Gly	Tyr	Glu	Gln	Pro
				165						170					175	
	Ser	Asp	Glu	Asp	Leu	Lys	Arg	Ile	Thr	Gln	Thr	Trp	Gln	Gln	Ala	Asp
				180					185					190		

55

EP 1 266 015 B1

Asp Glu Asn Glu Glu Ser Asp Thr Pro Phe Arg Gln Ile Thr Glu Met
 195 200 205
 Thr Ile Leu Thr Val Gln Leu Ile Val Glu Phe Ala Lys Gly Leu Pro
 210 215 220
 Gly Phe Ala Lys Ile Ser Gln Pro Asp Gln Ile Thr Leu Leu Lys Ala
 225 230 235 240
 Cys Ser Ser Glu Val Met Met Leu Arg Val Ala Arg Arg Tyr Asp Ala
 245 250 255
 Ala Ser Asp Ser Val Leu Phe Ala Asn Asn Gln Ala Tyr Thr Arg Asp
 260 265 270
 Asn Tyr Arg Lys Ala Gly Met Ala Tyr Val Ile Glu Asp Leu Leu His
 275 280 285
 Phe Cys Arg Cys Met Tyr Ser Met Ala Leu Asp Asn Ile His Tyr Ala
 290 295 300
 Leu Leu Thr Ala Val Val Ile Phe Ser Asp Arg Pro Gly Leu Glu Gln
 305 310 315 320
 Pro Gln Leu Val Glu Glu Ile Gln Arg Tyr Tyr Leu Asn Thr Leu Arg
 325 330 335
 Ile Tyr Ile Leu Asn Gln Leu Ser Gly Ser Ala Arg Ser Ser Val Ile
 340 345 350
 Tyr Gly Lys Ile Leu Ser Ile Leu Ser Glu Leu Arg Thr Leu Gly Met
 355 360 365
 Gln Asn Ser Asn Met Cys Ile Ser Leu Lys Leu Lys Asn Arg Lys Leu
 370 375 380
 Pro Pro Phe Leu Glu Glu Ile Trp Asp Val Ala Asp Met Ser His Thr
 385 390 395 400
 Gln Pro Pro Pro Ile Leu Glu Ser Pro Thr Asn Leu
 405 410

<210> 13
 <211> 334
 <212> PRT
 <213> Artificial Sequence
 <220>
 <221> misc feature
 <223> Novel Sequence
 <400> 13

Pro Glu Cys Val Val Pro Glu Thr Gln Cys Ala Met Lys Arg Lys Glu
 1 5 10 15
 Lys Lys Ala Gln Lys Glu Lys Asp Lys Leu Pro Val Ser Thr Thr Thr
 20 25 30
 Val Asp Asp His Met Pro Pro Ile Met Gln Cys Glu Pro Pro Pro Pro
 35 40 45
 Glu Ala Ala Arg Ile His Glu Val Val Pro Arg Phe Leu Ser Asp Lys
 50 55 60

EP 1 266 015 B1

5 Leu Leu Glu Thr Asn Arg Gln Lys Asn Ile Pro Gln Leu Thr Ala Asn
 65 70 75 80
 Gln Gln Phe Leu Ile Ala Arg Leu Ile Trp Tyr Gln Asp Gly Tyr Glu
 85 90 95
 Gln Pro Ser Asp Glu Asp Leu Lys Arg Ile Thr Gln Thr Trp Gln Gln
 100 105 110
 10 Ala Asp Asp Glu Asn Glu Glu Ser Asp Thr Pro Phe Arg Gln Ile Thr
 115 120 125
 Glu Met Thr Ile Leu Thr Val Gln Leu Ile Val Glu Phe Ala Lys Gly
 130 135 140
 15 Leu Pro Gly Phe Ala Lys Ile Ser Gln Pro Asp Gln Ile Thr Leu Leu
 145 150 155 160
 Lys Ala Cys Ser Ser Glu Val Met Met Leu Arg Val Ala Arg Arg Tyr
 165 170 175
 20 Asp Ala Ala Ser Asp Ser Val Leu Phe Ala Asn Asn Gln Ala Tyr Thr
 180 185 190
 Arg Asp Asn Tyr Arg Lys Ala Gly Met Ala Tyr Val Ile Glu Asp Leu
 195 200 205
 25 Leu His Phe Cys Arg Cys Met Tyr Ser Met Ala Leu Asp Asn Ile His
 210 215 220
 Tyr Ala Leu Leu Thr Ala Val Val Ile Phe Ser Asp Arg Pro Gly Leu
 225 230 235 240
 30 Glu Gln Pro Gln Leu Val Glu Glu Ile Gln Arg Tyr Tyr Leu Asn Thr
 245 250 255
 Leu Arg Ile Tyr Ile Leu Asn Gln Leu Ser Gly Ser Ala Arg Ser Ser
 260 265 270
 35 Val Ile Tyr Gly Lys Ile Leu Ser Ile Leu Ser Glu Leu Arg Thr Leu
 275 280 285
 Gly Met Gln Asn Ser Asn Met Cys Ile Ser Leu Lys Leu Lys Asn Arg
 290 295 300
 40 Lys Leu Pro Pro Phe Leu Glu Glu Ile Trp Asp Val Ala Asp Met Ser
 305 310 315 320
 His Thr Gln Pro Pro Pro Ile Leu Glu Ser Pro Thr Asn Leu
 325 330

<210> 14
 <211> 244
 <212> PRT
 <213> Artificial Sequence
 <220>
 <221> misc feature
 <223> Novel Sequence
 <400> 14

Tyr Gln Asp Gly Tyr Glu Gln Pro Ser Asp Glu Asp Leu Lys Arg Ile

EP 1 266 015 B1

	1		5		10		15									
	Thr	Gln	Thr	Trp	Gln	Gln	Ala	Asp	Asp	Glu	Asn	Glu	Glu	Ser	Asp	Thr
				20					25					30		
5	Pro	Phe	Arg	Gln	Ile	Thr	Glu	Met	Thr	Ile	Leu	Thr	Val	Gln	Leu	Ile
			35					40					45			
	Val	Glu	Phe	Ala	Lys	Gly	Leu	Pro	Gly	Phe	Ala	Lys	Ile	Ser	Gln	Pro
		50					55					60				
10	Asp	Gln	Ile	Thr	Leu	Leu	Lys	Ala	Cys	Ser	Ser	Glu	Val	Met	Met	Leu
	65				70						75					80
	Arg	Val	Ala	Arg	Arg	Tyr	Asp	Ala	Ala	Ser	Asp	Ser	Val	Leu	Phe	Ala
				85						90					95	
15	Asn	Asn	Gln	Ala	Tyr	Thr	Arg	Asp	Asn	Tyr	Arg	Lys	Ala	Gly	Met	Ala
				100					105					110		
	Tyr	Val	Ile	Glu	Asp	Leu	Leu	His	Phe	Cys	Arg	Cys	Met	Tyr	Ser	Met
			115					120					125			
20	Ala	Leu	Asp	Asn	Ile	His	Tyr	Ala	Ile	Leu	Thr	Ala	Val	Val	Ile	Phe
		130					135					140				
	Ser	Asp	Arg	Pro	Gly	Leu	Glu	Gln	Pro	Gln	Leu	Val	Glu	Glu	Ile	Gln
	145				150						155					160
25	Arg	Tyr	Tyr	Leu	Asn	Thr	Leu	Arg	Ile	Tyr	Ile	Leu	Asn	Gln	Leu	Ser
				165						170					175	
	Gly	Ser	Ala	Arg	Ser	Ser	Val	Ile	Tyr	Gly	Lys	Ile	Leu	Ser	Ile	Leu
			180						185					190		
30	Ser	Glu	Leu	Arg	Thr	Leu	Gly	Met	Gln	Asn	Ser	Asn	Met	Cys	Ile	Ser
			195					200					205			
	Leu	Lys	Leu	Lys	Asn	Arg	Lys	Leu	Pro	Pro	Phe	Leu	Glu	Glu	Ile	Trp
		210					215					220				
35	Asp	Val	Ala	Asp	Met	Ser	His	Thr	Gln	Pro	Pro	Pro	Ile	Leu	Glu	Ser
	225					230					235					240
	Pro	Thr	Asn	Leu												

40 <210> 15
 <211> 320
 <212> PRT
 <213> Artificial Sequence
 <220>
 45 <221> misc feature
 <223> Novel Sequence
 <400> 15

50	Pro	Glu	Cys	Val	Val	Pro	Glu	Thr	Gln	Cys	Ala	Met	Lys	Arg	Lys	Glu
	1				5					10					15	
	Lys	Lys	Ala	Gln	Lys	Glu	Lys	Asp	Lys	Leu	Pro	Val	Ser	Thr	Thr	Thr
				20					25					30		

55

EP 1 266 015 B1

Val Asp Asp His Met Pro Pro Ile Met Gln Cys Glu Pro Pro Pro Pro
35 40 45

5 Glu Ala Ala Arg Ile His Glu Val Val Pro Arg Phe Leu Ser Asp Lys
50 55 60

Leu Leu Glu Thr Asn Arg Gln Lys Asn Ile Pro Gln Leu Thr Ala Asn
65 70 75 80

10 Gln Gln Phe Leu Ile Ala Arg Leu Ile Trp Tyr Gln Asp Gly Tyr Glu
85 90 95

Gln Pro Ser Asp Glu Asp Leu Lys Arg Ile Thr Gln Thr Trp Gln Gln
100 105 110

15 Ala Asp Asp Glu Asn Glu Glu Ser Asp Thr Pro Phe Arg Gln Ile Thr
115 120 125

Glu Met Thr Ile Leu Thr Val Gln Leu Ile Val Glu Phe Ala Lys Gly
130 135 140

20 Leu Pro Gly Phe Ala Lys Ile Ser Gln Pro Asp Gln Ile Thr Leu Leu
145 150 155 160

Lys Ala Cys Ser Ser Glu Val Met Met Leu Arg Val Ala Arg Arg Tyr
165 170 175

25 Asp Ala Ala Ser Asp Ser Val Leu Phe Ala Asn Asn Gln Ala Tyr Thr
180 185 190

Arg Asp Asn Tyr Arg Lys Ala Gly Met Ala Tyr Val Ile Glu Asp Leu
195 200 205

30 Leu His Phe Cys Arg Cys Met Tyr Ser Met Ala Leu Asp Asn Ile His
210 215 220

Tyr Ala Leu Leu Thr Ala Val Val Ile Phe Ser Asp Arg Pro Gly Leu
225 230 235 240

Glu Gln Pro Gln Leu Val Glu Glu Ile Gln Arg Tyr Tyr Leu Asn Thr
245 250 255

35 Leu Arg Ile Tyr Ile Leu Asn Gln Leu Ser Gly Ser Ala Arg Ser Ser
260 265 270

Val Ile Tyr Gly Lys Ile Leu Ser Ile Leu Ser Glu Leu Arg Thr Leu
275 280 285

40 Gly Met Gln Asn Ser Asn Met Cys Ile Ser Leu Lys Leu Lys Asn Arg
290 295 300

Lys Leu Pro Pro Phe Leu Glu Glu Ile Trp Asp Val Ala Asp Met Ser
305 310 315 320

<210> 16
<211> 625
<212> PRT
<213> Artificial Sequence
50 <220>
<221> misc feature
<223> Novel Sequence
<400> 16

EP 1 266 015 B1

Gly Pro Ala Pro Arg Val Gln Glu Glu Leu Cys Leu Val Cys Gly Asp
 1 5 10 15
 Arg Ala Ser Gly Tyr His Tyr Asn Ala Leu Thr Cys Glu Gly Cys Lys
 5 20 25 30
 Gly Phe Phe Arg Arg Ser Val Thr Lys Ser Ala Val Tyr Cys Cys Lys
 35 40 45
 Phe Gly Arg Ala Cys Glu Met Asp Met Tyr Met Arg Arg Lys Cys Gln
 10 50 55 60
 Glu Cys Arg Leu Lys Lys Cys Leu Ala Val Gly Met Arg Pro Glu Cys
 65 70 75 80
 Val Val Pro Glu Asn Gln Cys Ala Met Lys Arg Arg Glu Lys Lys Ala
 15 85 90 95
 Gln Lys Glu Lys Asp Lys Met Thr Thr Ser Pro Ser Ser Gln His Gly
 100 105 110
 Gly Asn Gly Ser Leu Ala Ser Gly Gly Gly Gln Asp Phe Val Lys Lys
 115 120 125
 Glu Ile Leu Asp Leu Met Thr Cys Glu Pro Pro Gln His Ala Thr Ile
 130 135 140
 Pro Leu Leu Pro Asp Glu Ile Leu Ala Lys Cys Gln Ala Arg Asn Ile
 145 150 155 160
 Pro Ser Leu Thr Tyr Asn Gln Leu Ala Val Ile Tyr Lys Leu Ile Trp
 165 170 175
 Tyr Gln Asp Gly Tyr Glu Gln Pro Ser Glu Glu Asp Leu Arg Arg Ile
 180 185 190
 Met Ser Gln Pro Asp Glu Asn Glu Ser Gln Thr Asp Val Ser Phe Arg
 195 200 205
 His Ile Thr Glu Ile Thr Ile Leu Thr Val Gln Leu Ile Val Glu Phe
 210 215 220
 Ala Lys Gly Leu Pro Ala Phe Thr Lys Ile Pro Gln Glu Asp Gln Ile
 225 230 235 240
 Thr Leu Leu Lys Ala Cys Ser Ser Glu Val Met Met Leu Arg Met Ala
 245 250 255
 Arg Arg Tyr Asp His Ser Ser Asp Ser Ile Phe Phe Ala Asn Asn Arg
 260 265 270
 Ser Tyr Thr Arg Asp Ser Tyr Lys Met Ala Gly Met Ala Asp Asn Ile
 275 280 285
 Glu Asp Leu Leu His Phe Cys Arg Gln Met Phe Ser Met Lys Val Asp
 290 295 300
 Asn Val Glu Tyr Ala Leu Leu Thr Ala Ile Val Ile Phe Ser Asp Arg
 305 310 315 320
 Pro Gly Leu Glu Lys Ala Gln Leu Val Glu Ala Ile Gln Ser Tyr Tyr
 325 330 335
 Ile Asp Thr Leu Arg Ile Tyr Ile Leu Asn Arg His Cys Gly Asp Ser

EP 1 266 015 B1

	340	345	350
	Met Ser Leu Val Phe Tyr Ala Lys Leu Leu Ser Ile Leu Thr Glu Leu		
	355	360	365
5	Arg Thr Leu Gly Asn Gln Asn Ala Glu Met Cys Phe Ser Leu Lys Leu		
	370	375	380
	Lys Asn Arg Lys Leu Pro Lys Phe Leu Glu Glu Ile Trp Asp Val His		
	385	390	395
10	Ala Ile Pro Pro Ser Val Gln Ser His Leu Gln Ile Thr Gln Glu Glu		
	405	410	415
	Asn Glu Arg Leu Glu Arg Ala Glu Arg Met Arg Ala Ser Val Gly Gly		
	420	425	430
15	Ala Ile Thr Ala Gly Ile Asp Cys Asp Ser Ala Ser Thr Ser Ala Ala		
	435	440	445
	Ala Ala Ala Ala Gln His Gln Pro Gln Pro Gln Pro Gln Pro Gln Pro		
	450	455	460
20	Ser Ser Leu Thr Gln Asn Asp Ser Gln His Gln Thr Gln Pro Gln Leu		
	465	470	475
	Gln Pro Gln Leu Pro Pro Gln Leu Gln Gly Gln Leu Gln Pro Gln Leu		
	485	490	495
25	Gln Pro Gln Leu Gln Thr Gln Leu Gln Pro Gln Ile Gln Pro Gln Pro		
	500	505	510
	Gln Leu Leu Pro Val Ser Ala Pro Val Pro Ala Ser Val Thr Ala Pro		
	515	520	525
30	Gly Ser Leu Ser Ala Val Ser Thr Ser Ser Glu Tyr Met Gly Gly Ser		
	530	535	540
	Ala Ala Ile Gly Pro Ile Thr Pro Ala Thr Thr Ser Ser Ile Thr Ala		
	545	550	555
35	Ala Val Thr Ala Ser Ser Thr Thr Ser Ala Val Pro Met Gly Asn Gly		
	565	570	575
	Val Gly Val Gly Val Gly Val Gly Gly Asn Val Ser Met Tyr Ala Asn		
	580	585	590
40	Ala Gln Thr Ala Met Ala Leu Met Gly Val Ala Leu His Ser His Gln		
	595	600	605
	Glu Gln Leu Ile Gly Gly Val Ala Val Lys Ser Glu His Ser Thr Thr		
	610	615	620
45	Ala		
	625		

<210> 17
 <211> 583
 <212> PRT
 <213> Artificial Sequence
 <220>
 <221> misc feature
 <223> Novel Sequence
 <400> 17

EP 1 266 015 B1

Ala Val Tyr Cys Cys Lys Phe Gly Arg Ala Cys Glu Met Asp Met Tyr
1 5 10 15

Met Arg Arg Lys Cys Gln Glu Cys Arg Leu Lys Lys Cys Leu Ala Val
5 20 25 30

Gly Met Arg Pro Glu Cys Val Val Pro Glu Asn Gln Cys Ala Met Lys
35 40 45

Arg Arg Glu Lys Lys Ala Gln Lys Glu Lys Asp Lys Met Thr Thr Ser
10 50 55 60

Pro Ser Ser Gln His Gly Gly Asn Gly Ser Leu Ala Ser Gly Gly Gly
65 70 75 80

Gln Asp Phe Val Lys Lys Glu Ile Leu Asp Leu Met Thr Cys Glu Pro
15 85 90 95

Pro Gln His Ala Thr Ile Pro Leu Leu Pro Asp Glu Ile Leu Ala Lys
100 105 110

Cys Gln Ala Arg Asn Ile Pro Ser Leu Thr Tyr Asn Gln Leu Ala Val
20 115 120 125

Ile Tyr Lys Leu Ile Trp Tyr Gln Asp Gly Tyr Glu Gln Pro Ser Glu
130 135 140

Glu Asp Leu Arg Arg Ile Met Ser Gln Pro Asp Glu Asn Glu Ser Gln
25 145 150 155 160

Thr Asp Val Ser Phe Arg His Ile Thr Glu Ile Thr Ile Leu Thr Val
165 170 175

Gln Leu Ile Val Glu Phe Ala Lys Gly Leu Pro Ala Phe Thr Lys Ile
30 180 185 190

Pro Gln Glu Asp Gln Ile Thr Leu Leu Lys Ala Cys Ser Ser Glu Val
195 200 205

Met Met Leu Arg Met Ala Arg Arg Tyr Asp His Ser Ser Asp Ser Ile
35 210 215 220

Phe Phe Ala Asn Asn Arg Ser Tyr Thr Arg Asp Ser Tyr Lys Met Ala
225 230 235 240

Gly Met Ala Asp Asn Ile Glu Asp Leu Leu His Phe Cys Arg Gln Met
245 250 255

Phe Ser Met Lys Val Asp Asn Val Glu Tyr Ala Leu Leu Thr Ala Ile
40 260 265 270

Val Ile Phe Ser Asp Arg Pro Gly Leu Glu Lys Ala Gln Leu Val Glu
275 280 285

Ala Ile Gln Ser Tyr Tyr Ile Asp Thr Leu Arg Ile Tyr Ile Leu Asn
45 290 295 300

Arg His Cys Gly Asp Ser Met Ser Leu Val Phe Tyr Ala Lys Leu Leu
305 310 315 320

Ser Ile Leu Thr Glu Leu Arg Thr Leu Gly Asn Gln Asn Ala Glu Met
50 325 330 335

EP 1 266 015 B1

Cys Phe Ser Leu Lys Leu Lys Asn Arg Lys Leu Pro Lys Phe Leu Glu
 340 345 350
 Glu Ile Trp Asp Val His Ala Ile Pro Pro Ser Val Gln Ser His Leu
 5 355 360 365
 Gln Ile Thr Gln Glu Glu Asn Glu Arg Leu Glu Arg Ala Glu Arg Met
 370 375 380
 Arg Ala Ser Val Gly Gly Ala Ile Thr Ala Gly Ile Asp Cys Asp Ser
 10 385 390 395 400
 Ala Ser Thr Ser Ala Ala Ala Ala Ala Ala Gln His Gln Pro Gln Pro
 405 410 415
 Gln Pro Gln Pro Gln Pro Ser Ser Leu Thr Gln Asn Asp Ser Gln His
 15 420 425 430
 Gln Thr Gln Pro Gln Leu Gln Pro Gln Leu Pro Pro Gln Leu Gln Gly
 435 440 445
 Gln Leu Gln Pro Gln Leu Gln Pro Gln Leu Gln Thr Gln Leu Gln Pro
 20 450 455 460
 Gln Ile Gln Pro Gln Pro Gln Leu Leu Pro Val Ser Ala Pro Val Pro
 465 470 475 480
 Ala Ser Val Thr Ala Pro Gly Ser Leu Ser Ala Val Ser Thr Ser Ser
 25 485 490 495
 Glu Tyr Met Gly Gly Ser Ala Ala Ile Gly Pro Ile Thr Pro Ala Thr
 500 505 510
 Thr Ser Ser Ile Thr Ala Ala Val Thr Ala Ser Ser Thr Thr Ser Ala
 30 515 520 525
 Val Pro Met Gly Asn Gly Val Gly Val Gly Val Gly Val Gly Gly Asn
 530 535 540
 Val Ser Met Tyr Ala Asn Ala Gln Thr Ala Met Ala Leu Met Gly Val
 545 550 555 560
 Ala Leu His Ser His Gln Glu Gln Leu Ile Gly Gly Val Ala Val Lys
 565 570 575
 Ser Glu His Ser Thr Thr Ala
 580

<210> 18
 <211> 549
 <212> PRT
 <213> Artificial Sequence
 <400> 18

Arg Pro Glu Cys Val Val Pro Glu Asn Gln Cys Ala Met Lys Arg Arg
 1 5 10 15
 Glu Lys Lys Ala Gln Lys Glu Lys Asp Lys Met Thr Thr Ser Pro Ser
 20 25 30
 Ser Gln His Gly Gly Asn Gly Ser Leu Ala Ser Gly Gly Gly Gln Asp
 35 40 45

EP 1 266 015 B1

	Phe	Val	Lys	Lys	Glu	Ile	Leu	Asp	Leu	Met	Thr	Cys	Glu	Pro	Pro	Gln
	50						55					60				
5	His	Ala	Thr	Ile	Pro	Leu	Leu	Pro	Asp	Glu	Ile	Leu	Ala	Lys	Cys	Gln
	65					70					75					80
	Ala	Arg	Asn	Ile	Pro	Ser	Leu	Thr	Tyr	Asn	Gln	Leu	Ala	Val	Ile	Tyr
					85					90					95	
10	Lys	Leu	Ile	Trp	Tyr	Gln	Asp	Gly	Tyr	Glu	Gln	Pro	Ser	Glu	Glu	Asp
				100					105					110		
	Leu	Arg	Arg	Ile	Met	Ser	Gln	Pro	Asp	Glu	Asn	Glu	Ser	Gln	Thr	Asp
			115					120					125			
15	Val	Ser	Phe	Arg	His	Ile	Thr	Glu	Ile	Thr	Ile	Leu	Thr	Val	Gln	Leu
	130						135					140				
	Ile	Val	Glu	Phe	Ala	Lys	Gly	Leu	Pro	Ala	Phe	Thr	Lys	Ile	Pro	Gln
	145					150					155					160
20	Glu	Asp	Gln	Ile	Thr	Leu	Leu	Lys	Ala	Cys	Ser	Ser	Glu	Val	Met	Met
					165					170					175	
	Leu	Arg	Met	Ala	Arg	Arg	Tyr	Asp	His	Ser	Ser	Asp	Ser	Ile	Phe	Phe
				180					185					190		
25	Ala	Asn	Asn	Arg	Ser	Tyr	Thr	Arg	Asp	Ser	Tyr	Lys	Met	Ala	Gly	Met
			195					200					205			
	Ala	Asp	Asn	Ile	Glu	Asp	Leu	Leu	His	Phe	Cys	Arg	Gln	Met	Phe	Ser
	210						215					220				
30	Met	Lys	Val	Asp	Asn	Val	Glu	Tyr	Ala	Leu	Leu	Thr	Ala	Ile	Val	Ile
	225					230					235					240
	Phe	Ser	Asp	Arg	Pro	Gly	Leu	Glu	Lys	Ala	Gln	Leu	Val	Glu	Ala	Ile
					245					250					255	
35	Gln	Ser	Tyr	Tyr	Ile	Asp	Thr	Leu	Arg	Ile	Tyr	Ile	Leu	Asn	Arg	His
				260					265					270		
	Cys	Gly	Asp	Ser	Met	Ser	Leu	Val	Phe	Tyr	Ala	Lys	Leu	Leu	Ser	Ile
			275					280					285			
40	Leu	Thr	Glu	Leu	Arg	Thr	Leu	Gly	Asn	Gln	Asn	Ala	Glu	Met	Cys	Phe
	290						295					300				
	Ser	Leu	Lys	Leu	Lys	Asn	Arg	Lys	Leu	Pro	Lys	Phe	Leu	Glu	Glu	Ile
	305					310					315					320
	Trp	Asp	Val	His	Ala	Ile	Pro	Pro	Ser	Val	Gln	Ser	His	Leu	Gln	Ile
					325					330					335	
45	Thr	Gln	Glu	Glu	Asn	Glu	Arg	Leu	Glu	Arg	Ala	Glu	Arg	Met	Arg	Ala
					340				345					350		
	Ser	Val	Gly	Gly	Ala	Ile	Thr	Ala	Gly	Ile	Asp	Cys	Asp	Ser	Ala	Ser
			355					360					365			
50	Thr	Ser	Ala	Ala	Ala	Ala	Ala	Ala	Gln	His	Gln	Pro	Gln	Pro	Gln	Pro
	370						375					380				
	Gln	Pro	Gln	Pro	Ser	Ser	Leu	Thr	Gln	Asn	Asp	Ser	Gln	His	Gln	Thr
	385					390					395					400

EP 1 266 015 B1

5 Gln Pro Gln Leu Gln Pro Gln Leu Pro Pro Gln Leu Gln Gly Gln Leu
 405 410 415
 Gln Pro Gln Leu Gln Pro Gln Leu Gln Thr Gln Leu Gln Pro Gln Ile
 420 425 430
 Gln Pro Gln Pro Gln Leu Leu Pro Val Ser Ala Pro Val Pro Ala Ser
 435 440 445
 10 Val Thr Ala Pro Gly Ser Leu Ser Ala Val Ser Thr Ser Ser Glu Tyr
 450 455 460
 Met Gly Gly Ser Ala Ala Ile Gly Pro Ile Thr Pro Ala Thr Thr Ser
 465 470 475 480
 15 Ser Ile Thr Ala Ala Val Thr Ala Ser Ser Thr Thr Ser Ala Val Pro
 485 490 495
 Met Gly Asn Gly Val Gly Val Gly Val Gly Val Gly Gly Asn Val Ser
 500 505 510
 20 Met Tyr Ala Asn Ala Gln Thr Ala Met Ala Leu Met Gly Val Ala Leu
 515 520 525
 His Ser His Gln Glu Gln Leu Ile Gly Gly Val Ala Val Lys Ser Glu
 530 535 540
 25 His Ser Thr Thr Ala
 545

<210> 19

<211> 445

<212> PRT

<213> Artificial Sequence

<400> 19

35 Tyr Glu Gln Pro Ser Glu Glu Asp Leu Arg Arg Ile Met Ser Gln Pro
 1 5 10 15
 Asp Glu Asn Glu Ser Gln Thr Asp Val Ser Phe Arg His Ile Thr Glu
 20 25 30
 40 Ile Thr Ile Leu Thr Val Gln Leu Ile Val Glu Phe Ala Lys Gly Leu
 35 40 45
 Pro Ala Phe Thr Lys Ile Pro Gln Glu Asp Gln Ile Thr Leu Leu Lys
 50 55 60
 45 Ala Cys Ser Ser Glu Val Met Met Leu Arg Met Ala Arg Arg Tyr Asp
 65 70 75 80
 His Ser Ser Asp Ser Ile Phe Phe Ala Asn Asn Arg Ser Tyr Thr Arg
 85 90 95
 50 Asp Ser Tyr Lys Met Ala Gly Met Ala Asp Asn Ile Glu Asp Leu Leu
 100 105 110
 His Phe Cys Arg Gln Met Phe Ser Met Lys Val Asp Asn Val Glu Tyr
 115 120 125
 55 Ala Leu Leu Thr Ala Ile Val Ile Phe Ser Asp Arg Pro Gly Leu Glu
 130 135 140

EP 1 266 015 B1

Lys Ala Gln Leu Val Glu Ala Ile Gln Ser Tyr Tyr Ile Asp Thr Leu
 145 150 155 160
 Arg Ile Tyr Ile Leu Asn Arg His Cys Gly Asp Ser Met Ser Leu Val
 165 170 175
 Phe Tyr Ala Lys Leu Leu Ser Ile Leu Thr Glu Leu Arg Thr Leu Gly
 180 185 190
 Asn Gln Asn Ala Glu Met Cys Phe Ser Leu Lys Leu Lys Asn Arg Lys
 195 200 205
 Leu Pro Lys Phe Leu Glu Glu Ile Trp Asp Val His Ala Ile Pro Pro
 210 215 220
 Ser Val Gln Ser His Leu Gln Ile Thr Gln Glu Glu Asn Glu Arg Leu
 225 230 235 240
 Glu Arg Ala Glu Arg Met Arg Ala Ser Val Gly Gly Ala Ile Thr Ala
 245 250 255
 Gly Ile Asp Cys Asp Ser Ala Ser Thr Ser Ala Ala Ala Ala Ala Ala
 260 265 270
 Gln His Gln Pro Gln Pro Gln Pro Gln Pro Gln Pro Ser Ser Leu Thr
 275 280 285
 Gln Asn Asp Ser Gln His Gln Thr Gln Pro Gln Leu Gln Pro Gln Leu
 290 295 300
 Pro Pro Gln Leu Gln Gly Gln Leu Gln Pro Gln Leu Gln Pro Gln Leu
 305 310 315 320
 Gln Thr Gln Leu Gln Pro Gln Ile Gln Pro Gln Pro Gln Leu Leu Pro
 325 330 335
 Val Ser Ala Pro Val Pro Ala Ser Val Thr Ala Pro Gly Ser Leu Ser
 340 345 350
 Ala Val Ser Thr Ser Ser Glu Tyr Met Gly Gly Ser Ala Ala Ile Gly
 355 360 365
 Pro Ile Thr Pro Ala Thr Thr Ser Ser Ile Thr Ala Ala Val Thr Ala
 370 375 380
 Ser Ser Thr Thr Ser Ala Val Pro Met Gly Asn Gly Val Gly Val Gly
 385 390 395 400
 Val Gly Val Gly Gly Asn Val Ser Met Tyr Ala Asn Ala Gln Thr Ala
 405 410 415
 Met Ala Leu Met Gly Val Ala Leu His Ser His Gln Glu Gln Leu Ile
 420 425 430
 Gly Gly Val Ala Val Lys Ser Glu His Ser Thr Thr Ala
 435 440 445

<210> 20

<211> 323

<212> PRT

<213> Artificial Sequence

<400> 20

Arg Pro Glu Cys Val Val Pro Glu Asn Gln Cys Ala Met Lys Arg Arg

EP 1 266 015 B1

	1		5		10		15									
	Glu	Lys	Lys	Ala	Gln	Lys	Glu	Lys	Asp	Lys	Met	Thr	Thr	Ser	Pro	Ser
				20					25					30		
5	Ser	Gln	His	Gly	Gly	Asn	Gly	Ser	Leu	Ala	Ser	Gly	Gly	Gly	Gln	Asp
			35					40					45			
	Phe	Val	Lys	Lys	Glu	Ile	Leu	Asp	Leu	Met	Thr	Cys	Glu	Pro	Pro	Gln
		50					55					60				
10	His	Ala	Thr	Ile	Pro	Leu	Leu	Pro	Asp	Glu	Ile	Leu	Ala	Lys	Cys	Gln
	65					70					75					80
	Ala	Arg	Asn	Ile	Pro	Ser	Leu	Thr	Tyr	Asn	Gln	Leu	Ala	Val	Ile	Tyr
				85						90					95	
15	Lys	Leu	Ile	Trp	Tyr	Gln	Asp	Gly	Tyr	Glu	Gln	Pro	Ser	Glu	Glu	Asp
				100					105					110		
	Leu	Arg	Arg	Ile	Met	Ser	Gln	Pro	Asp	Glu	Asn	Glu	Ser	Gln	Thr	Asp
			115					120					125			
20	Val	Ser	Phe	Arg	His	Ile	Thr	Glu	Ile	Thr	Ile	Leu	Thr	Val	Gln	Leu
		130					135					140				
	Ile	Val	Glu	Phe	Ala	Lys	Gly	Leu	Pro	Ala	Phe	Thr	Lys	Ile	Pro	Gln
	145					150					155					160
25	Glu	Asp	Gln	Ile	Thr	Leu	Leu	Lys	Ala	Cys	Ser	Ser	Glu	Val	Met	Met
				165						170					175	
	Leu	Arg	Met	Ala	Arg	Arg	Tyr	Asp	His	Ser	Ser	Asp	Ser	Ile	Phe	Phe
				180					185					190		
30	Ala	Asn	Asn	Arg	Ser	Tyr	Thr	Arg	Asp	Ser	Tyr	Lys	Met	Ala	Gly	Met
			195					200					205			
	Ala	Asp	Asn	Ile	Glu	Asp	Leu	Leu	His	Phe	Cys	Arg	Gln	Met	Phe	Ser
		210					215					220				
35	Met	Lys	Val	Asp	Asn	Val	Glu	Tyr	Ala	Leu	Leu	Thr	Ala	Ile	Val	Ile
	225					230					235					240
	Phe	Ser	Asp	Arg	Pro	Gly	Leu	Glu	Lys	Ala	Gln	Leu	Val	Glu	Ala	Ile
				245						250					255	
40	Gln	Ser	Tyr	Tyr	Ile	Asp	Thr	Leu	Arg	Ile	Tyr	Ile	Leu	Asn	Arg	His
				260					265					270		
	Cys	Gly	Asp	Ser	Met	Ser	Leu	Val	Phe	Tyr	Ala	Lys	Leu	Leu	Ser	Ile
			275					280					285			
45	Leu	Thr	Glu	Leu	Arg	Thr	Leu	Gly	Asn	Gln	Asn	Ala	Glu	Met	Cys	Phe
		290					295					300				
	Ser	Leu	Lys	Leu	Lys	Asn	Arg	Lys	Leu	Pro	Lys	Phe	Leu	Glu	Glu	Ile
	305					310					315					320
50	Trp	Asp	Val													

<210> 21

<211> 987

<212> DNA

<213> Artificial Sequence

<400> 21

EP 1 266 015 B1

5 tgtgctatct gtggggaccg ctcctcaggc aaacactatg gggatacag ttgtgagggc 60
 tgcaagggtc tcttcaagag gacagtacgc aaagacctga cctacacctg ccgagacaac 120
 aaggactgcc tgatcgacaa gagacagcgg aaccgggtgtc agtactgccg ctaccagaag 180
 tgcctggcca tgggcatgaa gcgggaagct gtgcaggagg agcggcagcg gggcaaggac 240
 cggaatgaga acgaggtgga gtccaccagc agtgccaacg aggacatgcc tgtagagaag 300
 10 attctggaag ccgagcttgc tgtcgagccc aagactgaga catacgtgga ggcaaaccatg 360
 gggctgaacc ccagctcacc aaatgaccct gttaccaaca tctgtcaagc agcagacaag 420
 cagctcttca ctcttgtgga gtgggccaag aggatccac acttttctga gctgccccta 480
 15 gacgaccagg tcatcctgct acgggcaggc tggaacgagc tgctgatcgc ctcttctctcc 540
 caccgctcca tagctgtgaa agatgggatt ctcttgcca ccggcctgca cgtacaccgg 600
 aacagcgctc acagtgtctg ggtgggcgcc atctttgaca ggggtgctaac agagctggtg 660
 20 tctaagatgc gtgacatgca gatggacaag acggagctgg gctgcctgcg agccattgtc 720
 ctgttcaacc ctgactctaa ggggctctca aaccctgctg aggtggaggc gttgagggag 780
 aaggtgtatg cgtcactaga agcgtactgc aaacacaagt accctgagca gccgggcagg 840
 25 ttgccaagc tgetgctccg cctgcctgca ctgcgttcca tggggctcaa gtgcctggag 900
 cacctgttct tcttcaagct catcggggac acgccatcg acaccttct catggagatg 960
 ctggaggcac cacatcaagc cacctag 987

30 <210> 22
 <211> 789
 <212> DNA
 <213> Artificial Sequence
 <400> 22

35 aagcgggaag ctgtgcagga ggagcggcag cggggcaagg accggaatga gaacgaggtg 60
 gagtccacca gcagtccaa cgaggacatg cctgtagaga agattctgga agccgagctt 120
 40 gctgtcgagc ccaagactga gacatacgtg gaggcaaaca tggggctgaa cccagctca 180
 ccaaatgacc ctgttaccaa catctgtcaa gcagcagaca agcagctctt cactcttgtg 240
 gagtgggcca agaggatccc acacttttct gagctgcccc tagacgacca ggtcatcctg 300
 45 ctacgggcag gctggaacga gctgctgac gcctccttct cccaccgctc catagctgtg 360
 aaagatggga ttctcctggc caccggcctg cacgtacacc ggaacagcgc tcacagtget 420
 ggggtgggcg ccatctttga cagggtgcta acagagctgg tgtctaagat gcgtgacatg 480
 50 cagatggaca agacggagct gggctgcctg cgagccattg tcctgttcaa ccctgactct 540
 aaggggctct caaaccctgc tgaggtggag gcgttgaggg agaaggtgta tgcgtcacta 600
 gaagcgtact gcaaacacaa gtaccctgag cagccgggca ggtttgcaa gctgctgctc 660

55

EP 1 266 015 B1

cgctgcctg cactgcgttc catcgggctc aagtgcctgg agcacctgtt cttcttcaag 720
 ctcacgcggg acacgcccat cgacaccttc ctcacggaga tgctggaggc accacatcaa 780
 5 gccacctag 789

<210> 23
 <211> 714
 <212> DNA
 <213> Artificial Sequence
 <400> 23

15 gccaacgagg acatgcctgt agagaagatt ctggaagccg agcttgctgt cgagcccaag 60
 actgagacat acgtggaggc aaacatgggg ctgaaccca gctcaccaa tgaccctgtt 120
 accaacatct gtcaagcagc agacaagcag ctcttcactc ttgtggagtg ggccaagagg 180
 20 atcccacact tttctgagct gcccctagac gaccaggcca tcctgctacg ggagggctgg 240
 aacgagctgc tgatcgcttc cttctccac cgctccatag ctgtgaaaga tgggattctc 300
 ctggccaccg gcctgcacgt acaccggaac agcgctcaca gtgtgggggt gggcgccatc 360
 25 tttgacaggg tgctaacaga gctggtgtct aagatgcgtg acatgcagat ggacaagacg 420
 gagctgggct gcctgcgagc cattgtcctg ttcaaccctg actctaagg gctctcaaac 480
 cctgctgagg tggaggcgtt gagggagaag gtgtatgcgt cactagaagc gtactgcaaa 540
 30 cacaagtacc ctgagcagcc gggcagggtt gccaaagctgc tgctccgct gcctgcactg 600
 cgttccatcg ggctcaagtg cctggagcac ctgttcttct tcaagctcat cggggacacg 660
 cccatcgaca ccttctcat ggagatgctg gaggcaccac atcaagccac ctag 714

<210> 24
 <211> 536
 <212> DNA
 <213> Artificial Sequence
 <400> 24

35 ggatcccaca cttttctgag ctgcccctag acgaccaggc catcctgcta cgggcaggct 60
 ggaacgagct gctgatcgcc tccttctccc accgctccat agctgtgaaa gatgggatc 120
 45 tcctggccac cggcctgcac gtacaccgga acagcgctca cagtgtggg gtggcgcca 180
 tctttgacag ggtgctaaca gagctggtgt ctaagatgcg tgacatgcag atggacaaga 240
 cggagctggg ctgcctgcga gccattgtcc tgttcaacc tgactctaag gggctctcaa 300
 50 accctgctga ggtggaggcg ttgagggaga aggtgtatgc gtcactagaa gcgtactgca 360
 aacacaagta ccctgagcag ccgggcaggc ttgccaagct gctgctccgc ctgcctgcac 420
 tgcgttccat cgggctcaag tgccctggagc acctgttctt cttcaagctc atcggggaca 480
 55 cgcccatoga caccttctc atggagatgc tggaggcacc acatcaagcc acctag 536

<210> 25
 <211> 672

EP 1 266 015 B1

<212> DNA

<213> Artificial Sequence

<400> 25

```

5      gccaacgagg acatgcctgt agagaagatt ctggaagccg agcttgctgt cgagcccaag      60
      actgagacat acgtggaggc aaacatgggg ctgaaccca gctcaccaaa tgaccctgtt      120
      accaacatct gtcaagcagc agacaagcag ctcttcactc ttgtggagtg ggccaagagg      180
10     atcccacact tttctgagct gcccctagac gaccaggtea tcctgctacg ggcaggctgg      240
      aacgagctgc tgatgcctc cttetcccac cgctccatag ctgtgaaaga tgggattctc      300
      ctggccaccg gcctgcacgt acaccggaac agcgctcaca gtgctggggg gggcgccatc      360
15     tttgacaggg tgctaacaga gctggtgtct aagatgcgtg acatgcagat ggacaagacg      420
      gagctgggct gcctgcgagc cattgtcctg ttcaaccctg actctaaggg gctctcaaac      480
      cctgctgagg tggaggcggt gagggagaag gtgtatgcgt cactagaagc gtactgcaaa      540
20     cacaagtacc ctgagcagcc gggcaggttt gccaagctgc tgctccgct gcctgcactg      600
      cgttccatcg ggctcaagtg cctggagcac ctgttcttct tcaagctcat cggggacacg      660
      cccatcgaca cc                                     672

```

<210> 26

<211> 1123

<212> DNA

<213> Artificial Sequence

<400> 26

```

      tgcgccatct gcggggaccg ctccctcaggc aagcactatg gagtgtacag ctgcgagggg      60
      tgcaagggct tcttcaagcg gacggtgcgc aaggacctga cctacacctg ccgcgacaac      120
35     aaggactgcc tgattgacaa gcggcagcgg aaccggtgcc agtactgccg ctaccagaag      180
      tgccctggcca tgggcatgaa gcgggaagcc gtgcaggagg agcggcagcg tggcaaggac      240
      cggaacgaga atgaggtgga gtcgaccagc agcgccaacg aggacatgcc ggtggagagg      300
40     atcctggagg ctgagctggc cgtggagccc aagaccgaga cctacgtgga ggcaaaccatg      360
      gggctgaacc ccagctcgcc gaacgacctt gtcaccaaca tttgccaagc agccgacaaa      420
      cagcttttca ccctggtgga gtgggccaag cggatcccac acttctcaga gctgcccctg      480
45     gacgaccagg tcctcctgct gcgggcaggc tggaatgagc tgctcatcgc ctcttctcc      540
      caccgtcca tcgccgtgaa ggacgggatc ctctggcca ccgggctgca cgtccaccgg      600
      aacagcgccc acagcgcagg ggtgggcgcc atctttgaca gggctgctgac ggagcttggt      660
50     tccaagatgc gggacatgca gatggacaag acggagctgg gctgcctgcg cgccatcgtc      720
      ctctttaacc ctgactccaa ggggctctcg aaccggccg aggtggaggc gctgagggag      780
      aaggtctatg cgtccttgga ggcctactgc aagcacaagt acccagagca gccgggaagg      840
55     ttcgctaagc tcttgctccg cctgccggct ctgcgctcca tcgggctcaa atgcctggaa      900

```

EP 1 266 015 B1

catctcttct tcttcaagct catcggggac acaccattg acaccttct tatggagatg 960
 ctggaggcgc cgcaccaaatt gacttaggcc tgcgggcca tcctttgtgc ccaccgttc 1020
 5 tggccaccct gcctggacgc cagctgttct tctcagcctg agccctgtcc ctgcccttct 1080
 ctgcctggcc tgtttggact ttggggcaca gcctgtcact gct 1123

10 <210> 27
 <211> 925
 <212> DNA
 <213> Artificial Sequence
 <400> 27

15 aagcgggaag ccgtgcagga ggagcggcag cgtggcaagg accggaacga gaatgaggtg 60
 gagtcgacca gcagcgccaa cgaggacatg ccggtggaga ggatcctgga ggctgagctg 120
 20 gccgtggagc ccaagaccga gacctacgtg gaggcaaaca tggggctgaa cccagctcg 180
 ccgaacgacc ctgtcaccaa cttttgccaa gcagccgaca aacagctttt caccctgggtg 240
 gagtggggcca agcggatccc acacttctca gagctgcccc tggacgacca ggatcctctg 300
 25 ctgcggggcag gctggaatga gctgtctatc gcctccttct cccaccgtc catcgccgtg 360
 aaggacggga tcctcctggc caccgggctg cactgccacc ggaacagcgc ccacagcgca 420
 ggggtggggc ccatctttga cagggtgctg acggagcttg tgtccaagat gcgggacatg 480
 30 cagatggaca agacggagct gggctgcctg cgcgccatcg tcctctttaa ccctgactcc 540
 aaggggtctc cgaacccggc cgagggtggag gcgctgaggg agaaggtcta tgcgtccttg 600
 gaggcctact gcaagcaca gtacccagag cagccgggaa ggctcgctaa gctcttgctc 660
 35 cgcctgccgg ctctgcgctc catcgggctc aaatgcctgg aacatctctt cttcttcaag 720
 ctcatcgggg acacacccat tgacaccttc cttatggaga tgctggaggc gccgcaccaa 780
 atgacttagg cctgcggggc catcctttgt gcccaccctg tctggccacc ctgcctggac 840
 40 gccagctgtt cttctcagcc tgagccctgt ccctgccctt ctctgcctgg cctgtttgga 900
 ctttggggca cagcctgtca ctgct 925

45 <210> 28
 <211> 850
 <212> DNA
 <213> Artificial Sequence
 <400> 28

50 gccaacgagg acatgccggt ggagaggatc ctggaggctg agctggccgt ggagcccaag 60
 accgagacct acgtggaggc aaacatgggg ctgaaccca gctcgccgaa cgaccctgtc 120
 accaacattt gccaaagcgc cgacaaacag cttttcaacc tgggtggagtg ggccaagcgg 180
 55 atcccacact tctcagagct gcccctggac gaccagggtca tcctgctgcg ggcaggctgg 240
 aatgagctgc tcatgcctc cttctccac cgtccatcg ccgtgaagga cgggatcctc 300

EP 1 266 015 B1

ctggccaccg ggctgcacgt ccaccggaac agcgcccaca gcgcaggggt gggcgccatc 360
 tttgacaggg tgctgacgga gcttgtgtcc aagatgcggg acatgcagat ggacaagacg 420
 5 gagctgggct gcctgcgcgc catcgtcctc ttttaaccctg actccaaggg gctctcgaac 480
 ccggccgagg tggaggcgct gagggagaag gtctatgcgt ccttggaggc ctactgcaag 540
 cacaagtacc cagagcagcc gggaagggtc gctaagctct tgctccgcct gccggtcttg 600
 10 cgctccatcg ggctcaaatg cctggaacat ctcttcttct tcaagctcat cggggacaca 660
 cccattgaca ccttccttat ggagatgctg gaggcgcgc accaaatgac ttaggcctgc 720
 gggcccatcc tttgtgcca cccgttcttg ccaccctgcc tggacgccag ctgttcttct 780
 15 cagcctgagc cctgtccctg cccttctctg cctggcctgt ttggactttg gggcacagcc 840
 tgtcactgct 850

<210> 29
 <211> 670
 <212> DNA
 <213> Artificial Sequence
 <400> 29

atccacact tctcagagct gccctggac gaccagggtca tcctgctgcg ggcaggctgg 60
 aatgagctgc tcatcgctc cttctccac cgctccatcg ccgtgaagga cgggatccctc 120
 ctggccaccg ggctgcacgt ccaccggaac agcgcccaca gcgcaggggt gggcgccatc 180
 30 tttgacaggg tgctgacgga gcttgtgtcc aagatgcggg acatgcagat ggacaagacg 240
 gagctgggct gcctgcgcgc catcgtcctc ttttaaccctg actccaaggg gctctcgaac 300
 ccggccgagg tggaggcgct gagggagaag gtctatgcgt ccttggaggc ctactgcaag 360
 35 cacaagtacc cagagcagcc gggaagggtc gctaagctct tgctccgcct gccggtcttg 420
 cgctccatcg ggctcaaatg cctggaacat ctcttcttct tcaagctcat cggggacaca 480
 cccattgaca ccttccttat ggagatgctg gaggcgcgc accaaatgac ttaggcctgc 540
 40 gggcccatcc tttgtgcca cccgttcttg ccaccctgcc tggacgccag ctgttcttct 600
 cagcctgagc cctgtccctg cccttctctg cctggcctgt ttggactttg gggcacagcc 660
 tgtcactgct 670

<210> 30
 <211> 672
 <212> DNA
 <213> Artificial Sequence
 <400> 30

gccaacgagg acatgccggt ggagaggatc ctggaggctg agctggccgt ggagcccaag 60
 accgagacct acgtggaggc aaacatgggg ctgaaccca gctcgccgaa cgaccctgtc 120
 55 accaacattt gccaaagcag cgacaaacag cttttcaccc tgggtggagtg ggccaagcgg 180
 atccacact tctcagagct gccctggac gaccagggtca tcctgctgcg ggcaggctgg 240

EP 1 266 015 B1

aatgagctgc tcatcgectc cttctccac cgtccatcg ccgtgaagga cgggacctc 300
 ctggccaccg ggctgcacgt ccaccggaac agcgccaca gcgcaggggt gggcgccatc 360
 5 tttgacaggg tgctgacgga gcttggtgcc aagatgcggg acatgcagat ggacaagacg 420
 gagctgggct gcctgcgcgc catcgtcctc ttttaaccctg actccaaggg gctctcgaac 480
 ccggccgagg tggaggcgct gagggagaag gtctatgcgt ccttgagggc ctactgcaag 540
 10 cacaagtacc cagagcagcc gggaagggtc gctaagctct tgctccgcct gccggctctg 600
 cgctccatcg ggctcaaattg cctggaacat ctcttcttct tcaagctcat cggggacaca 660
 cccattgaca cc 672

<210> 31
 <211> 328
 <212> PRT
 <213> Artificial Sequence
 20 <400> 31

Cys Ala Ile Cys Gly Asp Arg Ser Ser Gly Lys His Tyr Gly Val Tyr
 1 5 10 15
 25 Ser Cys Glu Gly Cys Lys Gly Phe Phe Lys Arg Thr Val Arg Lys Asp
 20 25 30
 Leu Thr Tyr Thr Cys Arg Asp Asn Lys Asp Cys Leu Ile Asp Lys Arg
 35 40 45
 30 Gln Arg Asn Arg Cys Gln Tyr Cys Arg Tyr Gln Lys Cys Leu Ala Met
 50 55 60
 Gly Met Lys Arg Glu Ala Val Gln Glu Glu Arg Gln Arg Gly Lys Asp
 65 70 75 80
 35 Arg Asn Glu Asn Glu Val Glu Ser Thr Ser Ser Ala Asn Glu Asp Met
 85 90 95
 Pro Val Glu Lys Ile Leu Glu Ala Glu Leu Ala Val Glu Pro Lys Thr
 100 105 110
 40 Glu Thr Tyr Val Glu Ala Asn Met Gly Leu Asn Pro Ser Ser Pro Asn
 115 120 125
 Asp Pro Val Thr Asn Ile Cys Gln Ala Ala Asp Lys Gln Leu Phe Thr
 130 135 140
 45 Leu Val Glu Trp Ala Lys Arg Ile Pro His Phe Ser Glu Leu Pro Leu
 145 150 155 160
 Asp Asp Gln Val Ile Leu Leu Arg Ala Gly Trp Asn Glu Leu Leu Ile
 165 170 175
 50 Ala Ser Phe Ser His Arg Ser Ile Ala Val Lys Asp Gly Ile Leu Leu
 180 185 190
 Ala Thr Gly Leu His Val His Arg Asn Ser Ala His Ser Ala Gly Val
 195 200 205
 55 Gly Ala Ile Phe Asp Arg Val Leu Thr Glu Leu Val Ser Lys Met Arg

EP 1 266 015 B1

	210		215		220													
	Asp	Met	Gln	Met	Asp	Lys	Thr	Glu	Leu	Gly	Cys	Leu	Arg	Ala	Ile	Val		
	225					230					235					240		
5	Leu	Phe	Asn	Pro	Asp	Ser	Lys	Gly	Leu	Ser	Asn	Pro	Ala	Glu	Val	Glu		
					245					250					255			
	Ala	Leu	Arg	Glu	Lys	Val	Tyr	Ala	Ser	Leu	Glu	Ala	Tyr	Cys	Lys	His		
				260					265					270				
10	Lys	Tyr	Pro	Glu	Gln	Pro	Gly	Arg	Phe	Ala	Lys	Leu	Leu	Leu	Arg	Leu		
			275					280					285					
	Pro	Ala	Leu	Arg	Ser	Ile	Gly	Leu	Lys	Cys	Leu	Glu	His	Leu	Phe	Phe		
		290					295					300						
15	Phe	Lys	Leu	Ile	Gly	Asp	Thr	Pro	Ile	Asp	Thr	Phe	Leu	Met	Glu	Met		
	305					310					315					320		
	Leu	Glu	Ala	Pro	His	Gln	Ala	Thr										
					325													
20	<210> 32 <211> 262 <212> PRT <213> Artificial Sequence <400> 32																	
25																		
	Lys	Arg	Glu	Ala	Val	Gln	Glu	Glu	Arg	Gln	Arg	Gly	Lys	Asp	Arg	Asn		
	1				5					10				15				
30	Glu	Asn	Glu	Val	Glu	Ser	Thr	Ser	Ser	Ala	Asn	Glu	Asp	Met	Pro	Val		
				20					25					30				
	Glu	Lys	Ile	Leu	Glu	Ala	Glu	Leu	Ala	Val	Glu	Pro	Lys	Thr	Glu	Thr		
			35					40					45					
35	Tyr	Val	Glu	Ala	Asn	Met	Gly	Leu	Asn	Pro	Ser	Ser	Pro	Asn	Asp	Pro		
		50					55					60						
	Val	Thr	Asn	Ile	Cys	Gln	Ala	Ala	Asp	Lys	Gln	Leu	Phe	Thr	Leu	Val		
	65				70						75					80		
40	Glu	Trp	Ala	Lys	Arg	Ile	Pro	His	Phe	Ser	Glu	Leu	Pro	Leu	Asp	Asp		
				85						90					95			
	Gln	Val	Ile	Leu	Leu	Arg	Ala	Gly	Trp	Asn	Glu	Leu	Leu	Ile	Ala	Ser		
				100					105					110				
45	Phe	Ser	His	Arg	Ser	Ile	Ala	Val	Lys	Asp	Gly	Ile	Leu	Leu	Ala	Thr		
			115					120					125					
	Gly	Leu	His	Val	His	Arg	Asn	Ser	Ala	His	Ser	Ala	Gly	Val	Gly	Ala		
		130					135					140						
50	Ile	Phe	Asp	Arg	Val	Leu	Thr	Glu	Leu	Val	Ser	Lys	Met	Arg	Asp	Met		
	145					150					155					160		
	Gln	Met	Asp	Lys	Thr	Glu	Leu	Gly	Cys	Leu	Arg	Ala	Ile	Val	Leu	Phe		
					165					170					175			
55	Asn	Pro	Asp	Ser	Lys	Gly	Leu	Ser	Asn	Pro	Ala	Glu	Val	Glu	Ala	Leu		
				180					185					190				

EP 1 266 015 B1

Arg Glu Lys Val Tyr Ala Ser Leu Glu Ala Tyr Cys Lys His Lys Tyr
195 200 205

Pro Glu Gln Pro Gly Arg Phe Ala Lys Leu Leu Leu Arg Leu Pro Ala
210 215 220

Leu Arg Ser Ile Gly Leu Lys Cys Leu Glu His Leu Phe Phe Phe Lys
225 230 235 240

Leu Ile Gly Asp Thr Pro Ile Asp Thr Phe Leu Met Glu Met Leu Glu
245 250 255

Ala Pro His Gln Ala Thr
260

<210> 33
<211> 237
<212> PRT
<213> Artificial Sequence
<400> 33

Ala Asn Glu Asp Met Pro Val Glu Lys Ile Leu Glu Ala Glu Leu Ala
1 5 10 15

Val Glu Pro Lys Thr Glu Thr Tyr Val Glu Ala Asn Met Gly Leu Asn
20 25 30

Pro Ser Ser Pro Asn Asp Pro Val Thr Asn Ile Cys Gln Ala Ala Asp
35 40 45

Lys Gln Leu Phe Thr Leu Val Glu Trp Ala Lys Arg Ile Pro His Phe
50 55 60

Ser Glu Leu Pro Leu Asp Asp Gln Val Ile Leu Leu Arg Ala Gly Trp
65 70 75 80

Asn Glu Leu Leu Ile Ala Ser Phe Ser His Arg Ser Ile Ala Val Lys
85 90 95

Asp Gly Ile Leu Leu Ala Thr Gly Leu His Val His Arg Asn Ser Ala
100 105 110

His Ser Ala Gly Val Gly Ala Ile Phe Asp Arg Val Leu Thr Glu Leu
115 120 125

Val Ser Lys Met Arg Asp Met Gln Met Asp Lys Thr Glu Leu Gly Cys
130 135 140

Leu Arg Ala Ile Val Leu Phe Asn Pro Asp Ser Lys Gly Leu Ser Asn
145 150 155 160

Pro Ala Glu Val Glu Ala Leu Arg Glu Lys Val Tyr Ala Ser Leu Glu
165 170 175

Ala Tyr Cys Lys His Lys Tyr Pro Glu Gln Pro Gly Arg Phe Ala Lys
180 185 190

Leu Leu Leu Arg Leu Pro Ala Leu Arg Ser Ile Gly Leu Lys Cys Leu
195 200 205

Glu His Leu Phe Phe Phe Lys Leu Ile Gly Asp Thr Pro Ile Asp Thr
210 215 220

EP 1 266 015 B1

Phe Leu Met Glu Met Leu Glu Ala Pro His Gln Ala Thr
225 230 235

5 <210> 34
<211> 177
<212> PRT
<213> Artificial Sequence
10 <400> 34

Ile Pro His Phe Ser Glu Leu Pro Leu Asp Asp Gln Val Ile Leu Leu
1 5 10 15
Arg Ala Gly Trp Asn Glu Leu Leu Ile Ala Ser Phe Ser His Arg Ser
20 25 30
Ile Ala Val Lys Asp Gly Ile Leu Leu Ala Thr Gly Leu His Val His
35 40 45
Arg Asn Ser Ala His Ser Ala Gly Val Gly Ala Ile Phe Asp Arg Val
50 55 60
Leu Thr Glu Leu Val Ser Lys Met Arg Asp Met Gln Met Asp Lys Thr
65 70 75 80
Glu Leu Gly Cys Leu Arg Ala Ile Val Leu Phe Asn Pro Asp Ser Lys
85 90 95
Gly Leu Ser Asn Pro Ala Glu Val Glu Ala Leu Arg Glu Lys Val Tyr
100 105 110
Ala Ser Leu Glu Ala Tyr Cys Lys His Lys Tyr Pro Glu Gln Pro Gly
115 120 125
Arg Phe Ala Lys Leu Leu Leu Arg Leu Pro Ala Leu Arg Ser Ile Gly
130 135 140
Leu Lys Cys Leu Glu His Leu Phe Phe Phe Lys Leu Ile Gly Asp Thr
145 150 155 160
Pro Ile Asp Thr Phe Leu Met Glu Met Leu Glu Ala Pro His Gln Ala
165 170 175
Thr

40 <210> 35
<211> 224
<212> PRT
<213> Artificial Sequence
45 <400> 35

Ala Asn Glu Asp Met Pro Val Glu Lys Ile Leu Glu Ala Glu Leu Ala
1 5 10 15
Val Glu Pro Lys Thr Glu Thr Tyr Val Glu Ala Asn Met Gly Leu Asn
20 25 30
Pro Ser Ser Pro Asn Asp Pro Val Thr Asn Ile Cys Gln Ala Ala Asp
35 40 45
Lys Gln Leu Phe Thr Leu Val Glu Trp Ala Lys Arg Ile Pro His Phe
50 55 60

EP 1 266 015 B1

Ser Glu Leu Pro Leu Asp Asp Gln Val Ile Leu Leu Arg Ala Gly Trp
 65 70 75 80
 Asn Glu Leu Leu Ile Ala Ser Phe Ser His Arg Ser Ile Ala Val Lys
 85 90 95
 Asp Gly Ile Leu Leu Ala Thr Gly Leu His Val His Arg Asn Ser Ala
 100 105 110
 His Ser Ala Gly Val Gly Ala Ile Phe Asp Arg Val Leu Thr Glu Leu
 115 120 125
 Val Ser Lys Met Arg Asp Met Gln Met Asp Lys Thr Glu Leu Gly Cys
 130 135 140
 Leu Arg Ala Ile Val Leu Phe Asn Pro Asp Ser Lys Gly Leu Ser Asn
 145 150 155 160
 Pro Ala Glu Val Glu Ala Leu Arg Glu Lys Val Tyr Ala Ser Leu Glu
 165 170 175
 Ala Tyr Cys Lys His Lys Tyr Pro Glu Gln Pro Gly Arg Phe Ala Lys
 180 185 190
 Leu Leu Leu Arg Leu Pro Ala Leu Arg Ser Ile Gly Leu Lys Cys Leu
 195 200 205
 Glu His Leu Phe Phe Phe Lys Leu Ile Gly Asp Thr Pro Ile Asp Thr
 210 215 220

<210> 36
 <211> 328
 <212> PRT
 <213> Artificial Sequence
 <220>
 <221> misc feature
 <223> Novel Sequence
 <400> 36

Cys Ala Ile Cys Gly Asp Arg Ser Ser Gly Lys His Tyr Gly Val Tyr
 1 5 10 15
 Ser Cys Glu Gly Cys Lys Gly Phe Phe Lys Arg Thr Val Arg Lys Asp
 20 25 30
 Leu Thr Tyr Thr Cys Arg Asp Asn Lys Asp Cys Leu Ile Asp Lys Arg
 35 40 45
 Gln Arg Asn Arg Cys Gln Tyr Cys Arg Tyr Gln Lys Cys Leu Ala Met
 50 55 60
 Gly Met Lys Arg Glu Ala Val Gln Glu Glu Arg Gln Arg Gly Lys Asp
 65 70 75 80
 Arg Asn Glu Asn Glu Val Glu Ser Thr Ser Ser Ala Asn Glu Asp Met
 85 90 95
 Pro Val Glu Arg Ile Leu Glu Ala Glu Leu Ala Val Glu Pro Lys Thr
 100 105 110
 Glu Thr Tyr Val Glu Ala Asn Met Gly Leu Asn Pro Ser Ser Pro Asn
 115 120 125

EP 1 266 015 B1

Asp Pro Val Thr Asn Ile Cys Gln Ala Ala Asp Lys Gln Leu Phe Thr
 130 135 140
 5 Leu Val Glu Trp Ala Lys Arg Ile Pro His Phe Ser Glu Leu Pro Leu
 145 150 155 160
 Asp Asp Gln Val Ile Leu Leu Arg Ala Gly Trp Asn Glu Leu Leu Ile
 165 170 175
 10 Ala Ser Phe Ser His Arg Ser Ile Ala Val Lys Asp Gly Ile Leu Leu
 180 185 190
 Ala Thr Gly Leu His Val His Arg Asn Ser Ala His Ser Ala Gly Val
 195 200 205
 15 Gly Ala Ile Phe Asp Arg Val Leu Thr Glu Leu Val Ser Lys Met Arg
 210 215 220
 Asp Met Gln Met Asp Lys Thr Glu Leu Gly Cys Leu Arg Ala Ile Val
 225 230 235 240
 20 Leu Phe Asn Pro Asp Ser Lys Gly Leu Ser Asn Pro Ala Glu Val Glu
 245 250 255
 Ala Leu Arg Glu Lys Val Tyr Ala Ser Leu Glu Ala Tyr Cys Lys His
 260 265 270
 25 Lys Tyr Pro Glu Gln Pro Gly Arg Phe Ala Lys Leu Leu Leu Arg Leu
 275 280 285
 Pro Ala Leu Arg Ser Ile Gly Leu Lys Cys Leu Glu His Leu Phe Phe
 290 295 300
 30 Phe Lys Leu Ile Gly Asp Thr Pro Ile Asp Thr Phe Leu Met Glu Met
 305 310 315 320
 Leu Glu Ala Pro His Gln Met Thr
 325

35 <210> 37
 <211> 262
 <212> PRT
 <213> Artificial Sequence
 <220>
 <221> misc feature
 40 <223> Novel Sequence
 <400> 37

45 Lys Arg Glu Ala Val Gln Glu Glu Arg Gln Arg Gly Lys Asp Arg Asn
 1 5 10 15
 Glu Asn Glu Val Glu Ser Thr Ser Ser Ala Asn Glu Asp Met Pro Val
 20 25 30
 50 Glu Arg Ile Leu Glu Ala Glu Leu Ala Val Glu Pro Lys Thr Glu Thr
 35 40 45
 Tyr Val Glu Ala Asn Met Gly Leu Asn Pro Ser Ser Pro Asn Asp Pro
 50 55 60
 Val Thr Asn Ile Cys Gln Ala Ala Asp Lys Gln Leu Phe Thr Leu Val

EP 1 266 015 B1

65 70 75 80
 Glu Trp Ala Lys Arg Ile Pro His Phe Ser Glu Leu Pro Leu Asp Asp
 85 90 95
 5 Gln Val Ile Leu Leu Arg Ala Gly Trp Asn Glu Leu Leu Ile Ala Ser
 100 105 110
 Phe Ser His Arg Ser Ile Ala Val Lys Asp Gly Ile Leu Leu Ala Thr
 115 120 125
 10 Gly Leu His Val His Arg Asn Ser Ala His Ser Ala Gly Val Gly Ala
 130 135 140
 Ile Phe Asp Arg Val Leu Thr Glu Leu Val Ser Lys Met Arg Asp Met
 145 150 155 160
 15 Gln Met Asp Lys Thr Glu Leu Gly Cys Leu Arg Ala Ile Val Leu Phe
 165 170 175
 Asn Pro Asp Ser Lys Gly Leu Ser Asn Pro Ala Glu Val Glu Ala Leu
 180 185 190
 20 Arg Glu Lys Val Tyr Ala Ser Leu Glu Ala Tyr Cys Lys His Lys Tyr
 195 200 205
 Pro Glu Gln Pro Gly Arg Phe Ala Lys Leu Leu Leu Arg Leu Pro Ala
 210 215 220
 25 Leu Arg Ser Ile Gly Leu Lys Cys Leu Glu His Leu Phe Phe Phe Lys
 225 230 235 240
 Leu Ile Gly Asp Thr Pro Ile Asp Thr Phe Leu Met Glu Met Leu Glu
 245 250 255
 30 Ala Pro His Gln Met Thr
 260

<210> 38
 <211> 237
 35 <212> PRT
 <213> Artificial Sequence
 <220>
 <221> misc feature
 <223> Novel Sequence
 40 <400> 38

Ala Asn Glu Asp Met Pro Val Glu Arg Ile Leu Glu Ala Glu Leu Ala
 1 5 10 15
 45 Val Glu Pro Lys Thr Glu Thr Tyr Val Glu Ala Asn Met Gly Leu Asn
 20 25 30
 Pro Ser Ser Pro Asn Asp Pro Val Thr Asn Ile Cys Gln Ala Ala Asp
 35 40 45
 50 Lys Gln Leu Phe Thr Leu Val Glu Trp Ala Lys Arg Ile Pro His Phe
 50 55 60
 Ser Glu Leu Pro Leu Asp Asp Gln Val Ile Leu Leu Arg Ala Gly Trp
 65 70 75 80

EP 1 266 015 B1

Asn Glu Leu Leu Ile Ala Ser Phe Ser His Arg Ser Ile Ala Val Lys
85 90 95

5 Asp Gly Ile Leu Leu Ala Thr Gly Leu His Val His Arg Asn Ser Ala
100 105 110

His Ser Ala Gly Val Gly Ala Ile Phe Asp Arg Val Leu Thr Glu Leu
115 120 125

10 Val Ser Lys Met Arg Asp Met Gln Met Asp Lys Thr Glu Leu Gly Cys
130 135 140

Leu Arg Ala Ile Val Leu Phe Asn Pro Asp Ser Lys Gly Leu Ser Asn
145 150 155 160

15 Pro Ala Glu Val Glu Ala Leu Arg Glu Lys Val Tyr Ala Ser Leu Glu
165 170 175

Ala Tyr Cys Lys His Lys Tyr Pro Glu Gln Pro Gly Arg Phe Ala Lys
180 185 190

20 Leu Leu Leu Arg Leu Pro Ala Leu Arg Ser Ile Gly Leu Lys Cys Leu
195 200 205

Glu His Leu Phe Phe Phe Lys Leu Ile Gly Asp Thr Pro Ile Asp Thr
210 215 220

25 Phe Leu Met Glu Met Leu Glu Ala Pro His Gln Met Thr
225 230 235

<210> 39

<211> 177

<212> PRT

<213> Artificial Sequence

<220>

<221> misc_feature

<223> Novel Sequence

<400> 39

Ile Pro His Phe Ser Glu Leu Pro Leu Asp Asp Gln Val Ile Leu Leu
1 5 10 15

40 Arg Ala Gly Trp Asn Glu Leu Leu Ile Ala Ser Phe Ser His Arg Ser
20 25 30

Ile Ala Val Lys Asp Gly Ile Leu Leu Ala Thr Gly Leu His Val His
35 40 45

45 Arg Asn Ser Ala His Ser Ala Gly Val Gly Ala Ile Phe Asp Arg Val
50 55 60

Leu Thr Glu Leu Val Ser Lys Met Arg Asp Met Gln Met Asp Lys Thr
65 70 75 80

50 Glu Leu Gly Cys Leu Arg Ala Ile Val Leu Phe Asn Pro Asp Ser Lys
85 90 95

Gly Leu Ser Asn Pro Ala Glu Val Glu Ala Leu Arg Glu Lys Val Tyr
100 105 110

55 Ala Ser Leu Glu Ala Tyr Cys Lys His Lys Tyr Pro Glu Gln Pro Gly
115 120 125

EP 1 266 015 B1

Arg Phe Ala Lys Leu Leu Leu Arg Leu Pro Ala Leu Arg Ser Ile Gly
 130 135 140
 5 Leu Lys Cys Leu Glu His Leu Phe Phe Phe Lys Leu Ile Gly Asp Thr
 145 150 155 160
 Pro Ile Asp Thr Phe Leu Met Glu Met Leu Glu Ala Pro His Gln Met
 165 170 175
 10 Thr
 <210> 40
 <211> 224
 <212> PRT
 15 <213> Artificial Sequence
 <220>
 <221> misc feature
 <223> Novel Sequence
 <400> 40
 20
 Ala Asn Glu Asp Met Pro Val Glu Arg Ile Leu Glu Ala Glu Leu Ala
 1 5 10 15
 25 Val Glu Pro Lys Thr Glu Thr Tyr Val Glu Ala Asn Met Gly Leu Asn
 20 25 30
 Pro Ser Ser Pro Asn Asp Pro Val Thr Asn Ile Cys Gln Ala Ala Asp
 35 40 45
 30 Lys Gln Leu Phe Thr Leu Val Glu Trp Ala Lys Arg Ile Pro His Phe
 50 55 60
 Ser Glu Leu Pro Leu Asp Asp Gln Val Ile Leu Leu Arg Ala Gly Trp
 65 70 75 80
 35 Asn Glu Leu Leu Ile Ala Ser Phe Ser His Arg Ser Ile Ala Val Lys
 85 90 95
 Asp Gly Ile Leu Leu Ala Thr Gly Leu His Val His Arg Asn Ser Ala
 100 105 110
 40 His Ser Ala Gly Val Gly Ala Ile Phe Asp Arg Val Leu Thr Glu Leu
 115 120 125
 Val Ser Lys Met Arg Asp Met Gln Met Asp Lys Thr Glu Leu Gly Cys
 130 135 140
 45 Leu Arg Ala Ile Val Leu Phe Asn Pro Asp Ser Lys Gly Leu Ser Asn
 145 150 155 160
 Pro Ala Glu Val Glu Ala Leu Arg Glu Lys Val Tyr Ala Ser Leu Glu
 165 170 175
 Ala Tyr Cys Lys His Lys Tyr Pro Glu Gln Pro Gly Arg Phe Ala Lys
 180 185 190
 50 Leu Leu Leu Arg Leu Pro Ala Leu Arg Ser Ile Gly Leu Lys Cys Leu
 195 200 205
 Glu His Leu Phe Phe Phe Lys Leu Ile Gly Asp Thr Pro Ile Asp Thr
 55
 210 215 220

EP 1 266 015 B1

<210> 41
 <211> 441
 <212> DNA
 <213> Artificial Sequence
 <220>
 <221> misc feature
 <223> Novel Sequence
 <400> 41

```

atgaagctac tgtcttctat cgaacaagca tgcgatattt gccgacttaa aaagctcaag      60
tgctccaaag aaaaaccgaa gtgcgccaag tgtctgaaga acaactggga gtgtcgctac      120
tctcccaaaa ccaaaagggt tccgctgact agggcacatc tgacagaagt ggaatcaagg      180
ctagaaagac tggaacagct atttctactg atttttcctc gagaagacct tgacatgatt      240
ttgaaaatgg attctttaca ggatataaaa gcattgttaa caggattatt tgtacaagat      300
aatgtgaata aagatgccgt cacagataga ttggcttcag tggagactga tatgcctcta      360
acattgagac agcatagaat aagtgcgaca tcatcatcgg aagagagtag taacaaagggt      420
caaagacagt tgactgtatc g                                     441
    
```

<210> 42
 <211> 147
 <212> PRT
 <213> Artificial Sequence
 <220>
 <221> misc feature
 <223> Novel Sequence
 <400> 42

```

Met Lys Leu Leu Ser Ser Ile Glu Gln Ala Cys Asp Ile Cys Arg Leu
 1          5          10          15
Lys Lys Leu Lys Cys Ser Lys Glu Lys Pro Lys Cys Ala Lys Cys Leu
          20          25          30
Lys Asn Asn Trp Glu Cys Arg Tyr Ser Pro Lys Thr Lys Arg Ser Pro
          35          40          45
Leu Thr Arg Ala His Leu Thr Glu Val Glu Ser Arg Leu Glu Arg Leu
          50          55          60
Glu Gln Leu Phe Leu Leu Ile Phe Pro Arg Glu Asp Leu Asp Met Ile
65          70          75          80
Leu Lys Met Asp Ser Leu Gln Asp Ile Lys Ala Leu Leu Thr Gly Leu
          85          90          95
Phe Val Gln Asp Asn Val Asn Lys Asp Ala Val Thr Asp Arg Leu Ala
          100          105          110
Ser Val Glu Thr Asp Met Pro Leu Thr Leu Arg Gln His Arg Ile Ser
          115          120          125
    
```

EP 1 266 015 B1

Ala Thr Ser Ser Ser Glu Glu Ser Ser Asn Lys Gly Gln Arg Gln Leu
130 135 140

Thr Val Ser
145

<210> 43
<211> 606
<212> DNA
<213> Artificial Sequence
<220>
<221> misc_feature
<223> Novel Sequence
<400> 43

atgaaagcgt taacggccag gcaacaagag gtgtttgatc tcatccgtga tcacatcagc 60
cagacaggta tgccgccgac gcgtgcggaa atcgcgagc gtttggggtt ccgttcccca 120
aacgcggctg aagaacatct gaaggcgctg gcacgcaaag gcgttattga aattgtttcc 180
ggcgcacac gccgggattcg tctgttcgag gaagaggaag aagggttgcc gctggtaggt 240
cgtgtggctg ccggtgaacc acttctggcg caacagcata ttgaagggtca ttatcaggtc 300
gatccttcct tattcaagcc gaatgctgat ttcctgctgc gcgtcagcgg gatgtcgatg 360
aaagatatcg gcattatgga tgggtgacttg ctggcagtg ataaaactca ggatgtacgt 420
aacggtcagg tcgttgctgc acgtattgat gacgaagtta ccgttaagcg cctgaaaaaa 480
cagggcaata aagtcgaact gttgccagaa aatagcgagt ttaaaccaat tgtcgtagat 540
cttcgtcagc agagcttcac cattgaaggg ctggcggttg gggttattcg caacggcgac 600
tggtctg 606

<210> 44.
<211> 202
<212> PRT
<213> Artificial Sequence
<220>
<221> misc feature
<223> Novel Sequence
<400> 44

Met Lys Ala Leu Thr Ala Arg Gln Gln Glu Val Phe Asp Leu Ile Arg
1 5 10 15
Asp His Ile Ser Gln Thr Gly Met Pro Pro Thr Arg Ala Glu Ile Ala
20 25 30
Gln Arg Leu Gly Phe Arg Ser Pro Asn Ala Ala Glu Glu His Leu Lys
35 40 45
Ala Leu Ala Arg Lys Gly Val Ile Glu Ile Val Ser Gly Ala Ser Arg
50 55 60

EP 1 266 015 B1

Gly Ile Arg Leu Leu Gln Glu Glu Glu Glu Gly Leu Pro Leu Val Gly
 65 70 75 80
 Arg Val Ala Ala Gly Glu Pro Leu Leu Ala Gln Gln His Ile Glu Gly
 5 85 90 95
 His Tyr Gln Val Asp Pro Ser Leu Phe Lys Pro Asn Ala Asp Phe Leu
 100 105 110
 Leu Arg Val Ser Gly Met Ser Met Lys Asp Ile Gly Ile Met Asp Gly
 10 115 120 125
 Asp Leu Leu Ala Val His Lys Thr Gln Asp Val Arg Asn Gly Gln Val
 130 135 140
 Val Val Ala Arg Ile Asp Asp Glu Val Thr Val Lys Arg Leu Lys Lys
 15 145 150 155 160
 Gln Gly Asn Lys Val Glu Leu Leu Pro Glu Asn Ser Glu Phe Lys Pro
 165 170 175
 Ile Val Val Asp Leu Arg Gln Gln Ser Phe Thr Ile Glu Gly Leu Ala
 180 185 190
 Val Gly Val Ile Arg Asn Gly Asp Trp Leu
 195 200

<210> 45
 <211> 271
 <212> DNA
 <213> Artificial Sequence
 <220>
 <221> misc_feature
 <223> Novel Sequence
 <400> 45

atggggccta aaaagaagcg taaagtcgcc ccccgaccg atgtcagcct gggggacgag 60
 35 ctccacttag acggcgagga cgtggcgatg gcgcatgccg acgcgctaga cgatttcgat 120
 ctggacatgt tgggggacgg ggattccccg gggccgggat ttacccccca cgactccgcc 180
 ccctacggcg ctctggatat ggccgacttc gagtttgagc agatgtttac cgatgccctt 240
 40 ggaattgacg agtacggtgg ggaattcccg g 271

<210> 46
 <211> 90
 <212> PRT
 <213> Artificial Sequence
 <220>
 <221> misc feature
 <223> Novel Sequence
 <400> 46

Met Gly Pro Lys Lys Lys Arg Lys Val Ala Pro Pro Thr Asp Val Ser
 1 5 10 15
 55 Leu Gly Asp Glu Leu His Leu Asp Gly Glu Asp Val Ala Met Ala His

EP 1 266 015 B1

20 25 30

Ala Asp Ala Leu Asp Asp Phe Asp Leu Asp Met Leu Gly Asp Gly Asp
35 40 45

5 Ser Pro Gly Pro Gly Phe Thr Pro His Asp Ser Ala Pro Tyr Gly Ala
50 55 60

Leu Asp Met Ala Asp Phe Glu Phe Glu Gln Met Phe Thr Asp Ala Leu
65 70 75 80

10 Gly Ile Asp Glu Tyr Gly Gly Glu Phe Pro
85 90

15 <210> 47
<211> 19
<212> DNA
<213> Artificial Sequence
<220>
<221> misc feature
20 <223> Novel Sequence
<400> 47
ggagtactgt cctccgagc 19
<210> 48
<211> 666
25 <212> DNA
<213> Artificial Sequence
<220>
<221> misc feature
<223> Novel Sequence
30 <400> 48

ggatccccag cttggaattc gacaggttat cagcaacaac acagtcatat ccatttctcaa 60

ttagctctac cacagtgtgt gaaccaatgt atccagcacc acctgtaacc aaaacaattt 120

35 tagaagtact ttcactttgt aactgagctg tcattttatat tgaattttca aaaatttctta 180

cttttttttt ggatggacgc aaagaagttt aataatcata ttacatggca ttaccaccat 240

atacatatcc atatacatat ccatactctaa tcttacctcg actgctgtat ataaaaccag 300

40 tgggttatatg tacagtactg ctgtatataa aaccagtggt tatatgtaca gtacgtcgac 360

tgctgtatat aaaaccagtg gttatatgta cagtactgct gtatataaaa ccagtggtta 420

tatgtacagt acgtcgaggg atgataatgc gattagtttt ttagccttat ttctggggta 480

45 attaatacagc gaagcgatga tttttgatct attaacagat atataaatgc aaaaactgca 540

taaccacttt aactaatact ttcaacattt tcggtttgta ttactttetta ttcaaatgta 600

ataaaagtat caacaaaaaa ttgttaatat acctctatac tttaacgtca aggagaaaaa 660

50 actata 666

55 <210> 49
<211> 1542
<212> DNA
<213> Artificial Sequence
<220>
<221> misc feature

EP 1 266 015 B1

<223> Novel Sequence

<400> 49

5	ctggacctga aacacgaagt ggcttaccga ggggtgctcc caggccaggt gaaggccgaa	60
	ccgggggtcc acaacggcca ggtcaacggc cacgtgaggg actggatggc aggcggcgct	120
	ggtgccaatc cgccgtctcc gggagcgggt gctcaacccc agcctaacaa tgggtattcg	180
10	tcgccactct cctcgggaag ctacgggccc tacagtccaa atgggaaaat aggcggtgag	240
	gaactgtcgc cagcttcaag tataaatggg tgcagtacag atggcgaggc acgacgtcag	300
	aagaagggcc ctgcgccccg tcagcaagag gaactgtgtc tggtatgcgg ggacagagcc	360
15	tccggatacc actacaatgc gctcacgtgt gaaggggtga aagggttctt cagacggagt	420
	gttaccaaaa atgcggttta tttttgtaaa ttcgggtcacg cttgcgaaat ggacatgtac	480
	atgcgacgga aatgccagga gtgcgcctg aagaagtgtc tagctgtagg catgaggcct	540
20	gagtgcgtag tacccgagac tcagtgcgcc atgaagcggg aagagaagaa agcacagaag	600
	gagaaggaca aactgcctgt cagcacgacg acggtggacg accacatgcc gccattatg	660
	cagtgtgaac ctccacctcc tgaagcagca aggattcacg aagtgggtccc aaggtttctc	720
	tccgacaagc tgttggagac aaaccggcag aaaaacatcc ccagttgac agccaaccag	780
25	cagttcctta tcgccaggct catctggtac caggacgggt acgagcagcc ttctgatgaa	840
	gatttgaaga ggattacgca gacgtggcag caagcggacg atgaaaacga agagtctgac	900
	actoccttcc gccagatcac agagatgact atcctcacgg tccaacttat cgtggagttc	960
30	gcgaagggat tgccagggtt cgccaagatc tcgcagcctg atcaaattac gctgcttaag	1020
	gcttgctcaa gtgaggtaat gatgtccga gtcgcgcgac gatacgatgc ggcctcagac	1080
	agtgttctgt tcgcgaacaa ccaagcgtac actcgcgaca actaccgcaa ggctggcatg	1140
35	gcctacgtca tcgaggatct actgcacttc tgccgggtgca tgtactctat ggcgttggac	1200
	aacatccatt acgcgtgct caccggtgtc gtcatctttt ctgaccggcc agggttggag	1260
	cagccgcaac tgggtggaaga aatccagcgg tactacctga atacgctccg catctatate	1320
40	ctgaaccagc tgagcgggtc ggcgcgttcg tccgtcatat acggcaagat cctctcaatc	1380
	ctctctgagc tacgcacgct cggcatgcaa aactccaaca tgtgcatctc cctcaagctc	1440
	aagaacagaa agctgccgcc tttcctcgag gagatctggg atgtggcgga catgtcgcac	1500
45	acccaaccgc cgcctatcct cgagtcccc acgaatctct ag	1542

<210> 50

<211> 513

<212> PRT

50 <213> Artificial Sequence

<220>

<221> misc feature

<223> Novel Sequence

<400> 50

55

EP 1 266 015 B1

Leu Asp Leu Lys His Glu Val Ala Tyr Arg Gly Val Leu Pro Gly Gln
 1 5 10 15
 Val Lys Ala Glu Pro Gly Val His Asn Gly Gln Val Asn Gly His Val
 5 20 25 30
 Arg Asp Trp Met Ala Gly Gly Ala Gly Ala Asn Ser Pro Ser Pro Gly
 35 40 45
 Ala Val Ala Gln Pro Gln Pro Asn Asn Gly Tyr Ser Ser Pro Leu Ser
 10 50 55 60
 Ser Gly Ser Tyr Gly Pro Tyr Ser Pro Asn Gly Lys Ile Gly Arg Glu
 65 70 75 80
 Glu Leu Ser Pro Ala Ser Ser Ile Asn Gly Cys Ser Thr Asp Gly Glu
 15 85 90 95
 Ala Arg Arg Gln Lys Lys Gly Pro Ala Pro Arg Gln Gln Glu Glu Leu
 100 105 110
 Cys Leu Val Cys Gly Asp Arg Ala Ser Gly Tyr His Tyr Asn Ala Leu
 20 115 120 125
 Thr Cys Glu Gly Cys Lys Gly Phe Phe Arg Arg Ser Val Thr Lys Asn
 130 135 140
 Ala Val Tyr Ile Cys Lys Phe Gly His Ala Cys Glu Met Asp Met Tyr
 25 145 150 155 160
 Met Arg Arg Lys Cys Gln Glu Cys Arg Leu Lys Lys Cys Leu Ala Val
 165 170 175
 Gly Met Arg Pro Glu Cys Val Val Pro Glu Thr Gln Cys Ala Met Lys
 180 185 190
 Arg Lys Glu Lys Lys Ala Gln Lys Glu Lys Asp Lys Leu Pro Val Ser
 30 195 200 205
 Thr Thr Thr Val Asp Asp His Met Pro Pro Ile Met Gln Cys Glu Pro
 210 215 220
 Pro Pro Pro Glu Ala Ala Arg Ile His Glu Val Val Pro Arg Phe Leu
 35 225 230 235 240
 Ser Asp Lys Leu Leu Glu Thr Asn Arg Gln Lys Asn Ile Pro Gln Leu
 245 250 255
 Thr Ala Asn Gln Gln Phe Leu Ile Ala Arg Leu Ile Trp Tyr Gln Asp
 40 260 265 270
 Gly Tyr Glu Gln Pro Ser Asp Glu Asp Leu Lys Arg Ile Thr Gln Thr
 275 280 285
 Trp Gln Gln Ala Asp Asp Glu Asn Glu Glu Ser Asp Thr Pro Phe Arg
 45

50

55

EP 1 266 015 B1

	290	295	300	
	Gln Ile Thr Glu Met Thr	Ile Leu Thr Val	Gln Leu Ile Val Glu Phe	
	305	310	315	320
5	Ala Lys Gly Leu Pro Gly Phe Ala Lys	Ile	Ser Gln Pro Asp Gln Ile	
		325	330	335
	Thr Leu Leu Lys Ala Cys Ser Ser Glu Val Met Met Leu Arg Val Ala			
		340	345	350
10	Arg Arg Tyr Asp Ala Ala Ser Asp Ser Val Leu Phe Ala Asn Asn Gln			
		355	360	365
	Ala Tyr Thr Arg Asp Asn Tyr Arg Lys Ala Gly Met Ala Tyr Val Ile			
		370	375	380
15	Glu Asp Leu Leu His Phe Cys Arg Cys Met Tyr Ser Met Ala Leu Asp			
		385	390	395
	Asn Ile His Tyr Ala Leu Leu Thr Ala Val Val Ile Phe Ser Asp Arg			
		405	410	415
20	Pro Gly Leu Glu Gln Pro Gln Leu Val Glu Glu Ile Gln Arg Tyr Tyr			
		420	425	430
	Leu Asn Thr Leu Arg Ile Tyr Ile Leu Asn Gln Leu Ser Gly Ser Ala			
		435	440	445
25	Arg Ser Ser Val Ile Tyr Gly Lys Ile Leu Ser Ile Leu Ser Glu Leu			
		450	455	460
	Arg Thr Leu Gly Met Gln Asn Ser Asn Met Cys Ile Ser Leu Lys Leu			
		465	470	475
30	Lys Asn Arg Lys Leu Pro Pro Phe Leu Glu Glu Ile Trp Asp Val Ala			
		485	490	495
	Asp Met Ser His Thr Gln Pro Pro Pro Ile Leu Glu Ser Pro Thr Asn			
		500	505	510
35	Leu			

<210> 51
 <211> 4375
 <212> DNA
 <213> Artificial Sequence
 <220>
 <221> misc_feature
 <223> Novel Sequence
 <400> 51

	tgtaattttg atgggcgccg tgatgcaccg tgtgccatat tgccatccag tcgaatagaa	60
	aaaaaaaaaa aaaaaaaaaat atcagttggtt ttgtccctcg ctgcctttcg agtgtattcg	120
50	gaatattaga cgtcataatt cagcagtggtc ttttaaattt atatagcgat tagcggggcc	180
	gtttgttgga cgtgcgcttg cgttttagtgg agtgcaggga tagtgaggcg agtatggtag	240
	ttcgtggtca tgtcaagtgt ggcgaagaaa gacaagccga cgatgtcggg gacggcgctg	300

EP 1 266 015 B1

	atcaactggg	cgcgcccgcc	gccgccaggc	ccgccgcagc	cgcagtcagc	gtcgccctgcg	360
	ccggcagcca	tgctgcagca	gctcccgcag	cagtcaatgc	agtcgttaaa	ccacatccca	420
5	actgtcgatt	gctcgctcga	tatgcagtgg	cttaatttag	aacctggatt	catgtcgcct	480
	atgtcacctc	ctgagatgaa	accagacacc	gccatgcttg	atgggctacg	agacgacgcc	540
	acttcgccgc	ctaacttcaa	gaactaccgc	cctaataacc	ccctgagtgg	ctccaaacac	600
10	ctatgctcta	tatgcggcga	cagggcgctc	gggaagcact	atgggggtga	cagttgcgaa	660
	ggatgcaagg	gtttcttcaa	gcggaccgtc	cggaaggacc	tgctgtacgc	ttgccgggag	720
	gagcgggaact	gcatcataga	caagcgacaa	aggaaccgat	gccagtactg	ccgctatcaa	780
15	aagtgtttgg	cttgccgtat	gaagcgagag	gcgggtgcaag	aggagcgcca	gaggaatgct	840
	cgcgccgcgg	aggatgcgca	cccagtagtc	tccgtgcagg	taagcgatga	gctgtcaatc	900
	gagcgcctaa	cggagatgga	gtctttgggt	gcagatccca	gcgaggagtt	ccagttcctc	960
20	cgcggtgggc	ctgacagcaa	cgtgcctcca	cgttaccgcg	cgcccgtctc	ctccctctgc	1020
	caaataggca	acaagcaaat	agcggcggtg	gtgggtatgg	cgcgcgacat	ccctcatttc	1080
	gggcagctgg	agctggacga	tcaagtggta	ctcatcaagg	cctcctggaa	tgagctgcta	1140
25	ctcttcgccca	tcgcctggcg	ctctatggag	tattttggaag	atgagagggga	gaacggggac	1200
	ggaacgcgga	gcaccactca	gccacaactg	atgtgtctca	tgccctggcat	gacgttgcac	1260
	cgcaactcgg	cgcgagcagg	gggcgtgggc	gccatcttcg	accgcgtgct	gtccgagctc	1320
30	agtctgaaga	tgcgacacct	gcgcattggc	caggccgagt	acgtcgcgct	caaagccatc	1380
	gtgctgctca	accctgatgt	gaaaggactg	aagaatcggc	aagaagttga	cgttttgcga	1440
	gaaaaaatgt	tctcttgcc	ggacgactac	tgccggcggt	cgcgaaagcaa	cgaggaaggc	1500
	cggtttgcgt	ccttgctgct	gcggctgcca	gctctccgct	ccatctcgct	caagagcttc	1560
35	gaacacctct	actttctcca	cctcgtggcc	gaaggctcca	tcagcggata	catagcagag	1620
	gcgctccgaa	accacgcgcc	tccgatcgac	gtcaatgccca	tgatgtaaag	tgcgatacac	1680
	gccctgccga	tgtgagaaga	actatggcta	atagaagcga	aactgaatac	atctaggggtg	1740
40	ggacttaact	tgggactatc	attaaagtat	cacgcaaatt	atgcgtagtc	agaaagtcgc	1800
	gtcgatcaaa	cttttttata	aacgaattga	gtttctaacg	actgcaacac	agcggagttt	1860
	tgcttctgat	agttttttatt	ctaattggta	agatgcttta	cacgggcatt	attgacattc	1920
45	aagtgtaaagt	ggaagttgac	aaccttgaca	tttatatcac	gtttgttaatt	ggttaaataa	1980
	attaattaat	cacaagtaag	actaacatca	acgtcacgat	actaacgccca	tttagtgata	2040
	tttttcatgt	caagaaactc	attgttttga	taaaatattt	ttctaattac	tccagtgaac	2100
50	tcatccaaat	gtgaccaggt	ttcccgacga	gttgcccggt	taaaatcatc	tttagggaca	2160
	tatccccgcg	tatctcatga	aattccaagg	atcagtaggg	gccaattccc	ccgatgtggt	2220
	gggaggcaga	attttcgata	atctacgact	attgttagcc	tacgaattag	ttgaattttt	2280

55

EP 1 266 015 B1

tgaaattatt tttattaagt cgccactttc caaacacatc agcaggggat atgtgcaatt 2340
 ttgtaacgat aactctattc atttctgata tttatcgaaa ttttatctta cataacatgc 2400
 5 tggctgggcc aggtgtttgg tagttacata tgtatctacg gtttgtttta aattatagct 2460
 tttttattgt aatctgtata aaattgagtt atcttacttc acactacgat cgagtaaacc 2520
 catcgtcagc tacgaaaaac taatcgtata aggcgtaaga gtaaataact aattgacaac 2580
 10 cagcaacgag gaccacctca gtccctcgtc ttacattgtg ccgtagctta atatgatgga 2640
 agctgtcgtc gttacgacat tagataaagt gcatgaatac caaaaatgta ccatcccgtg 2700
 ctgatctctc atgctctcgc tgcgtgggac ccgtgtcgag tgtcgtgaagg actgactaat 2760
 15 attttagact aggcgtctat gcttcagtaa ttccttatac atattataag tcatccaaat 2820
 aacgagtaag gcggcatgtt gagatcagca ttcgagagat caaagagccc ctaacgtgac 2880
 tgagaagtag agacaataca ctgattttct gagatgaacg caaccgagat tgacactaaa 2940
 20 aatctattta tggatttcaa aatggcgatg cttgattgtc tgcggcggtg atagactgaa 3000
 atgggtttgc ttaacactgg atattgtttt tattagttaa tagtcttaca ttgcaagttg 3060
 gtaattcggg gctaataatcg accggtttgt taactatcta acggttccca gtgtcaggca 3120
 25 cacatctttc ccaagcagac aacgcaagag tgtacaaaat gtacatgtta caaaaataagg 3180
 aacattcgtc ggataagtgt aacagttgat aggtaaagaa aatggggccg cctctttatt 3240
 attacgtagc cgtaaaatta ttaacgtatt tagtttagat gttcagctaa ttaggataat 3300
 30 tctatttgtc gagtacctag atgtccatag tgaattaata taataattag actgttacgc 3360
 gtaggtaatt ataaagttta ccaaactctc cttcaaagca aaaactttgt acacttccgt 3420
 actgagacgt cgtagcttat tctgattcac gaaatatttg gatcacattg ttacaaggcg 3480
 35 accgtcacgt agtatatgat tatttacaaa tgacacgtat gtatcaatgc tataagtgtt 3540
 ttcgttacat atgtcgtgac tttaacgtgc atttcgatgt gcagattaaa aatagcaaga 3600
 aatcttgaaa ttgtttttaga aaatatttga tttccttatt gaaagtattt tttaaatgta 3660
 40 aatatttcgt aatcataata attatgtatt gtgtagttat ttcacctta cggttgggat 3720
 attatttaat ggtggcctac gaaagtgtt ataaccatcc gcgtcctcaa aaaggccagt 3780
 ttatttttgt acctcataca tactaattac gtaagtaata tcaggcgaat ggttgactaa 3840
 caactaacca gtattaaaaa ttaaaagact tcgtcctaata aaaatgtaat atctatgtat 3900
 45 aaaaatgaaa aatctggcgt ataataggta aaattaaact agattgttaa tgaatgtgat 3960
 gtctcataaa cgttttagttt ttaatgagaa acatgttttag tcgcctacta taagacgaga 4020
 cggcaagctc accgagttaa ctctgtaaaca ggaatgttga aaaagatgac acaatttata 4080
 50 tttggtattg aaattatgac taaccatgcg ctctatcgtt tggtatggat gcatagtatt 4140
 gctgttgaaa ataatggaat taggtaatta ctgcattaat gttgaaaact tgatattatt 4200

55

EP 1 266 015 B1

ctatggttgg gtatgaattc tatgttgga gttgtgcagc ggttgtaaag atgatttata 4260
atgatgttca ctaaataatct gactaaatgt aagttatttt tttttgtata gacatagctt 4320
5 taagatgaag gtgattaaac tttatcctta tcacaataaa aaaaaaaaaa aaaaa 4375

<210> 52
<211> 472
10 <212> PRT
<213> Artificial Sequence
<220>
<221> misc_feature
<223> Novel Sequence
15 <400> 52

Met Ser Ser Val Ala Lys Lys Asp Lys Pro Thr Met Ser Val Thr Ala
1 5 10 15
20 Leu Ile Asn Trp Ala Arg Pro Ala Pro Pro Gly Pro Pro Gln Pro Gln
20 20 25 30
Ser Ala Ser Pro Ala Pro Ala Ala Met Leu Gln Gln Leu Pro Thr Gln
35 35 40 45
25 Ser Met Gln Ser Leu Asn His Ile Pro Thr Val Asp Cys Ser Leu Asp
50 55 60
Met Gln Trp Leu Asn Leu Glu Pro Gly Phe Met Ser Pro Met Ser Pro
65 70 75 80
30 Pro Glu Met Lys Pro Asp Thr Ala Met Leu Asp Gly Leu Arg Asp Asp
85 90 95
Ala Thr Ser Pro Pro Asn Phe Lys Asn Tyr Pro Pro Asn His Pro Leu
100 105 110
35 Ser Gly Ser Lys His Leu Cys Ser Ile Cys Gly Asp Arg Ala Ser Gly
115 120 125
Lys His Tyr Gly Val Tyr Ser Cys Glu Gly Cys Lys Gly Phe Phe Lys
130 135 140
40 Arg Thr Val Arg Lys Asp Leu Ser Tyr Ala Cys Arg Glu Glu Arg Asn
145 150 155 160
Cys Ile Ile Asp Lys Arg Gln Arg Asn Arg Cys Gln Tyr Cys Arg Tyr
165 170 175
45 Gln Lys Cys Leu Ala Cys Gly Met Lys Arg Glu Ala Val Gln Glu Glu
180 185 190
Arg Gln Arg Asn Ala Arg Gly Ala Glu Asp Ala His Pro Ser Ser Ser
195 200 205
50 Val Gln Val Ser Asp Glu Leu Ser Ile Glu Arg Leu Thr Glu Met Glu
210 215 220
Ser Leu Val Ala Asp Pro Ser Glu Glu Phe Gln Phe Leu Arg Val Gly
225 230 235 240
55 Pro Asp Ser Asn Val Pro Pro Arg Tyr Arg Ala Pro Val Ser Ser Leu

EP 1 266 015 B1

	245	250	255
	Cys Gln Ile Gly Asn Lys Gln Ile Ala Ala Leu Val Val Trp Ala Arg	260	265 270
5	Asp Ile Pro His Phe Gly Gln Leu Glu Leu Asp Asp Gln Val Val Leu	275	280 285
	Ile Lys Ala Ser Trp Asn Glu Leu Leu Leu Phe Ala Ile Ala Trp Arg	290	295 300
10	Ser Met Glu Tyr Leu Glu Asp Glu Arg Glu Asn Gly Asp Gly Thr Arg	305	310 315 320
	Ser Thr Thr Gln Pro Gln Leu Met Cys Leu Met Pro Gly Met Thr Leu	325	330 335
15	His Arg Asn Ser Ala Gln Gln Ala Gly Val Gly Ala Ile Phe Asp Arg	340	345 350
	Val Leu Ser Glu Leu Ser Leu Lys Met Arg Thr Leu Arg Met Asp Gln	355	360 365
20	Ala Glu Tyr Val Ala Leu Lys Ala Ile Val Leu Leu Asn Pro Asp Val	370	375 380
	Lys Gly Leu Lys Asn Arg Gln Glu Val Asp Val Leu Arg Glu Lys Met	385	390 395 400
25	Phe Ser Cys Leu Asp Asp Tyr Cys Arg Arg Ser Arg Ser Asn Glu Glu	405	410 415
	Gly Arg Phe Ala Ser Leu Leu Leu Arg Leu Pro Ala Leu Arg Ser Ile	420	425 430
30	Ser Leu Lys Ser Phe Glu His Leu Tyr Phe Phe His Leu Val Ala Glu	435	440 445
	Gly Ser Ile Ser Gly Tyr Ile Arg Glu Ala Leu Arg Asn His Ala Pro	450	455 460
35	Pro Ile Asp Val Asn Ala Met Met	465	470

<210> 53
 <211> 1404
 <212> DNA
 <213> Artificial Sequence
 <220>
 <221> misc feature
 <223> Novel Sequence
 <400> 53

	atggacacca aacatttcct gccgctcgac ttctctaccc aggtgaactc ttcgtccctc	60
50	aactctccaa cgggtcgagg ctccatggct gtcccctcgc tgcaccctc cttgggtccg	120
	ggaatcggct ctccactggg ctgcctggg cagctgcact ctctatcag caccctgagc	180
	tcccccatca atggcatggg tccgcccttc tctgtcatca gtcccccat gggcccgac	240
55	tccatgtcgg taccaccac accacattg ggcttcggga ctggtagccc ccagctcaat	300

EP 1 266 015 B1

	tcacccatga	accctgtgag	cgcactgag	gatatcaagc	cgccactagg	cctcaatggc	360
	gtcctcaagg	ttcctgcccc	tccctcagga	aatatggcct	ccttcaccaa	gcacatctgt	420
5	gctatctgtg	gggaccgctc	ctcaggcaaa	cactatgggg	tatacagttg	tgaggggctgc	480
	aagggcttct	tcaagaggac	agtacgcaa	gacctgacct	acacctgccg	agacaacaag	540
	gactgcctga	tcgacaagag	acagcggaac	cggtgtcagt	actgccgcta	ccagaagtgc	600
10	ctggccatgg	gcatgaagcg	ggaagctgtg	caggaggagc	ggcagcgggg	caaggaccgg	660
	aatgagaacg	aggtggagtc	caccagcagt	gccaacgagg	acatgcctgt	agagaagatt	720
	ctggaagccg	agcttgctgt	cgagcccaa	actgagacat	acgtggaggc	aaacatgggg	780
15	ctgaacccca	gctcaccaaa	tgacctgtt	accaacatct	gtcaagcagc	agacaagcag	840
	ctcttcactc	ttgtggagtg	ggccaagagg	atcccaact	tttctgagct	gcccttagac	900
	gaccaggtca	tcctgctacg	ggcaggctgg	aacgagctgc	tgatcgccctc	cttctccac	960
20	cgctccatag	ctgtgaaaga	tgggattctc	ctggccaccg	gcctgcacgt	acaccggaac	1020
	agcgctcaca	gtgctggggg	ggcgccatc	tttgacaggg	tgctaacaga	gctggtgtct	1080
	aagatgcgtg	acatgcagat	ggacaagacg	gagctgggct	gcctgcgagc	cattgtcctg	1140
25	ttcaaccctg	actctaaggg	gctctcaaac	cctgctgagg	tggaggcggt	gagggagaag	1200
	gtgtatgcgt	cactagaagc	gtactgcaa	cacaagtacc	ctgagcagcc	gggcaggttt	1260
	gccaaactgc	tgctccgcct	gcctgcactg	cgttccatcg	ggctcaagtg	cctggagcac	1320
30	ctgttcttct	tcaagctcat	cggggacacg	cccatcgaca	ccttcctcat	ggagatgctg	1380
	gaggcaccac	atcaagccac	ctag				1404

```
<210> 54
<211> 467
<212> PRT
<213> Artificial Sequence
<220>
<221> misc feature
<223> Novel Sequence
<400> 54
```

	Met	Asp	Thr	Lys	His	Phe	Leu	Pro	Leu	Asp	Phe	Ser	Thr	Gln	Val	Asn
	1				5					10					15	
45	Ser	Ser	Ser	Leu	Asn	Ser	Pro	Thr	Gly	Arg	Gly	Ser	Met	Ala	Val	Pro
				20					25					30		
	Ser	Leu	His	Pro	Ser	Leu	Gly	Pro	Gly	Ile	Gly	Ser	Pro	Leu	Gly	Ser
			35					40					45			
50	Pro	Gly	Gln	Leu	His	Ser	Pro	Ile	Ser	Thr	Leu	Ser	Ser	Pro	Ile	Asn
		50					55					60				
	Gly	Met	Gly	Pro	Pro	Phe	Ser	Val	Ile	Ser	Ser	Pro	Met	Gly	Pro	His
	65					70					75					80

EP 1 266 015 B1

Ser Met Ser Val Pro Thr Thr Pro Thr Leu Gly Phe Gly Thr Gly Ser
 85 90 95
 5 Pro Gln Leu Asn Ser Pro Met Asn Pro Val Ser Ser Thr Glu Asp Ile
 100 105 110
 Lys Pro Pro Leu Gly Leu Asn Gly Val Leu Lys Val Pro Ala His Pro
 115 120 125
 10 Ser Gly Asn Met Ala Ser Phe Thr Lys His Ile Cys Ala Ile Cys Gly
 130 135 140
 Asp Arg Ser Ser Gly Lys His Tyr Gly Val Tyr Ser Cys Glu Gly Cys
 145 150 155 160
 15 Lys Gly Phe Phe Lys Arg Thr Val Arg Lys Asp Leu Thr Tyr Thr Cys
 165 170 175
 Arg Asp Asn Lys Asp Cys Leu Ile Asp Lys Arg Gln Arg Asn Arg Cys
 180 185 190
 20 Gln Tyr Cys Arg Tyr Gln Lys Cys Leu Ala Met Gly Met Lys Arg Glu
 195 200 205
 Ala Val Gln Glu Glu Arg Gln Arg Gly Lys Asp Arg Asn Glu Asn Glu
 210 215 220
 25 Val Glu Ser Thr Ser Ser Ala Asn Glu Asp Met Pro Val Glu Lys Ile
 225 230 235 240
 Leu Glu Ala Glu Leu Ala Val Glu Pro Lys Thr Glu Thr Tyr Val Glu
 245 250 255
 30 Ala Asn Met Gly Leu Asn Pro Ser Ser Pro Asn Asp Pro Val Thr Asn
 260 265 270
 Ile Cys Gln Ala Ala Asp Lys Gln Leu Phe Thr Leu Val Glu Trp Ala
 275 280 285
 35 Lys Arg Ile Pro His Phe Ser Glu Leu Pro Leu Asp Asp Gln Val Ile
 290 295 300
 Leu Leu Arg Ala Gly Trp Asn Glu Leu Leu Ile Ala Ser Phe Ser His
 305 310 315 320
 Arg Ser Ile Ala Val Lys Asp Gly Ile Leu Leu Ala Thr Gly Leu His
 325 330 335
 40 Val His Arg Asn Ser Ala His Ser Ala Gly Val Gly Ala Ile Phe Asp
 340 345 350
 Arg Val Leu Thr Glu Leu Val Ser Lys Met Arg Asp Met Gln Met Asp
 355 360 365
 45 Lys Thr Glu Leu Gly Cys Leu Arg Ala Ile Val Leu Phe Asn Pro Asp
 370 375 380
 Ser Lys Gly Leu Ser Asn Pro Ala Glu Val Glu Ala Leu Arg Glu Lys
 385 390 395 400
 50 Val Tyr Ala Ser Leu Glu Ala Tyr Cys Lys His Lys Tyr Pro Glu Gln
 405 410 415
 Pro Gly Arg Phe Ala Lys Leu Leu Leu Arg Leu Pro Ala Leu Arg Ser

EP 1 266 015 B1

		420		425		430	
	Ile Gly Leu Lys Cys Leu Glu His Leu Phe Phe Phe Lys Leu Ile Gly						
		435		440		445	
5	Asp Thr Pro Ile Asp Thr Phe Leu Met Glu Met Leu Glu Ala Pro His						
		450		455		460	
	Gln Ala Thr						
10	465						
	<210> 55						
	<211> 309						
	<212> DNA						
15	<213> Artificial Sequence						
	<220>						
	<221> misc_feature						
	<223> Novel Sequence						
	<400> 55						
20							
	gggtgtggaaa gtccccaggc tccccagcag gcagaagtat gcaaagcatg catctcaatt					60	
	agtcagcaac caggtgtgga aagtccccag gctccccagc aggcagaagt atgcaaagca					120	
25	tgcattctcaa ttagtcagca accatagtcc cgcccctaac tccgcccata ccgcccctaa					180	
	ctccgcccag ttccgcccata tctccgcccc atggctgact aatttttttt atttatgcag					240	
	aggccgaggc cgcctcgggc tctgagctat tccagaagta gtgaggaggc ttttttggag					300	
30	gcctaggct					309	
	<210> 56						
	<211> 24						
	<212> DNA						
35	<213> Artificial Sequence						
	<220>						
	<221> misc feature						
	<223> Novel-Sequence						
	<400> 56						
40	tatataatgg atccccgggt accg 24						
	<210> 57						
	<211> 1653						
	<212> DNA						
	<213> Artificial Sequence						
45	<220>						
	<221> misc feature						
	<223> Novel Sequence						
	<400> 57						
50							
	atggaagacg ccaaaaacat aaagaaaggc ccggcgccat tctatcctct agaggatgga					60	
	accgctggag agcaactgca taaggctatg aagagatacg ccctggttcc tggaacaatt					120	
55	gcttttacag atgcacatat cgagggtgaac atcacgtaag cggaataact cgaaatgtcc					180	

EP 1 266 015 B1

gttcggttgg cagaagctat gaaacgatat gggctgaata caaatcacag aatcgtcgtta 240
 tgcagtgaaa actctcttca attctttatg ccggtgttgg gcgcgttatt tatcggagtt 300
 5 gcagttgcgc ccgcgaacga catttataat gaacgtgaat tgctcaacag tatgaacatt 360
 tcgcagccta ccgtagtggt tgtttccaaa aaggggttgc aaaaaatttt gaacgtgcaa 420
 aaaaaattac caataatcca gaaaattatt atcatggatt ctaaaacgga ttaccaggga 480
 10 tttcagtcga tgtacacgtt cgtcacatct catctacctc ccggttttaa tgaatacgat 540
 tttgtaccag agtcctttga tcgtgacaaa acaattgcac tgataatgaa ttcctctgga 600
 tctactgggt tacctaagggt tgtggccctt ccgcatagaa ctgcctgcgt cagattctcg 660
 15 catgccagag atcctatttt tggcaatcaa atcattccgg atactgcat ttttaagtgtt 720
 gttccattcc atcacggttt tggaaatgttt actacactcg gatatttgat atgtggattt 780
 cgagtcgtct taatgtatag atttgaagaa gagctgtttt tacgatccct tcaggattac 840
 20 aaaattcaaa gtgcgttgct agtaccaacc ctattttcat tcttcgcaa aagcactctg 900
 attgacaaat acgatttatc taatttacac gaaattgctt ctgggggagc acctctttcg 960
 aaagaagtcg gggaagcgggt tgcaaaacgc ttccatcttc cagggatacg acaaggatat 1020
 25 gggctcactg agactacatc agctattctg attacacccg agggggatga taaaccgggc 1080
 gcggtcggta aagttgttcc attttttgaa gcgaagggtt tggtatctgga taccgggaaa 1140
 acgctgggag ttaatcagag aggcgaatta tgtgtcagag gacctatgat tatgtccggt 1200
 30 tatgtaaaca atccggaagc gaccaacgcc ttgattgaca aggatggatg gctacattct 1260
 ggagacatag cttactggga cgaagacgaa cacttcttca tagttgaccg cttgaagtct 1320
 ttaattaaat acaaaggata tcaggtggcc cccgctgaat tggaatcgat attgttacia 1380
 35 caccocaaca tcttcgacgc gggcgtggca ggtcttcccg acgatgacgc cgggtgaactt 1440
 ccgcgcgccc ttgttgtttt ggagcacgga aagacgatga cggaaaaaga gatcgtggat 1500
 tacgtcgcca gtcaagtaac aaccgcgaaa aagttgcgag gaggagttgt gtttgtggac 1560
 40 gaagtaccga aaggtcttac cggaaaactc gacgcaagaa aaatcagaga gatcctcata 1620
 aaggccaaga agggcggaag gtccaaattg taa 1653

45 <210> 58
 <211> 867
 <212> DNA
 <213> Artificial Sequence
 <220>
 <221> misc feature
 50 <223> Novel Sequence
 <400> 58

aagcgagagg cgggtgcaaga ggagcgccag aggaatgctc gcggcgcgga ggatgcgcac 60
 55

EP 1 266 015 B1

ccgagtagct cgggtgcaggt aagcgatgag ctgtcaatcg agcgcctaac ggagatggag 120
tctttggtgg cagatcccag cgaggagttc cagttcctcc gcgtggggcc tgacagcaac 180
5 gtgcctccac gttaccgcgc gcccgctctcc tccctctgcc aaataggcaa caagcaaata 240
gcggcggttg tggatatggc gcgcgacatc cctcatttcg ggcagctgga gctggacgat 300
caagtggtag tcatcaaggc ctcttggaaat gagctgctac tcttcgccat cgcttggcgc 360
10 tctatggagt atttggaaga tgagagggag aacggggacg gaacgcggag caccactcag 420
ccacaactga tgtgtctcat gcttggcatg acgttgaccc gcaactcggc gcagcaggcg 480
ggcgtggggc ccatcttcga ccgcgtgctg tccgagctca gtctgaagat gcgcaccttg 540
15 cgcattggacc aggccgagta cgtcgcgctc aaagccatcg tgctgctcaa ccctgatgtg 600
aaaggactga agaatcggca agaagttgac gttttgcgag aaaaaatgtt ctcttgctg 660
gacgactact gccggcggtc gcgaagcaac gaggaaggcc gggttgctgc cttgctgctg 720
20 cggttgccag ctctccgctc catctcgtc aagagdttcg aacacctcta cttcttccac 780
ctcgtggccg aaggctccat cagcggatac atacgagagg cgctccgaaa ccacgcgcct 840
ccgatcgacg tcaatgccat gatgtaa 867

25
<210> 59
<211> 225
<212> DNA
<213> Artificial Sequence
30 <220>
<221> misc_feature
<223> Novel Sequence
<400> 59

35 tcgacattgg acaagtgcac tgaacccttg tctctcgaga gacaaggggg ttcaatgcac 60
ttgtccaatg tcgagagaca aggggggttc atgcacttgt ccaatgtcga gagacaaggg 120
ggttcaatgc acttgtccaa tctcgagaga caaggggggt caatgcactt gtccaatgtc 180
40 gagagacaag ggggttcaat gcacttgtcc aatgtcgact ctaga 225

45
<210> 60
<211> 619
<212> DNA
<213> Artificial Sequence
<220>
<221> misc feature
<223> Novel Sequence
50 <400> 60

cgttacataa cttacggtaa atggcccgcc tggctgaccg cccaacgacc cccgcccatt 60
gacgtcaata atgacgtatg ttcccatagt aacgccataa gggactttcc attgacgtca 120
55 atgggtggag tatttacggt aaactgccca cttggcagta catcaagtgt atcatatgcc 180

EP 1 266 015 B1

aagtacgccc cctattgacg tcaatgacgg taaatggccc gcctggcatt atgccagta 240
 catgacctta tgggactttc ctacttggca gtacatctac gtattagtca tcgctattac 300
 5 catggtgatg cgggttttggc agtacatcaa tgggcgtgga tagcggtttg actcacgggg 360
 atttccaagt ctccacccca ttgacgtcaa tgggagtttg ttttggcacc aaaatcaacg 420
 ggactttcca aaatgtcgt acaactccgc ccattgacg caaatgggcg gtaggcgtgt 480
 10 acggtgggag gtctatataa gcagagctcg tttagtgaac cgtcagatcg cctggagacg 540
 ccatccacgc tgttttgacc tccatagaag acaccgggac cgatccagcc tccgcggccg 600
 ggaacggtgc attggaacg 619

15
 <210> 61
 <211> 262
 <212> DNA
 <213> Artificial Sequence
 20 <220>
 <221> misc_feature
 <223> Novel Sequence
 <400> 61

25 atgtagtctt atgcaatact cttgtagtct tgcaacatgg taacgatgag ttagcaacat 60
 gccttacaag gagagaaaaa gcaccgtgca tgccgatagg tggaagtaag gtggtacgat 120
 cgtgccttat taggaaggca acagacgggt ctgacatgga ttggacgaac cactgaattc 180
 30 cgcatcag agatattgta tttaagtgcc tagctcgata caataaacgc catttgacca 240
 ttcaccacat tggagtgcac ct 262

35 <210> 62
 <211> 1247
 <212> DNA
 <213> Artificial Sequence
 <220>
 <221> misc feature
 40 <223> Novel Sequence
 <400> 62

tctatttcct caggccgtga ggaactgtcg ccagcttcaa gtataaatgg gtgcagtaca 60
 45 gatggcgagg cagcagctca gaagaagggc cctgogcccc gtcagcaaga ggaactgtgt 120
 ctggtatgcg gggacagagc ctccggatac cactacaatg cgctcacgtg tgaagggtgt 180
 aaagggttct tcagacggag tgttaccaaa aatgcggttt atatttgtaa attcggtcac 240
 50 gcttgcgaaa tggacatgta catgcgacgg aaatgccagg agtgccgcct gaagaagtgc 300
 ttagctgtag gcatgaggcc tgagtgcgta gtacccgaga ctcagtgcgc catgaagcgg 360
 aaagagaaga aagcacagaa ggagaaggac aaactgcctg tcagcacgac gacggtggac 420

55

EP 1 266 015 B1

gaccacatgc cgccattat gcagtgtgaa cctccacctc ctgaagcagc aaggattcac 480
gaagtggccc caaggtttct ctccgacaag ctgttggaga caaaccggca gaaaaacatc 540
5 ccccgattga cagccaacca gcagttcctt atcgccaggc tcatctggta ccaggacggg 600
tacgagcagc cttctgatga agatttgaag aggattacgc agacgtggca gcaagcggac 660
gatgaaaacg aagagtctga cactcccttc cgccagatca cagagatgac taccctcacg 720
10 gtccaactta tcgtggagtt cgcgaaggga ttgccagggt tcgccaagat ctgcgagcct 780
gatcaaatta cgctgcttaa ggcttgetca agtgaggtaa tgatgctccg agtcgcgcga 840
cgatacgatg cggcctcaga cagtgttctg ttcgcgaaca accaagcgta cactcgcgac 900
15 aactaccgca aggtctggcat ggcctacgtc atcgaggatc tactgcactt ctgccggtgc 960
atgtactcta tggcgttgga caacatccat tacgcgctgc tcacggctgt cgtcatcttt 1020
tctgaccggc cagggttgga gcagccgcaa ctggtggaag aaatccagcg gtactacctg 1080
20 aatacgctcc gcatctatat cctgaaccag ctgagcgggt cggcgcgctc gtccgctcata 1140
tacggcaaga tcctctcaat cctctctgag ctacgcacgc tcggcatgca aaactccaac 1200
atgtgcatct ccctcaagct caagaacaga aagctgccgc ctttcct 1247

25
<210> 63
<211> 440
<212> PRT
<213> Artificial Sequence
30 <220>
<221> misc feature
<223> Novel Sequence
<400> 63

35
Ser Ile Ser Ser Gly Arg Glu Glu Leu Ser Pro Ala Ser Ser Ile Asn
1 5 10 15
Gly Cys Ser Thr Asp Gly Glu Ala Arg Arg Gln Lys Lys Gly Pro Ala
20 25 30
40 Pro Arg Gln Gln Glu Glu Leu Cys Leu Val Cys Gly Asp Arg Ala Ser
35 40 45
Gly Tyr His Tyr Asn Ala Leu Thr Cys Glu Gly Cys Lys Gly Phe Phe
50 55 60
45 Arg Arg Ser Val Thr Lys Asn Ala Val Tyr Ile Cys Lys Phe Gly His
65 70 75 80
Ala Cys Glu Met Asp Met Tyr Met Arg Arg Lys Cys Gln Glu Cys Arg
85 90 95
50 Leu Lys Lys Cys Leu Ala Val Gly Met Arg Pro Glu Cys Val Val Pro
100 105 110
Glu Thr Gln Cys Ala Met Lys Arg Lys Glu Lys Lys Ala Gln Lys Glu
115 120 125
55

EP 1 266 015 B1

	Lys	Asp	Lys	Leu	Pro	Val	Ser	Thr	Thr	Thr	Val	Asp	Asp	His	Met	Pro
	130						135					140				
5	Pro	Ile	Met	Gln	Cys	Glu	Pro	Pro	Pro	Pro	Glu	Ala	Ala	Arg	Ile	His
	145					150					155					160
	Glu	Val	Val	Pro	Arg	Phe	Leu	Ser	Asp	Lys	Leu	Leu	Glu	Thr	Asn	Arg
					165					170					175	
10	Gln	Lys	Asn	Ile	Pro	Gln	Leu	Thr	Ala	Asn	Gln	Gln	Phe	Leu	Ile	Ala
				180					185					190		
	Arg	Leu	Ile	Trp	Tyr	Gln	Asp	Gly	Tyr	Glu	Gln	Pro	Ser	Asp	Glu	Asp
			195					200					205			
15	Leu	Lys	Arg	Ile	Thr	Gln	Thr	Trp	Gln	Gln	Ala	Asp	Asp	Glu	Asn	Glu
	210						215					220				
	Glu	Ser	Asp	Thr	Pro	Phe	Arg	Gln	Ile	Thr	Glu	Met	Thr	Ile	Leu	Thr
	225					230					235					240
20	Val	Gln	Leu	Ile	Val	Glu	Phe	Ala	Lys	Gly	Leu	Pro	Gly	Phe	Ala	Lys
					245					250					255	
	Ile	Ser	Gln	Pro	Asp	Gln	Ile	Thr	Leu	Leu	Lys	Ala	Cys	Ser	Ser	Glu
				260					265					270		
25	Val	Met	Met	Leu	Arg	Val	Ala	Arg	Arg	Tyr	Asp	Ala	Ala	Ser	Asp	Ser
			275					280					285			
	Val	Leu	Phe	Ala	Asn	Asn	Gln	Ala	Tyr	Thr	Arg	Asp	Asn	Tyr	Arg	Lys
	290						295					300				
30	Ala	Gly	Met	Ala	Tyr	Val	Ile	Glu	Asp	Leu	Leu	His	Phe	Cys	Arg	Cys
	305					310					315					320
	Met	Tyr	Ser	Met	Ala	Leu	Asp	Asn	Ile	His	Tyr	Ala	Leu	Leu	Thr	Ala
					325					330					335	
35	Val	Val	Ile	Phe	Ser	Asp	Arg	Pro	Gly	Leu	Glu	Gln	Pro	Gln	Leu	Val
				340					345					350		
	Glu	Glu	Ile	Gln	Arg	Tyr	Tyr	Leu	Asn	Thr	Leu	Arg	Ile	Tyr	Ile	Leu
			355					360					365			
40	Asn	Gln	Leu	Ser	Gly	Ser	Ala	Arg	Ser	Ser	Val	Ile	Tyr	Gly	Lys	Ile
	370						375					380				
	Leu	Ser	Ile	Leu	Ser	Glu	Leu	Arg	Thr	Leu	Gly	Met	Gln	Asn	Ser	Asn
	385					390					395					400
45	Met	Cys	Ile	Ser	Leu	Lys	Leu	Lys	Asn	Arg	Lys	Leu	Pro	Pro	Phe	Leu
					405					410					415	
	Glu	Glu	Ile	Trp	Asp	Val	Ala	Asp	Met	Ser	His	Thr	Gln	Pro	Pro	Pro
				420					425					430		
50	Ile	Leu	Glu	Ser	Pro	Thr	Asn	Leu								
			435					440								

<210> 64
 <211> 943
 <212> DNA
 <213> Artificial Sequence
 <220>
 <221> misc feature
 <223> Novel Sequence
 <400> 64

atgacttcga aagtttatga tccagaacaa aggaaacgga tgataactgg tccgcagtgg 60
 tgggccagat gtaaacaat gaatgttctt gattcattta ttaattatta tgattcagaa 120
 5 aaacatgcag aaaatgctgt tattttttta catggtaacg cggcctcttc ttatttatgg 180
 cgacatgttg tgccacatat tgagccagta gcgcggtgta ttataccaga ccttattggt 240
 atgggcaaat caggcaaadc tggtaatggt tcttataggt tacttgatca ttacaaatat 300
 10 cttactgcat ggtttgaact tcttaattta ccaaagaaga tcatttttgt cggccatgat 360
 tggggtgctt gtttggcatt tcattatagc tatgagcatc aagataagat caaagcaata 420
 gttcacgctg aaagtgtagt agatgtgatt gaatcatggg atgaatggcc tgatattgaa 480
 15 gaagatattg cgttgatcaa atctgaagaa ggagaaaaaa tggttttgga gaataacttc 540
 ttcgtggaaa ccatgttgcc atcaaaaadc atgagaaagt tagaaccaga agaatttgca 600
 gcatactctg aaccattcaa agagaaaggt gaagttcgtc gtccaacatt atcatggcct 660
 cgtgaaatcc cgtagtaaa aggtggtaaa cctgacgttg taaaaattgt taggaattat 720
 20 aatgcttadc tacgtgcaag tgatgattta caaaaaatgt ttattgaadc ggaccaggga 780
 ttcttttcca atgctattgt tgaaggtgcc aagaagtttc ctaatactga atttgcaaaa 840
 gtaaaaggtc ttcatttttc gcaagaagat gcacctgatg aaatgggaaa atatataaaa 900
 25 tcgttcgttg agcgagttct caaaaatgaa caataattct aga 943

30 Claims

1. A gene expression modulation system comprising:

- 35 a) a first gene expression cassette that is capable of being expressed in a host cell comprising a polynucleotide sequence that encodes a first polypeptide comprising:
- i) a DNA-binding domain that recognizes a response element associated with a gene whose expression is to be modulated;
 - 40 ii) a ligand binding domain comprising a ligand binding domain from an ecdysone receptor polypeptide;
- b) a second gene expression cassette that is capable of being expressed in the host cell comprising a polynucleotide sequence that encodes a second polypeptide comprising:
- i) a transactivation domain; and
 - 45 ii) a ligand binding domain comprising a ligand binding domain from a truncated retinoid X receptor comprising a deletion of at least the A and B domains;

50 wherein the transactivation domain is from a nuclear receptor other than an ecdysone receptor, a retinoid X receptor, or an ultraspiracle receptor, and wherein the ligand binding domains from the first polypeptide and the second polypeptide are different and dimerize.

2. The gene expression modulation system according to claim 1, further comprising a third gene expression cassette comprising:

- 55 i) a response element to which the DNA-binding domain of the polypeptide binds;
- ii) a promoter that is activated by the transactivation domain of the second polypeptide; and
 - iii) the gene whose expression is to be modulated.

3. The gene expression modulation system according to claim 1, wherein the ligand binding domain of the first polypeptide is encoded by a polynucleotide comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, and SEQ ID NO: 10.
4. The gene expression modulation system according to claim 1, wherein the ligand binding domain of the first polypeptide comprises an amino acid sequence selected from the group consisting of SEQ ID NO: 11, SEQ ID NO: 12, SEQ ID NO: 13, SEQ ID NO: 14, SEQ ID NO: 15, SEQ ID NO: 16, SEQ ID NO: 17, SEQ ID NO: 18, SEQ ID NO: 19, and SEQ ID NO: 20.
5. The gene expression modulation system according to claim 1, wherein the ligand binding domain of the second polypeptide is encoded by a polynucleotide comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO: 21, SEQ ID NO: 22, SEQ ID NO: 23, SEQ ID NO: 24, SEQ ID NO: 25, SEQ ID NO: 26, SEQ ID NO: 27, SEQ ID NO: 28, SEQ ID NO: 29, and SEQ ID NO: 30.
6. The gene expression modulation system according to claim 1, wherein the ligand binding domain of the second polypeptide comprises an amino acid sequence selected from the group consisting of SEQ ID NO: 31, SEQ ID NO: 32, SEQ ID NO: 33, SEQ ID NO: 34, SEQ ID NO: 35, SEQ ID NO: 36, SEQ ID NO: 37, SEQ ID NO: 38, SEQ ID NO: 39, and SEQ ID NO: 40.
7. A gene expression modulation system comprising:
 - a) a first gene expression cassette that is capable of being expressed in a host cell comprising a polynucleotide sequence that encodes a first polypeptide comprising:
 - i) a DNA-binding domain that recognizes a response element associated with a gene whose expression is to be modulated; and
 - ii) a ligand binding domain comprising a ligand binding domain from a truncated retinoid X receptor comprising a deletion of at least the A and B domains; and
 - b) a second gene expression cassette that is capable of being expressed in a host cell comprising a polynucleotide sequence that encodes a second polypeptide comprising:
 - i) a transactivation domain; and
 - ii) a ligand binding domain comprising a ligand binding domain from an ecdysone receptor;

wherein the ligand binding domains from the first polypeptide and the second polypeptide are different and dimerize.
8. The gene expression modulation system according to claim 7, further comprising a third gene expression cassette comprising:
 - i) a response element to which the DNA-binding domain of the first polypeptide binds;
 - ii) a promoter that is activated by the transactivation domain of the second polypeptide; and
 - iii) the gene whose expression is being modulated.
9. The gene expression modulation system according to claim 7, wherein the ligand binding domain of the first polypeptide is encoded by a polynucleotide comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO: 21, SEQ ID NO: 22, SEQ ID NO: 23, SEQ ID NO: 24, SEQ ID NO: 25, SEQ ID NO: 26, SEQ ID NO: 27, SEQ ID NO: 28, SEQ ID NO: 29, and SEQ ID NO: 30.
10. The gene expression modulation system according to claim 7, wherein the ligand binding domain of the first polypeptide comprises an amino acid sequence selected from the group consisting of SEQ ID NO: 31, SEQ ID NO: 32, SEQ ID NO: 33, SEQ ID NO: 34, SEQ ID NO: 35, SEQ ID NO: 36, SEQ ID NO: 37, SEQ ID NO: 38, SEQ ID NO: 39, and SEQ ID NO: 40.
11. The gene expression modulation system according to claim 7, wherein the ligand binding domain of the second polypeptide is encoded by a polynucleotide comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, and SEQ ID NO: 10.

ID NO: 8, SEQ ID NO: 9, and SEQ ID NO: 10.

12. The gene expression modulation system according to claim 7, wherein the ligand binding domain of the second polypeptide comprises an amino acid sequence selected from the group consisting of SEQ ID No 11, SEQ ID NO: 12, SEQ ID NO: 13, SEQ ID NO: 14, SEQ ID NO: 15, SEQ ID NO: 16, SEQ ID NO: 17, SEQ ID NO: 18, SEQ ID NO: 19, and SEQ ID NO: 20.

13. A gene expression cassette comprising a polynucleotide encoding a hybrid polypeptide selected from the group consisting of:

- a) a DNA-binding domain and an ecdysone receptor ligand binding domain, wherein the DNA binding domain is from a nuclear receptor other than an ecdysone receptor,
- b) a DNA-binding domain and a ligand binding domain from a truncated retinoid X receptor comprising a deletion of at least the A and B domains, wherein the DNA-binding domain is from a nuclear receptor other than a retinoid X receptor,
- c) a transactivation domain and an ecdysone receptor ligand binding domain, wherein the transactivation domain is from a nuclear receptor other than an ecdysone receptor, and
- d) a transactivation domain and a ligand binding domain from a truncated retinoid X receptor comprising a deletion of at least the A and B domains, wherein the transactivation domain is from a nuclear receptor other than a retinoid X receptor.

14. The gene expression cassette according to claim 13, wherein said expression cassette comprises a polynucleotide encoding a hybrid polypeptide comprising:

- a DNA-binding domain and an ecdysone receptor ligand binding domain wherein the DNA binding domain is from a nuclear receptor other than an ecdysone receptor; or
- a DNA-binding domain and a retinoid X receptor ligand binding domain, wherein the DNA-binding domain is from a nuclear receptor other than a retinoid X receptor; and

wherein the DNA-binding domain is GAL4 DNA-binding domain or a LexA DNA-binding domain.

15. The gene expression cassette according to claim 13, wherein said expression cassette comprises a polynucleotide encoding a hybrid polypeptide comprising:

- a transactivation domain and an ecdysone receptor ligand binding domain, wherein the transactivation domain is from a nuclear receptor other than an ecdysone receptor;
- or a transactivation domain and a retinoid X receptor ligand binding domain, wherein the transactivation domain is from a nuclear receptor other than a retinoid X receptor;

and wherein the transactivation domain is a VP16 transactivation domain.

16. A gene expression cassette comprising a polynucleotide encoding a hybrid polypeptide comprising a DNA-binding domain encoded by a polynucleotide comprising a nucleic acid sequence selected from the group consisting of GAL4 DBD (SEQ ID NO: 41) or a LexA DBD (SEQ ID NO: 43) and an ecdysone receptor ligand binding domain encoded by a polynucleotide comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, and SEQ ID NO: 10.

17. A gene expression cassette comprising a polynucleotide encoding a hybrid polypeptide comprising a DNA-binding domain comprising an amino acid sequence selected from the group consisting of a GAL4 DBD (SEQ ID NO: 42) or a LexA DBD (SEQ ID NO: 44) and an ecdysone receptor ligand binding domain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 11, SEQ ID NO: 12, SEQ ID NO: 13, SEQ ID NO: 14, SEQ ID NO: 15, SEQ ID NO: 16, SEQ ID NO: 17, SEQ ID NO: 18, SEQ ID NO: 19, and SEQ ID NO: 20.

18. A gene expression cassette comprising a polynucleotide encoding a hybrid polypeptide comprising a DNA-binding domain encoded by a polynucleotide comprising a nucleic acid sequence selected from the group consisting of a GAL4 DBD (SEQ ID NO: 41) or a LexA DBD (SEQ ID NO: 43) and a retinoid X receptor ligand binding domain encoded by a polynucleotide comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO:

21, SEQ ID NO: 22, SEQ ID NO: 23, SEQ ID NO: 24, SEQ ID NO: 25, SEQ ID NO: 26, SEQ ID NO: 27, SEQ ID NO: 28, SEQ ID NO: 29, and SEQ ID NO: 30.

19. A gene expression cassette comprising a polynucleotide encoding a hybrid polypeptide comprising a DNA-binding domain comprising an amino acid sequence selected from the group consisting of a GAL4 DBD (SEQ ID NO: 42) or a LexA DBD (SEQ ID NO: 44) and a retinoid X receptor ligand binding domain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 31, SEQ ID NO: 32, SEQ ID NO: 33, SEQ ID NO: 34, SEQ ID NO: 35, SEQ ID NO: 36, SEQ ID NO 37, SEQ ID NO 38, SEQ ID NO: 39, and SEQ ID NO: 40.

20. A gene expression cassette comprising a polynucleotide encoding a hybrid polypeptide comprising a transactivation domain encoded by a polynucleotide comprising a nucleic acid sequence of SEQ ID NO: 45 and an ecdysone receptor ligand binding domain encoded by a polynucleotide comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, and SEQ ID NO: 10.

21. A gene expression cassette comprising a polynucleotide encoding a hybrid polypeptide comprising a transactivation domain comprising an amino acid sequence of SEQ ID NO: 46 and an ecdysone receptor ligand binding domain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 11, SEQ ID NO: 12, SEQ ID NO: 13, SEQ ID NO: 14, SEQ ID NO: 15, SEQ ID NO: 16, SEQ ID NO: 17, SEQ ID NO: 18, SEQ ID NO: 19, and SEQ ID NO: 20.

22. A gene expression cassette comprising a polynucleotide encoding a hybrid polypeptide comprising a transactivation domain encoded by a polynucleotide comprising a nucleic acid sequence of SEQ ID NO: 45 and a retinoid X receptor ligand binding domain encoded by a polynucleotide comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO: 21, SEQ ID NO: 22, SEQ ID NO: 23, SEQ ID NO: 24, SEQ ID NO: 25, SEQ ID NO: 26, SEQ ID NO: 27, SEQ ID NO: 28, SEQ ID NO: 29, and SEQ ID NO: 30.

23. A gene expression cassette comprising a polynucleotide encoding a hybrid polypeptide comprising a transactivation domain comprising an amino acid sequence of SEQ ID NO: 46 and a retinoid X receptor ligand binding domain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 31, SEQ ID NO: 32, SEQ ID NO: 33, SEQ ID NO: 34, SEQ ID NO: 35, SEQ ID NO: 36, SEQ ID NO 37, SEQ ID NO 38, SEQ ID NO: 39, and SEQ ID NO: 40.

24. An isolated polynucleotide encoding a truncated retinoid X receptor polypeptide wherein the polynucleotide comprises a nucleic acid sequence selected from the group consisting of SEQ ID NO: 21, SEQ ID NO: 22, SEQ ID NO: 23, SEQ ID NO: 24, SEQ ID NO: 25, SEQ ID NO: 26, SEQ ID NO: 27, SEQ ID NO: 28, SEQ ID NO: 29, and SEQ ID NO: 30.

25. An isolated truncated retinoid X receptor polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 31, SEQ ID NO: 32, SEQ ID NO: 33, SEQ ID NO: 34, SEQ ID NO: 35, SEQ ID NO: 36, SEQ ID NO 37, SEQ ID NO 38, SEQ ID NO: 39, and SEQ ID NO: 40.

26. An isolated polynucleotide comprising a polynucleotide sequence which encodes the isolated polypeptide according to claim 25.

27. A method of modulating the expression of a gene in a host cell comprising the gene to be modulated comprising the steps of:


- a) introducing into the host cell the gene expression modulation system according to claim 1 or 2; and
- b) introducing into the host cell a ligand that independently combines with the ligand binding domains of the first polypeptide and the second polypeptide;

wherein the gene to be expressed is a component of a chimeric gene comprising:

- i) a response element to which the DNA binding domain from the first polypeptide binds;
- ii) a promoter that is activated by the transactivation domain of the second polypeptide; and
- iii) a gene whose expression is to be modulated.

whereby a complex is formed comprising the ligand, the first polypeptide, and the second polypeptide, and whereby the complex modulates expression of the gene in the host cell.

28. The method according to claim 27, wherein the ligand is a compound of the formula:



wherein:

E is a (C₄-C₆)alkyl containing a tertiary carbon or a cyano(C₃-C₅)alkyl containing a tertiary carbon:

R¹ is H, Me, Et, i-Pr, F, formyl, CF₃, CHF₂, CHCl₂, CH₂F, CH₂Cl, CH₂OH, CH₂OMe, CH₂CN, CN, <CCH>, 1-propynyl, 2-propynyl, vinyl, OH, OMe, OEt, cyclopropyl, CF₂CF₃, CH=CHCN, allyl, azido, SCN, or SCHF₂; R₂ is H, Me, Et, n-Pr, i-Pr, formyl, CF₃, CHF₂, CHCl₂, CH₂F, CH₂Cl, CH₂OH, CH₂OMe, CH₂CN, CN, <CCH>, 1-propynyl, 2-propynyl, vinyl, Ac, F, Cl, OH, OMe, OEt, O-n-Pr, OAc, NMe₂, NEt₂, SMe, SEt, SOCF₃, OCF₂CF₂H, COEt, cyclopropyl, CF₂CF₃, CH=CHCN, allyl, azido, OCF₃, OCHF₂, O-i-Pr, SCN, SCHF₂, SMe, NH-CN, or joined with R³ and the phenyl carbons to which R² and R³ are attached to form an ethylenedioxy, a dihydrofuryl ring with the oxygen adjacent to a phenyl carbon, or a dihydropyryl ring with the oxygen adjacent to a phenyl carbon;

R³ is H, ET, or joined with R² and the phenyl carbons to which R² and R³ are attached to form an ethylenedioxy, a dihydrofuryl ring with the oxygen adjacent to a phenyl carbon, or a dihydropyryl ring with the oxygen adjacent to a phenyl carbon;

R⁴, R⁵, and R⁶ are independently H, Me, Et, F, Cl, Br, formyl, CF₃, CHF₂, CHCl₂, CH₂F, CH₂Cl, CH₂OH, CN, C^oCH, 1-propynyl, 2-propynyl, vinyl, OMe, OEt, SMe, or SEt..

29. An isolated host cell into which the gene expression modulation system according to claim 1 or 7 has been introduced.

30. The isolated host cell according to claim 29, wherein the host cell is selected from the group consisting of a bacterial cell, a fungal cell, a yeast cell, a plant cell, an animal cell, and a mammalian cell.

31. The isolated host cell according to claim 30, wherein the mammalian cell is a murine cell or a human cell.

32. A non-human organism comprising a host cell into which the gene expression modulation system according to claim 1 or 7 has been introduced.

33. The non-human organism according to claim 32, wherein the non-human organism is selected from the group consisting of a bacterium, a fungus, a yeast, a plant, an animal, and a mammal.

34. The non-human organism according to claim 33, wherein the mammal is selected from the group consisting of a mouse, a rat, a rabbit, a cat, a dog, a bovine, a goat, a pig, a horse, a sheep, a monkey, and a chimpanzee.

Patentansprüche

1. Die Genexpression modulierendes System, welches folgendes umfaßt:

a) eine erste Genexpressionkassette, die in einer Wirtszelle exprimiert werden kann, umfassend eine Polynukleotidsequenz, die ein erstes Polypeptid codiert, welches folgendes umfaßt:

- i) eine DNA-Bindungsdomäne, die ein Antwortelement, das mit einem Gen, dessen Expression moduliert werden soll, assoziiert ist, erkennt,
- ii) eine Ligandenbindungsdomäne, die eine Ligandenbindungsdomäne eines Ecdysonrezeptorpolypeptids umfaßt,

5

b) eine zweite Genexpressionskassette, die in der Wirtszelle exprimiert werden kann, umfassend eine Polynukleotidsequenz, die ein zweites Polypeptid codiert, welches folgendes umfaßt:

10

- i) eine Transaktivierungsdomäne und
- ii) eine Ligandenbindungsdomäne, die eine Ligandenbindungsdomäne eines verkürzten Retinoid X-Rezeptors, der eine Deletion wenigstens der A- und der B-Domäne aufweist, umfaßt,

15

wobei die Transaktivierungsdomäne von einem Kernrezeptor stammt, der kein Ecdysonrezeptor, kein Retinoid X-Rezeptor oder kein Ultraspiracle-Rezeptor ist, und wobei die Ligandenbindungsdomänen des ersten Polypeptids und des zweiten Polypeptids verschieden sind und dimerisieren.

2. Die Genexpression modulierendes System nach Anspruch 1, welches weiterhin eine dritte Genexpressionskassette umfaßt, die folgendes umfaßt:

20

- i) ein Antwortelement, an das die DNA-Bindungsdomäne des Polypeptids bindet,
- ii) einen Promotor, der durch die Transaktivierungsdomäne des zweiten Polypeptids aktiviert wird, und
- iii) das Gen, dessen Expression moduliert werden soll.

25

3. Die Genexpression modulierendes System nach Anspruch 1, wobei die Ligandenbindungsdomäne des ersten Polypeptids durch ein Polynukleotid codiert wird, welches eine Nukleinsäuresequenz umfaßt, die aus der Gruppe ausgewählt ist, bestehend aus SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9 und SEQ ID NO:10.

30

4. Die Genexpression modulierendes System nach Anspruch 1, wobei die Ligandenbindungsdomäne des ersten Polypeptids eine Aminosäuresequenz umfaßt, die aus der Gruppe ausgewählt ist, bestehend aus SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19 und SEQ ID NO:20.

35

5. Die Genexpression modulierendes System nach Anspruch 1, wobei die Ligandenbindungsdomäne des zweiten Polypeptids durch ein Polynukleotid codiert wird, welches eine Nukleinsäuresequenz umfaßt, die aus der Gruppe ausgewählt ist, bestehend aus SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29 und SEQ ID NO:30.

40

6. Die Genexpression modulierendes System nach Anspruch 1, wobei die Ligandenbindungsdomäne des zweiten Polypeptids eine Aminosäuresequenz umfaßt, die aus der Gruppe ausgewählt ist, bestehend aus SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:36, SEQ ID NO:37, SEQ ID NO:38, SEQ ID NO:39 und SEQ ID NO:40.

45

7. Die Genexpression modulierendes System, welches folgendes umfaßt:

a) eine erste Genexpressionskassette, die in einer Wirtszelle exprimiert werden kann, umfassend eine Polynukleotidsequenz, welche ein erstes Polypeptid codiert, das folgendes umfaßt:

50

- i) eine DNA-Bindungsdomäne, die ein Antwortelement, das mit einem Gen, dessen Expression moduliert werden soll, assoziiert ist, erkennt, und
- ii) eine Ligandenbindungsdomäne, die eine Ligandenbindungsdomäne eines verkürzten Retinoid X-Rezeptors, der eine Deletion wenigstens der A- und der B-Domäne aufweist, umfaßt und

55

b) eine zweite Genexpressionskassette, die in einer Wirtszelle exprimiert werden kann, umfassend eine Polynukleotidsequenz, die ein zweites Polypeptid codiert, das folgendes umfaßt:

- i) eine Transaktivierungsdomäne und
- ii) eine Ligandenbindungsdomäne, die eine Ligandenbindungsdomäne eines Ecdysonrezeptors umfaßt,

wobei die Ligandenbindungsdomänen des ersten Polypeptids und des zweiten Polypeptids verschieden sind und dimerisieren.

8. Die Genexpression modulierendes System nach Anspruch 7, welches weiterhin eine dritte Genexpressionkassette umfaßt, die folgendes umfaßt:

- i) ein Antwortelement, an das die DNA-Bindungsdomäne des ersten Polypeptids bindet,
- ii) einen Promotor, der durch die Transaktivierungsdomäne des zweiten Polypeptids aktiviert wird, und
- iii) das Gen, dessen Expression moduliert werden soll.

9. Die Genexpression modulierendes System nach Anspruch 7, wobei die Ligandenbindungsdomäne des ersten Polypeptids durch ein Polynukleotid codiert wird, welches eine Nukleinsäuresequenz umfaßt, die aus der Gruppe ausgewählt ist, bestehend aus SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29 und SEQ ID NO:30.

10. Die Genexpression modulierendes System nach Anspruch 7, wobei die Ligandenbindungsdomäne des ersten Polypeptids eine Aminosäuresequenz umfaßt, die aus der Gruppe ausgewählt ist, bestehend aus SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:36, SEQ ID NO:37, SEQ ID NO:38, SEQ ID NO:39 und SEQ ID NO:40.

11. Die Genexpression modulierendes System nach Anspruch 7, wobei die Ligandenbindungsdomäne des zweiten Polypeptids durch ein Polynukleotid codiert wird, welches eine Nukleinsäuresequenz umfaßt, die aus der Gruppe ausgewählt ist, bestehend aus SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9 und SEQ ID NO:10.

12. Die Genexpression modulierendes System nach Anspruch 7, wobei die Ligandenbindungsdomäne des zweiten Polypeptids eine Aminosäuresequenz umfaßt, die aus der Gruppe ausgewählt ist, bestehend aus SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19 und SEQ ID NO:20.

13. Genexpressionskassette, welche ein Polynukleotid umfaßt, das ein hybrides Polypeptid codiert, ausgewählt aus der Gruppe, bestehend aus:

- a) einer DNA-Bindungsdomäne und einer Ecdysonrezeptor-Ligandenbindungsdomäne, wobei die DNA-Bindungsdomäne von einem Kernrezeptor stammt, der kein Ecdysonrezeptor ist,
- b) einer DNA-Bindungsdomäne und einer Ligandenbindungsdomäne eines verkürzten Retinoid X-Rezeptors, welcher eine Deletion wenigstens der A- und der B-Domäne aufweist, wobei die DNA-Bindungsdomäne von einem Kernrezeptor stammt, der kein Retinoid X-Rezeptor ist,
- c) einer Transaktivierungsdomäne und einer Ecdysonrezeptor-Ligandenbindungsdomäne, wobei die Transaktivierungsdomäne von einem Kernrezeptor stammt, der kein Ecdysonrezeptor ist, und
- d) einer Transaktivierungsdomäne und einer Ligandenbindungsdomäne eines verkürzten Retinoid X-Rezeptors, welcher eine Deletion wenigstens der A- und der B-Domäne aufweist, wobei die Transaktivierungsdomäne von einem Kernrezeptor stammt, der kein Retinoid X-Rezeptor ist.

14. Genexpressionskassette nach Anspruch 13, wobei die Expressionskassette ein Polynukleotid umfaßt, welches ein hybrides Polypeptid codiert, das folgendes aufweist:

- eine DNA-Bindungsdomäne und eine Ecdysonrezeptor-Ligandenbindungsdomäne, wobei die DNA-Bindungsdomäne von einem Kernrezeptor stammt, der kein Ecdysonrezeptor ist,
- eine DNA-Bindungsdomäne und eine Retinoid X-Rezeptor-Ligandenbindungsdomäne, wobei die DNA-Bindungsdomäne von einem Kernrezeptor stammt, der kein Retinoid X-Rezeptor ist, und

wobei die DNA-Bindungsdomäne eine GAL4-DNA-Bindungsdomäne oder eine LexA-DNA-Bindungsdomäne ist.

15. Genexpressionskassette nach Anspruch 13, wobei die Expressionskassette ein Polynukleotid umfaßt, welches ein hybrides Polypeptid codiert, das folgendes aufweist:

- eine Transaktivierungsdomäne und eine Ecdysonrezeptor-Ligandenbindungsdomäne, wobei die Transaktivie-

rungsdomäne von einem Kernrezeptor stammt, der kein Ecdysonrezeptor ist, oder eine Transaktivierungsdomäne und eine Retinoid X-Rezeptor-Ligandenbindungsdomäne, wobei die Transaktivierungsdomäne von einem Kernrezeptor stammt, der kein Retinoid X-Rezeptor ist,

und wobei die Transaktivierungsdomäne eine VP16-Transaktivierungsdomäne ist.

16. Genexpressionskassette, welche ein Polynukleotid umfaßt, das ein hybrides Polypeptid codiert, welches eine DNA-Bindungsdomäne umfaßt, die durch ein Polynukleotid codiert ist, welches eine Nukleinsäuresequenz umfaßt, die aus der Gruppe ausgewählt ist, bestehend aus GAL4 DBD (SEQ ID NO:41) oder LexA DBD (SEQ ID NO:43), und eine Ecdysonrezeptor-Ligandenbindungsdomäne umfaßt, die durch ein Polynukleotid codiert ist, welches eine Nukleinsäuresequenz umfaßt, die aus der Gruppe ausgewählt ist, bestehend aus SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9 und SEQ ID NO:10.
17. Genexpressionskassette, welche ein Polynukleotid umfaßt, das ein hybrides Polypeptid codiert, welches eine DNA-Bindungsdomäne umfaßt, die eine Aminosäuresequenz umfaßt, ausgewählt aus der Gruppe, bestehend aus GAL4 DBD (SEQ ID NO:42) oder LexA DBD (SEQ ID NO:44), umfaßt, und eine Ecdysonrezeptor-Ligandenbindungsdomäne, die eine Aminosäuresequenz umfaßt, ausgewählt aus der Gruppe, bestehend aus SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19 und SEQ ID NO:20.
18. Genexpressionskassette, welche ein Polynukleotid umfaßt, das ein hybrides Polypeptid codiert, welches eine DNA-Bindungsdomäne umfaßt, die durch ein Polynukleotid codiert wird, welches eine Nukleinsäuresequenz umfaßt, ausgewählt aus der Gruppe, bestehend aus GAL4 DBD (SEQ ID NO:41) oder LexA DBD (SEQ ID NO:43), und eine Retinoid X-Rezeptor-Ligandenbindungsdomäne umfaßt, die durch ein Polynukleotid codiert wird, welches eine Nukleinsäuresequenz umfaßt, ausgewählt aus der Gruppe, bestehend aus SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29 und SEQ ID NO:30.
19. Genexpressionskassette, welche ein Polynukleotid umfaßt, das ein hybrides Polypeptid codiert, welches eine DNA-Bindungsdomäne umfaßt, die eine Aminosäuresequenz umfaßt, ausgewählt aus der Gruppe, bestehend aus GAL4 DBD (SEQ ID NO:42) oder LexA DBD (SEQ ID NO:44), und eine Retinoid X-Rezeptor-Ligandenbindungsdomäne umfaßt, die eine Aminosäuresequenz umfaßt, ausgewählt aus der Gruppe, bestehend aus SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:36, SEQ ID NO:37, SEQ ID NO:38, SEQ ID NO:39 und SEQ ID NO:40.
20. Genexpressionskassette, welche ein Polynukleotid umfaßt, das ein hybrides Polypeptid codiert, welches eine Transaktivierungsdomäne umfaßt, die durch ein Polynukleotid codiert wird, welches eine Nukleinsäuresequenz der SEQ ID NO:45 umfaßt, und eine Ecdysonrezeptor-Ligandenbindungsdomäne umfaßt, die durch ein Polynukleotid codiert wird, welches eine Nukleinsäuresequenz umfaßt, ausgewählt aus der Gruppe, bestehend aus SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9 und SEQ ID NO:10.
21. Genexpressionskassette, welche ein Polynukleotid umfaßt, das ein hybrides Polypeptid codiert, welches eine Transaktivierungsdomäne umfaßt, die eine Aminosäuresequenz der SEQ ID NO:46 umfaßt, und eine Ecdysonrezeptor-Ligandenbindungsdomäne umfaßt, die eine Aminosäuresequenz umfaßt, ausgewählt aus der Gruppe, bestehend aus SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19 und SEQ ID NO:20.
22. Genexpressionskassette, welche ein Polynukleotid umfaßt, das ein hybrides Polypeptid codiert, welches eine Transaktivierungsdomäne umfaßt, die durch ein Polynukleotid codiert wird, welches eine Nukleinsäuresequenz der SEQ ID NO:45 umfaßt, und eine Retinoid X-Rezeptor-Ligandenbindungsdomäne umfaßt, die durch ein Polynukleotid codiert wird, welches eine Nukleinsäuresequenz umfaßt, ausgewählt aus der Gruppe, bestehend aus SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29 und SEQ ID NO:30.
23. Genexpressionskassette, welche ein Polynukleotid umfaßt, das ein hybrides Polypeptid codiert, welches eine Transaktivierungsdomäne umfaßt, die eine Aminosäuresequenz der SEQ ID NO:46 umfaßt, und eine Retinoid X-Rezeptor-Ligandenbindungsdomäne umfaßt, die eine Aminosäuresequenz umfaßt, ausgewählt aus der Gruppe, bestehend

aus SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:36, SEQ ID NO:37, SEQ ID NO:38, SEQ ID NO:39 und SEQ ID NO:40.

24. Isoliertes Polypeptid, welches ein verkürztes Retinoid X-Rezeptorpolypeptid codiert, wobei das Polynukleotid eine Nukleinsäuresequenz umfaßt, ausgewählt aus der Gruppe, bestehend aus SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29 und SEQ ID NO:30.

25. Isoliertes verkürztes Retinoid X-Rezeptorpolypeptid, welches eine Aminosäuresequenz umfaßt, ausgewählt aus der Gruppe, bestehend aus SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:36, SEQ ID NO:37, SEQ ID NO:38, SEQ ID NO:39 und SEQ ID NO:40.

26. Isoliertes Polynukleotid, welches eine Polynukleotidsequenz umfaßt, die das isolierte Polypeptid nach Anspruch 25 codiert.

27. Verfahren zum Modulieren der Expression eines Gens in einer Wirtszelle, die das zu modulierende Gen enthält, welches die folgenden Stufen umfaßt:


- Einbringen des die Genexpression modulierenden Systems nach einem der Ansprüche 1 oder 2 in die Wirtszelle und
- Einbringen eines Liganden, der sich unabhängig mit den Ligandenbindungsdomänen des ersten Polypeptids und des zweiten Polypeptids vereinigt, in die Wirtszelle,

wobei das zu exprimierende Gen eine Komponente eines chimären Gens ist, welches folgendes umfaßt:

- ein Antwortelement, an das die DNA-Bindungsdomäne des ersten Polypeptids bindet,
- einen Promotor, der durch die Transaktivierungsdomäne des zweiten Polypeptids aktiviert wird, und
- das Gen, dessen Expression moduliert werden soll,

wobei ein Komplex gebildet wird, der den Liganden, das erste Polypeptid und das zweite Polypeptid umfaßt, und wobei der Komplex die Expression des Gens in der Wirtszelle moduliert.

28. Verfahren nach Anspruch 27, wobei der Ligand eine Verbindung der Formel



ist, wobei:

E ein (C₄-C₆)-Alkyl, das ein tertiäres Kohlenstoffatom enthält, oder ein Cyano-(C₃-C₅)-alkyl, das ein tertiäres Kohlenstoffatom enthält, ist,
 R¹ H, Me, Et, i-Pr, F, Formyl, CF₃, CHF₂, CHCl₂, CH₂F, CH₂Cl, CH₂OH, CH₂OMe, CH₂CN, CN, CCH, 1-Propinyl, 2-Propinyl, Vinyl, OH, OMe, OEt, Cyclopropyl, CF₂CF₃, CH=CHCN, Allyl, Azido, SCN oder SCHF₂ ist,
 R² H, Me, Et, n-Pr, i-Pr, Formyl, CF₃, CHF₂, CHCl₂, CH₂F, CH₂Cl, CH₂OH, CH₂OMe, CH₂CN, CN, CCH, 1-Propinyl, 2-Propinyl, Vinyl, Ac, F, Cl, OH, OMe, OEt, O-n-Pr, OAc, NMe₂, NEt₂, SMe, SET, SOCF₃, OCF₂CF₂H, COEt, Cyclopropyl, CF₂CF₃, CH=CHCN, Allyl, Azido, OCF₃, OCHF₂, O-i-Pr, SCN, SCHF₂, SOMe, NH-CN ist oder mit R³ und den Phenylkohlenstoffen, an die R² und R³ angebunden sind, unter Bildung von Ethylendioxy, eines Dihydrofurylring, bei dem der Sauerstoff zu einem Phenylkohlenstoff benachbart ist, oder eines Dihy-

dropyrylrings, bei dem der Sauerstoff zu einem Phenylkohlenstoff benachbart ist, verbunden ist, R^3 H, Et oder mit R^2 und den Phenylkohlenstoffen, an die R^2 und R^3 angebunden sind, unter Bildung von Ethylendioxy, eines Dihydrofurylrings, bei dem der Sauerstoff zu einem Phenylkohlenstoff benachbart ist, oder eines Dihydropyrylrings, bei dem der Sauerstoff zu einem Phenylkohlenstoff benachbart ist, verbunden ist, R^4 , R^5 und R^6 unabhängig voneinander H, Me, Et, F, Cl, Br, Formyl, CF_3 , CHF_2 , $CHCl_2$, CH_2F , CH_2Cl , CH_2OH , CN, $C^\circ CH$, 1-Propinyl, 2-Propinyl, Vinyl, OMe, OEt, SMe oder SEt sind.

29. Isolierte Wirtszelle, in die das die Genexpression modulierende System nach Anspruch 1 oder 7 eingebracht wurde.

30. Isolierte Wirtszelle nach Anspruch 29, wobei die Wirtszelle aus der Gruppe ausgewählt ist, bestehend aus einer Bakterienzelle, einer Pilzzelle, einer Hefezelle, einer Pflanzenzelle, einer Tierzelle und einer Säugerzelle.

31. Isolierte Wirtszelle nach Anspruch 30, wobei die Säugerzelle eine Mauszelle oder eine menschliche Zelle ist.

32. Nicht-menschlicher Organismus, welcher eine Wirtszelle, in die das die Genexpression modulierende System nach Anspruch 1 oder 7 eingebracht wurde, umfaßt.

33. Nicht-menschlicher Organismus nach Anspruch 32, wobei der nicht-menschliche Organismus aus der Gruppe ausgewählt ist, bestehend aus einem Bakterium, einem Pilz, einer Hefe, einer Pflanze, einem Tier und einem Säuger.

34. Nicht-menschlicher Organismus nach Anspruch 33, wobei der Säuger aus der Gruppe ausgewählt ist, bestehend aus einer Maus, einer Ratte, einem Kaninchen, einer Katze, einem Hund, einem Rind, einer Ziege, einem Schwein, einem Pferd, einem Schaf, einem Affen und einem Schimpansen.

Revendications

1. Système de modulation d'expression génique comprenant :

a) une première cassette d'expression génique qui est capable d'être exprimée dans une cellule hôte comprenant une séquence polynucléotidique qui code pour un premier polypeptide comprenant :

i) un domaine de liaison à l'ADN qui reconnaît un élément de réponse associé à un gène dont l'expression doit être modulée ;

ii) un domaine de liaison au ligand comprenant un domaine de liaison au ligand issu d'un polypeptide du récepteur de l'ecdysone ;

b) une deuxième cassette d'expression génique qui est capable d'être exprimée dans la cellule hôte comprenant une séquence polynucléotidique qui code pour un second polypeptide comprenant :

i) un domaine de transactivation ; et

ii) un domaine de liaison au ligand comprenant un domaine de liaison au ligand issu d'un récepteur X des rétinoïdes tronqué comprenant une délétion d'au moins les domaines A et B ;

dans lequel le domaine de transactivation provient d'un récepteur nucléaire autre qu'un récepteur de l'ecdysone, d'un récepteur X des rétinoïdes ou d'un récepteur ultraspiracle ; et dans lequel les domaines de liaison au ligand issus du premier polypeptide et du second polypeptide sont différents et se dimérisent.

2. Système de modulation d'expression génique selon la revendication 1, comprenant de plus une troisième cassette d'expression génique comprenant :

i) un élément de réponse auquel se lie le domaine de liaison à l'ADN du polypeptide ;

ii) un promoteur qui est activé par le domaine de transactivation du second polypeptide ; et

iii) le gène dont l'expression doit être modulée.

3. Système d'expression génique selon la revendication 1, dans lequel le domaine de liaison au ligand du premier polypeptide est codé par un polynucléotide comprenant une séquence d'acide nucléique choisie dans le groupe formé par SEQ ID N° 1, SEQ ID N° 2, SEQ ID N° 3, SEQ ID N° 4, SEQ ID N° 5, SEQ ID N° 6, SEQ ID N° 7, SEQ

ID N° 8, SEQ ID N° 9 et SEQ ID N° 10.

4. Système de modulation d'expression génique selon la revendication 1, dans lequel le domaine de liaison au ligand du premier polypeptide comprend une séquence d'acides aminés choisie dans le groupe formé par SEQ ID N° 11, SEQ ID N° 12, SEQ ID N° 13, SEQ ID N° 14, SEQ ID N° 15, SEQ ID N° 16, SEQ ID N° 17, SEQ ID N° 18, SEQ ID N° 19 et SEQ ID N° 20.
5. Système de modulation d'expression génique selon la revendication 1, dans lequel le domaine de liaison au ligand du second polypeptide est codé par un polynucléotide comprenant une séquence d'acide nucléique choisie dans le groupe formé par SEQ ID N° 21, SEQ ID N° 22, SEQ ID N° 23, SEQ ID N° 24, SEQ ID N° 25, SEQ ID N° 26, SEQ ID N° 27, SEQ ID N° 28, SEQ ID N° 29 et SEQ ID N° 30.
6. Système de modulation d'expression génique selon la revendication 1, dans lequel le domaine de liaison au ligand du second polypeptide comprend une séquence d'acides aminés choisie dans le groupe formé par SEQ ID N° 31, SEQ ID N° 32, SEQ ID N° 33, SEQ ID N° 34, SEQ ID N° 35, SEQ ID N° 36, SEQ ID N° 37, SEQ ID N° 38, SEQ ID N° 39 et SEQ ID N° 40.
7. Système de modulation d'expression génique comprenant :
 - a) une première cassette d'expression génique qui est capable d'être exprimée dans une cellule hôte comprenant une séquence polynucléotidique qui code pour un premier polypeptide comprenant :
 - i) un domaine de liaison à l'ADN qui reconnaît un élément de réponse associé à un gène dont l'expression doit être modulée ; et
 - ii) un domaine de liaison au ligand comprenant un domaine de liaison au ligand issu d'un récepteur X des rétinoïdes tronqué comprenant une délétion d'au moins les domaines A et B ; et
 - b) une deuxième cassette d'expression génique qui est capable d'être exprimée dans une cellule hôte comprenant une séquence polynucléotidique qui code pour un second polypeptide comprenant :
 - i) un domaine de transactivation ; et
 - ii) un domaine de liaison au ligand comprenant un domaine de liaison au ligand issu d'un récepteur de l'ecdysone ;
8. Système de modulation d'expression génique selon la revendication 7, comprenant de plus une troisième cassette d'expression génique comprenant :
 - i) un élément de réponse auquel se lie le domaine de liaison à l'ADN du premier polypeptide ;
 - ii) un promoteur qui est activé par le domaine de transactivation du second polypeptide ; et
 - iii) le gène dont l'expression est modulée.
9. Système de modulation d'expression génique selon la revendication 7, dans lequel le domaine de liaison au ligand du premier polypeptide est codé par un polynucléotide comprenant une séquence d'acide nucléique choisie dans le groupe formé par SEQ ID N° 21, SEQ ID N° 22, SEQ ID N° 23, SEQ ID N° 24, SEQ ID N° 25, SEQ ID N° 26, SEQ ID N° 27, SEQ ID N° 28, SEQ ID N° 29 et SEQ ID N° 30.
10. Système de modulation d'expression génique selon la revendication 7, dans lequel le domaine de liaison au ligand du premier polypeptide comprend une séquence d'acides aminés choisie dans le groupe formé par SEQ ID N° 31, SEQ ID N° 32, SEQ ID N° 33, SEQ ID N° 34, SEQ ID N° 35, SEQ ID N° 36, SEQ ID N° 37, SEQ ID N° 38, SEQ ID N° 39 et SEQ ID N° 40.
11. Système de modulation d'expression génique selon la revendication 7, dans lequel le domaine de liaison au ligand du second polypeptide est codé par un polynucléotide comprenant une séquence d'acide nucléique choisie dans le groupe formé par SEQ ID N° 1, SEQ ID N° 2, SEQ ID N° 3, SEQ ID N° 4, SEQ ID N° 5, SEQ ID N° 6, SEQ ID N° 7, SEQ ID N° 8, SEQ ID N° 9 et SEQ ID N° 10.

12. Système de modulation d'expression génique selon la revendication 7, dans lequel le domaine de liaison au ligand du second polypeptide comprend une séquence d'acides aminés choisie dans le groupe formé par SEQ ID N° 11, SEQ ID N° 12, SEQ ID N° 13, SEQ ID N° 14, SEQ ID N° 15, SEQ ID N° 16, SEQ ID N° 17, SEQ ID N° 18, SEQ ID N° 19 et SEQ ID N° 20.

13. Cassette d'expression génique comprenant un polynucléotide codant pour un polypeptide hybride choisi dans le groupe formé par :

a) un domaine de liaison à l'ADN et un domaine de liaison au ligand de récepteur de l'ecdysone, le domaine de liaison à l'ADN étant issu d'un récepteur nucléaire autre qu'un récepteur de l'ecdysone ;

b) un domaine de liaison à l'ADN et un domaine de liaison au ligand issu d'un récepteur X des rétinoïdes tronqué comprenant une délétion d'au moins les domaines A et B, le domaine de liaison à l'ADN étant issu d'un récepteur nucléaire autre qu'un récepteur X des rétinoïdes ;

c) un domaine de transactivation et un domaine de liaison au ligand de récepteur de l'ecdysone, le domaine de transactivation étant issu d'un récepteur nucléaire autre qu'un récepteur de l'ecdysone ; et

d) un domaine de transactivation et un domaine de liaison au ligand issu d'un récepteur X des rétinoïdes tronqué comprenant une délétion d'au moins les domaines A et B, le domaine de transactivation étant issu d'un récepteur nucléaire autre qu'un récepteur X des rétinoïdes.

14. Cassette d'expression génique selon la revendication 13, dans laquelle ladite cassette d'expression génique comprend un polynucléotide codant pour un polypeptide hybride comprenant :

un domaine de liaison à l'ADN et un domaine de liaison au ligand de récepteur de l'ecdysone, le domaine de liaison à l'ADN étant issu d'un récepteur nucléaire autre qu'un récepteur de l'ecdysone ; ou

un domaine de liaison à l'ADN et un domaine de liaison au ligand de récepteur X des rétinoïdes, le domaine de liaison à l'ADN étant issu d'un récepteur nucléaire autre qu'un récepteur X des rétinoïdes ; et

dans laquelle le domaine de liaison à l'ADN est un domaine de liaison à l'ADN de GAL4 ou un domaine de liaison à l'ADN de LexA.

15. Cassette d'expression génique selon la revendication 13, dans laquelle ladite cassette d'expression génique comprend un polynucléotide codant pour un polypeptide hybride comprenant :

un domaine de transactivation et un domaine de liaison au ligand de récepteur de l'ecdysone, le domaine de transactivation étant issu d'un récepteur nucléaire autre qu'un récepteur de l'ecdysone ;

ou un domaine de transactivation et un domaine de liaison au ligand de récepteur X des rétinoïdes, le domaine de transactivation étant issu d'un récepteur nucléaire autre qu'un récepteur X des rétinoïdes ;

et dans laquelle le domaine de transactivation est un domaine de transactivation de VP16.

16. Cassette d'expression génique comprenant un polynucléotide codant pour un polypeptide hybride comprenant un domaine de liaison à l'ADN codé par un polynucléotide comprenant une séquence d'acide nucléique choisie dans le groupe formé par un DBD de GAL4 (SEQ ID N° 41) ou un DBD de LexA (SEQ ID N° 43) et un domaine de liaison au ligand de récepteur de l'ecdysone codé par un polynucléotide comprenant une séquence d'acide nucléique choisie dans le groupe formé par SEQ ID N° 1, SEQ ID N° 2, SEQ ID N° 3, SEQ ID N° 4, SEQ ID N° 5, SEQ ID N° 6, SEQ ID N° 7, SEQ ID N° 8, SEQ ID N° 9 et SEQ ID N° 10.

17. Cassette d'expression génique comprenant un polynucléotide codant pour un polypeptide hybride comprenant un domaine de liaison à l'ADN comprenant une séquence d'acides aminés choisie dans le groupe formé par un DBD de GAL4 (SEQ ID N° 42) ou un DBD de LexA (SEQ ID N° 44) et un domaine de liaison au ligand de récepteur de l'ecdysone comprenant une séquence d'acides aminés choisie dans le groupe formé par SEQ ID N° 11, SEQ ID N° 12, SEQ ID N° 13, SEQ ID N° 14, SEQ ID N° 15, SEQ ID N° 16, SEQ ID N° 17, SEQ ID N° 18, SEQ ID N° 19 et SEQ ID N° 20.

18. Cassette d'expression génique comprenant un polynucléotide codant pour un polypeptide hybride comprenant un domaine de liaison à l'ADN codé par un polynucléotide comprenant une séquence d'acide nucléique choisie dans le groupe formé par un DBD de GAL4 (SEQ ID N° 41) ou un DBD de LexA (SEQ ID N° 43) et un domaine de liaison au ligand de récepteur X des rétinoïdes codé par un polynucléotide comprenant une séquence d'acide nucléique

choisie dans le groupe formé par SEQ ID N° 21, SEQ ID N° 22, SEQ ID N° 23, SEQ ID N° 24, SEQ ID N° 25, SEQ ID N° 26, SEQ ID N° 27, SEQ ID N° 28, SEQ ID N° 29 et SEQ ID N° 30.


- 5 19. Cassette d'expression génique comprenant un polynucléotide codant pour un polypeptide hybride comprenant un domaine de liaison à l'ADN comprenant une séquence d'acides aminés choisie dans le groupe formé par un DBD de GAL4 (SEQ ID N° 42) ou un DBD de LexA (SEQ ID N° 44) et un domaine de liaison au ligand de récepteur X des rétinoïdes comprenant une séquence d'acides aminés choisie dans le groupe formé par SEQ ID N° 31, SEQ ID N° 32, SEQ ID N° 33, SEQ ID N° 34, SEQ ID N° 35, SEQ ID N° 36, SEQ ID N° 37, SEQ ID N° 38, SEQ ID N° 39 et SEQ ID N° 40.
- 10 20. Cassette d'expression génique comprenant un polynucléotide codant pour un polypeptide hybride comprenant un domaine de transactivation codé par un polynucléotide comprenant une séquence d'acide nucléique de SEQ ID N° 45 et un domaine de liaison au ligand de récepteur de l'ecdysone codé par un polynucléotide comprenant une séquence d'acide nucléique choisie dans le groupe formé par SEQ ID N° 1, SEQ ID N° 2, SEQ ID N° 3, SEQ ID N° 4, SEQ ID N° 5, SEQ ID N° 6, SEQ ID N° 7, SEQ ID N° 8, SEQ ID N° 9 et SEQ ID N° 10.
- 15 21. Cassette d'expression génique comprenant un polynucléotide codant pour un polypeptide hybride comprenant un domaine de transactivation comprenant une séquence d'acides aminés de SEQ ID N° 46 et un domaine de liaison au ligand de récepteur de l'ecdysone comprenant une séquence d'acides aminés choisie dans le groupe formé par SEQ ID N° 11, SEQ ID N° 12, SEQ ID N° 13, SEQ ID N° 14, SEQ ID N° 15, SEQ ID N° 16, SEQ ID N° 17, SEQ ID N° 18, SEQ ID N° 19 et SEQ ID N° 20.
- 20 22. Cassette d'expression génique comprenant un polynucléotide codant pour un polypeptide hybride comprenant un domaine de transactivation codé par un polynucléotide comprenant une séquence d'acide nucléique de SEQ ID N° 45 et un domaine de liaison au ligand de récepteur X des rétinoïdes codé par un polynucléotide comprenant une séquence d'acide nucléique choisie dans le groupe formé par SEQ ID N° 21, SEQ ID N° 22, SEQ ID N° 23, SEQ ID N° 24, SEQ ID N° 25, SEQ ID N° 26, SEQ ID N° 27, SEQ ID N° 28, SEQ ID N° 29 et SEQ ID N° 30.
- 25 23. Cassette d'expression génique comprenant un polynucléotide codant pour un polypeptide hybride comprenant un domaine de transactivation comprenant une séquence d'acides aminés de SEQ ID N° 46 et un domaine de liaison au ligand de récepteur X des rétinoïdes comprenant une séquence d'acides aminés choisie dans le groupe formé par SEQ ID N° 31, SEQ ID N° 32, SEQ ID N° 33, SEQ ID N° 34, SEQ ID N° 35, SEQ ID N° 36, SEQ ID N° 37, SEQ ID N° 38, SEQ ID N° 39 et SEQ ID N° 40.
- 30 24. Polynucléotide isolé codant pour un polypeptide de récepteur X des rétinoïdes tronqué, dans lequel le polynucléotide comprend une séquence d'acide nucléique choisie dans le groupe formé par SEQ ID N° 21, SEQ ID N° 22, SEQ ID N° 23, SEQ ID N° 24, SEQ ID N° 25, SEQ ID N° 26, SEQ ID N° 27, SEQ ID N° 28, SEQ ID N° 29 et SEQ ID N° 30.
- 35 25. Polypeptide de récepteur X des rétinoïdes tronqué isolé comprenant une séquence d'acides aminés choisie dans le groupe formé par SEQ ID N° 31, SEQ ID N° 32, SEQ ID N° 33, SEQ ID N° 34, SEQ ID N° 35, SEQ ID N° 36, SEQ ID N° 37, SEQ ID N° 38, SEQ ID N° 39 et SEQ ID N° 40.
- 40 26. Polynucléotide isolé comprenant une séquence polynucléotidique qui code pour le polypeptide isolé selon la revendication 25.
- 45 27. Procédé de modulation de l'expression d'un gène dans une cellule hôte comprenant le gène à moduler, comprenant les étapes de :
 - 50 a) introduction dans la cellule hôte du système de modulation d'expression génique selon la revendication 1 ou 2 ; et
 - b) introduction dans la cellule hôte d'un ligand qui se combine indépendamment avec les domaines de liaison au ligand du premier polypeptide et du second polypeptide ;

dans lequel le gène à exprimer est un composant d'un gène chimère comprenant :

 - 55 i) un élément de réponse auquel se lie le domaine de liaison à l'ADN issu du premier polypeptide ;
 - ii) un promoteur qui est activé par le domaine de transactivation du second polypeptide ; et
 - iii) un gène dont l'expression doit être modulée,

en sorte que soit formé un complexe comprenant le ligand, le premier polypeptide et le second polypeptide, et en sorte que le complexe module l'expression du gène dans la cellule hôte.

28. Procédé selon la revendication 27, dans lequel le ligand est un composé de formule :



dans laquelle :

E est un groupe alkyle en C₄ à C₆ contenant un atome de carbone tertiaire ou un groupe cyano(alkyle en C₃ à C₅) contenant un atome de carbone tertiaire ;

R¹ est H, Me, Et, i-Pr, F, un groupe formyle, CF₃, CHF₂, CHCl₂, CH₂F, CH₂Cl, CH₂OH, CH₂OMe, CH₂CN, CN, CCH, 1-propynyle, 2-propynyle, vinyle, OH, OMe, OEt, cyclopropyle, CF₂CF₃, CH=CHCN, allyle, azido, SCN ou SCHF₂ ;

R² est H, Me, Et, n-Pr, i-Pr, un groupe formyle, CF₃, CHF₂, CHCl₂, CH₂F, CH₂Cl, CH₂OH, CH₂OMe, CH₂CN, CN, CCH, 1-propynyle, 2-propynyle, vinyle, Ac, F, Cl, OH, OMe, OEt, O-n-Pr, OAc, NMe₂, NEt₂, SMe, SEt, SOCF₃, OCF₂CF₂H, COEt, cyclopropyle, CF₂CF₃, CH=CHCN, allyle, azido, OCF₃, OCHF₂, O-i-Pr, SCN, SCHF₂, SOMe, NH-CN, ou est joint à R³ et aux atomes de carbone phényles auxquels R² et R³ sont attachés pour former un groupe éthylènedioxy, un cycle dihydrofuryle dont l'atome d'oxygène est adjacent à un atome de carbone phényle ;

R³ est H, Et, ou est joint à R² et aux atomes de carbone phényles auxquels R² et R³ sont attachés pour former un groupe éthylènedioxy, un cycle dihydrofuryle dont l'oxygène est adjacent à un atome de carbone phényle ;

R⁴, R⁵ et R⁶ sont indépendamment H, Me, Et, F, Cl, Br, un groupe formyle, CF₃, CHF₂, CHCl₂, CH₂F, CH₂Cl, CH₂OH, CN, C^oCH, 1-propynyle, 2-propynyle, vinyle, OMe, OEt, SMe ou SEt.

29. Cellule hôte isolée dans laquelle a été introduit le système de modulation d'expression génique selon la revendication 1 ou 7.

30. Cellule hôte isolée selon la revendication 29, dans laquelle la cellule hôte est choisie dans le groupe formé par une cellule bactérienne, une cellule fongique, une cellule de levure, une cellule végétale, une cellule animale et une cellule de mammifère.

31. Cellule hôte isolée selon la revendication 30, dans laquelle la cellule de mammifère est une cellule murine ou une cellule humaine.

32. Organisme non humain comprenant une cellule hôte dans laquelle a été introduit le système de modulation d'expression génique selon la revendication 1 ou 7.

33. Organisme non humain selon la revendication 32, dans lequel l'organisme non humain est choisi dans le groupe formé par une bactérie, un champignon, une levure, une plante, un animal et un mammifère.

34. Organisme non humain selon la revendication 33, dans lequel le mammifère est choisi dans le groupe formé par une souris, un rat, un lapin, un chat, un chien, un bovin, une chèvre, un porc, un cheval, un mouton, un singe et un chimpanzé.




Figure 1




Figure 2




Figure 3




Figure 4

GAL4USP 

VP16CfEcR 

pGAL4RELuc 

Figure 5

GAL4CfEcRVP16 

pGAL4RELuc 

Figure 6

VP16CfEcR 


pEcRERELuc 

Figure 7

VP16DmEcR 

RXR 

pE/GRELacZ 

Figure 8

VP16CfEcR 

RXR 

pE/GRELacZ 

Figure 9

VP16CfEcR 

pE/GRELacZ 

Figure 10

Analysis of CIEoR Truncations with MmRXRDE in 3T3 Cells




Figure 11

Analysis of CIEoR Truncations with MmRXRE in 3T3 Cells




Figure 12

Analysis of MmRXR Truncations with CfEcRCDEF in 3T3 Cells




Figure 13

Analysis of MmRXR Truncations with CfEcRCDEF in 3T3 Cells




Figure 14




Figure 15